

Learning Normal Representations for Blood Biomarkers

Aashna P. Shah^{1,2,6}, Michelle M. Li^{1,6}, Yash Lal⁴, Seffi Cohen^{1,6,7}, Liat F. Antwarg^{1,6,7}, Morgan Sanchez^{1,6}, James A. Diao^{1,3,6}, Chirag J. Patel¹, Ben Y. Reis^{1,5,6,7}, Ran D. Balicer^{6,7*}, Noa Dagan^{6,7,8*}, and Arjun K. Manrai^{1,6*}

1. Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA
2. Department of Systems Biology, Harvard Medical School, Boston, MA, USA
3. Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA
4. Department of Mathematics, Johns Hopkins University, Baltimore, MD, USA
5. Computational Health Informatics Program (CHIP), Boston Children's Hospital, Boston, MA, USA
6. The Ivan and Francesca Berkowitz Family Living Laboratory Collaboration at Harvard Medical School and Clalit Research Institute, USA and Israel
7. Clalit Research Institute, Innovation Division, Clalit Health Services, Ramat-Gan, Israel
8. Faculty of Computer and Information Science, Ben Gurion University, Be'er Sheva, Israel

* Co-senior authors

Correspondence: Arjun_Manrai@hms.harvard.edu

Abstract

Blood-based biomarkers underpin clinical diagnosis and management, yet their interpretation relies largely on fixed population reference intervals that ignore stable, intra-patient variability. As such, population-based interpretation can mask meaningful deviation from an individual's baseline, risking delayed disease detection. To remedy this, there have been increasing efforts to personalize blood biomarker interpretation using individual testing histories. However, these methods may overfit to sparse data, inflating false-positive rates and unnecessary follow-up, and can also unwittingly include unrecognized or subclinical disease. Here, we leverage nearly 2 billion longitudinal laboratory measurements from over 1.6 million individuals across North America, the Middle East, and East Asia, to show that while laboratory values are highly individual, purely personalized intervals routinely overfit, classifying up to 68% of measurements as abnormal, without corresponding associations with adverse clinical outcomes. We then introduce NORMA, a conditional transformer-based framework that generates reference intervals by conditioning on both a patient's history and population-level data about "normal" variation. NORMA-derived intervals achieve higher precision for predicting outcomes, including mortality, acute kidney injury, and chronic disease. These findings caution against over-personalization in laboratory medicine and demonstrate that anchoring individual trajectories to population-level priors outperforms either approach alone. To promote transparency, we publicly release the model, code, and an interactive user interface for accessible, individualized laboratory interpretation.

Introduction

Laboratory testing is among the most frequently performed medical tests, with more than 14 billion tests ordered annually in the United States alone¹. Yet interpretation has changed remarkably little since the 1960s; values are classified as "low," "normal," or "high" against fixed population reference intervals (Pop_{RI}) that represent the central 95% of measurements from ostensibly healthy individuals^{2–8}. Prior studies have recognized that many routine biomarkers fluctuate around narrow individual "setpoints" and that within-person change carries prognostic information that population intervals miss^{9–20}. Despite this, clinical practice often still relies on universal thresholds, partly because of the ease of a one-size-fits-all approach, and because operationalizing individualized interpretation requires modeling patient-specific trajectories at scale.

Efforts to move beyond universal thresholds have taken two directions. The first refines reference intervals for predefined subgroups, for example, sex-specific hematologic ranges or adjusted HbA1c thresholds for patients with hemoglobin variants^{12,14,21–24}. Defining population subgroups introduces several challenges—such groups are often defined coarsely or arbitrarily, may encode existing biases, and cannot capture the full spectrum of individual variation embedded in longitudinal trajectories. The practice of race adjustment, in particular, has been increasingly challenged with major clinical and non-clinical implications for patients^{25–27}. The second approach derives personalized reference intervals (Per_{RI}) directly from each patient's laboratory testing history. Foy et al., for example, showed that deviations from individually derived hematologic setpoints associate more strongly with mortality than deviations from population thresholds, underscoring the prognostic value of within-person change¹⁶. However, purely individualized approaches that derive baselines entirely from a patient's own data risk incorporating unrecognized chronic disease into the estimated setpoint and overfitting to sparse histories, producing overly narrow intervals that label benign physiological variation as pathological or vice versa^{16,18,28,29}.

Here, we ask whether these competing approaches can be combined using nearly 2 billion measurements across 30 analytes, spanning two decades of national longitudinal care, dense ICU monitoring, and perioperative data across three countries (Fig. 1). We first quantify the individuality of routine blood tests and then systematically compare population and personalized reference intervals. We then introduce NORMA (Normal Outcome Range Modeling with Attention), a new conditional transformer framework that combines both approaches to generate individualized reference intervals by anchoring each patient's trajectory to population-level expectations for a healthy state. Rather than choosing between the false dichotomy of purely population and personalized approaches, NORMA learns where each patient's interval is expected to fall between the two extremes. We show that purely personalized intervals overcall abnormalities and lose prognostic signal, that NORMA resolves this tradeoff, and test whether the resulting intervals detect clinically meaningful change months to years before population intervals.

Results

Study populations

We assessed population-based (Pop_{RI}), personalized (Per_{RI}), and NORMA-derived ($NORMA_{RI}$) reference intervals across 30 routine laboratory analytes spanning four clinical panels: complete blood count, comprehensive metabolic panel, lipid panel, and hemoglobin A1c (HbA1c). These panels are used in routine health evaluation, cardiometabolic risk assessment, and diabetes screening or monitoring, as detailed in the Methods. Within this analyte set, NORMA was trained on 3.4 million longitudinal laboratory sequences from two development cohorts: MIMIC-IV (179,601 patients) and EHRSHOT (5,676 patients); no development data were used in external validation (Supplementary Tables 1–3).

We validated NORMA in three independent cohorts that differed in temporal scale and clinical setting. The Clalit Health Services (CHS) cohort included 1,450,862 adults (63% female; mean age 55.1 ± 18.6 years) contributing approximately 1.9 billion laboratory measurements over more than two decades of outpatient follow-up (Fig. 2a; Supplementary Table 3). To ensure stable baseline estimation, we required at least five outpatient measurements per analyte spaced at least 90 days apart, yielding trajectories spanning a median of more than four years.

The eICU Collaborative Research Database (eICU-CRD) cohort included 98,432 ICU patients (46% female; mean age 63.3 ± 16.1 years) across 208 U.S. hospitals, contributing over 20 million measurements with dense sampling (median number of measurements per patient ranged between 8–35 for high-frequency analytes) over a median stay of approximately 8 days.

The INSPIRE cohort included 51,159 surgical patients (52% female; mean age 57.6 ± 14.9 years) undergoing perioperative care in South Korea, contributing over 10 million laboratory measurements collected across preoperative, intraoperative, and postoperative timepoints, with a median follow-up of approximately 175 days.

Laboratory distributions in CHS, EHRSHOT, and INSPIRE largely overlapped population reference intervals, whereas eICU-CRD and MIMIC-IV exhibited shifts consistent with greater rates of critical illness, including lower albumin (ALB) and higher glucose (GLU) variability (Supplementary Table 3).

Lab values are individualized

Across all three cohorts, within-person variability was substantially smaller than between-person variability for most analytes, confirming that laboratory values fluctuate around narrow individual setpoints (Fig. 2b; Supplementary Table 4). We quantified this finding using the individuality index (II), the ratio of within-person to between-person coefficient of variation, where values below 0.6 indicate high biological individuality.

Hematologic indices showed the lowest intra-individual variability relative to inter-individual variability. Mean corpuscular hemoglobin (MCH), mean corpuscular volume (MCV), and red blood cell count (RBC) had individuality indices ranging from approximately 0.2 to 0.6 across CHS, eICU-CRD, and INSPIRE, with values in eICU-CRD generally equal to or lower than those observed in CHS and INSPIRE (Supplementary Table 4). In contrast, electrolytes and metabolic analytes exhibited lower biological individuality; potassium (K), glucose (GLU), sodium (NA), and calcium (CA) had individuality indices ranging from approximately 0.7 to 1.2 across cohorts, with within-person variability approaching the population spread (Fig. 2b; Supplementary Table 4). INSPIRE showed patterns consistent with an intermediate regime between outpatient and ICU settings.

Within-person stability carried prognostic signal. Greater deviation from a patient's baseline, quantified as the absolute z-score of each index measurement relative to that patient's baseline average, was monotonically associated with mortality across all cohorts (Fig. 2c; Supplementary Fig. 3). When stratified by raw analyte values, several biomarkers, including white blood cell count (WBC), glucose (GLU), aspartate aminotransferase (AST), and hemoglobin A1c (HbA1c), exhibited U-shaped mortality curves, in which both abnormally high and low values were associated with elevated risk.

NORMA generates calibrated individualized intervals

NORMA is a conditional, autoregressive transformer framework with multiple configurations that models the distribution of a patient's next laboratory value given their longitudinal measurement history (Fig. 3a; Methods) and population data on "healthy" variation. The model is trained to forecast the next observed value. To derive a personalized reference interval, we condition the query token on values within the population reference range, so that the resulting 95% prediction interval represents the expected range for that patient given their prior trajectory and population context, rather than a strictly disease-free state. We trained NORMA on 3.4 million longitudinal sequences from MIMIC-IV and EHRSHOT under two output parameterizations (Gaussian and quantile), which share the same architecture but differ in distributional assumptions (Methods and Supplementary Table 2).

On a held-out test set, NORMA outperformed all baselines in next-step forecasting accuracy across both output parameterizations. Baseline models included last value carried forward, autoregressive integrated moving average (ARIMA), and patient-specific mean. Mean absolute error was lowest for NORMA quantile (5.9 [IQR 3.6–9.1]) and NORMA Gaussian (6.0 [2.6–10.2]), compared with ARIMA (6.7 [4.3–11.9]), last value carried forward (7.3 [3.9–10.7]), and patient-specific mean (8.4 [4.5–13.6]; Supplementary Table 5; Fig. 3b). Forecasting accuracy varied by analyte, with mean platelet volume (MPV) showing the weakest performance, consistent with its having the smallest training sample size (Fig. 3c; Supplementary Fig. 2).

NORMA's prediction intervals responded appropriately to the factors that govern uncertainty (Fig. 3d; Supplementary Table 6). More prior measurements narrowed intervals, longer prediction horizons widened them, and greater within-person variability produced substantially wider intervals. The quantile parameterization was more responsive than the Gaussian to changes in input features; for example, doubling within-person variability roughly doubled the predicted interval width under the quantile model, compared with minimal change under the Gaussian. Importantly, interval width stabilized after approximately 30 prior measurements and did not continue to narrow with additional data, indicating that the model does not overweight longer histories.

NORMA_{RI} detects abnormalities earlier than population intervals

In all three external validation cohorts, Per_{RI} flagged the most tests as abnormal, Pop_{RI} the fewest, and NORMA_{RI} was intermediate. In CHS, abnormality rates were 29.6% (Pop_{RI}), 39.1% (NORMA_{RI}), and 46.8% (Per_{RI}). In eICU-CRD, where baseline abnormality rates were higher due to acute illness, the same ordering held: 50.2% (Pop_{RI}), 55.8% (NORMA_{RI}), and 68.1% (Per_{RI}). In INSPIRE, abnormality rates followed the same pattern, with values of 35.1% (Pop_{RI}), 42.5% (NORMA_{RI}), and 57.0% (Per_{RI}), consistent with its intermediate clinical setting (Fig. 4a; Fig. 5a; Fig. 6a; Supplementary Table 7).

Among tests classified as normal by population intervals, NORMA_{RI} reclassified 12, 9, and 14 per 100 as abnormal in CHS, eICU-CRD, and INSPIRE, respectively, compared with 27, 37, and 34 per 100 for Per_{RI} (Supplementary Tables 7–8).

NORMA_{RI} flagged abnormalities earlier than population intervals across all cohorts. In CHS, 49% of tests NORMA_{RI} flagged as abnormal were later also flagged by Pop_{RI}, with a median lead time of 8.7 months (IQR 1.7–33.8; Supplementary Table 20; Fig. 4e). In eICU-CRD, 25% were later confirmed by Pop_{RI}, with a median lead time of 34.8 hours (IQR 23.1–72.6; Supplementary Table 20; Fig. 5e). In INSPIRE, 30% were later confirmed by Pop_{RI}, with a median lead time of 23.7 hours (IQR 11.4–54.7), consistent with its perioperative timescale (Fig. 6e; Supplementary Table 20).

NORMA_{RI} detects clinically meaningful abnormalities missed by population intervals

We next asked whether the additional abnormalities detected by NORMA_{RI} carried clinical meaning. Restricting to measurements classified as normal by population intervals, where personalization could add signal beyond standard practice, NORMA_{RI} reclassifications carried substantially higher positive predictive value than Per_{RI}. For every 100 patients Pop_{RI} classified as normal but NORMA_{RI} flagged as abnormal in eICU-CRD, 13 died in hospital versus 10 under Per_{RI} ($\Delta = +3$), 16 developed AKI versus 15 ($\Delta = +1$), and 27 had prolonged ICU stays versus 23 ($\Delta = +4$). In CHS, 30 died versus 24 under Per_{RI} ($\Delta = +6$), 46 developed CKD versus 39 ($\Delta = +7$), and 80 had type 2 diabetes versus 75 ($\Delta = +5$). In INSPIRE, NORMA_{RI} identified 6 more unplanned ICU admissions, 1 more in-hospital death, and 1 more perioperative infection per 100 reclassified patients than Per_{RI} (Supplementary Tables 17–19). Across all assessed outcomes – mortality, type 2 diabetes, and CKD over 10 years in CHS; in-hospital mortality, AKI, sepsis, and prolonged LOS in eICU-CRD; and perioperative mortality, prolonged LOS, unplanned ICU admission, and postoperative infection in INSPIRE – Per_{RI} achieved higher sensitivity but NORMA_{RI} achieved substantially higher specificity and positive predictive value (Figs. 4c,d, 5c,d, 6c,d and Supplementary Figs. 9–11).

NORMA_{RI} improves clinical prediction

In time-to-event analyses across all cohorts, NORMA_{RI} and Pop_{RI} abnormality flags showed comparable prognostic associations with clinical outcomes, while Per_{RI} flags were substantially weaker (Figs. 4b, 5b, 6b and Supplementary Tables 21–29). This was most pronounced in CHS, where NORMA_{RI} abnormality flags

were strongly associated with all-cause mortality for hematologic and metabolic analytes, with several showing 2- to 4-fold elevated risk after adjusting for age and sex – including creatinine (HR 3.30), hemoglobin (HR 3.72), and hematocrit (HR 3.36). Effect sizes were comparable to or slightly exceeded those of Pop_{RI} for most analytes, while Per_{RI} flags carried essentially no prognostic signal.

Mean concordance indices for mortality were similar between NORMA_{RI} and Pop_{RI} across all three cohorts (CHS: 0.752 vs. 0.758; eICU-CRD: 0.567 vs. 0.571; INSPIRE: 0.730 vs. 0.731), while Per_{RI} was consistently lowest (CHS: 0.748; eICU-CRD: 0.562; INSPIRE: 0.695). The gap was most pronounced in INSPIRE, where Per_{RI} concordance (median 0.718, IQR [0.673, 0.742]) fell well below both NORMA_{RI} (0.754 [0.712, 0.780]) and Pop_{RI} (0.742 [0.722, 0.778]; Supplementary Tables 21, 22, 26).

Discussion

Routine laboratory testing is becoming more frequent, more accessible, and more central to medical decision-making^{30,31}. Beyond tests that clinicians order within the healthcare system, patients are increasingly seeking out repeated panel-based screening through direct-to-consumer companies. In parallel, interpretation is shifting away from static cutoffs and monolithic “normal ranges” towards “personalized” ranges that leverage trajectories of variation and individual baselines, yet the implications and risks of this shift remain largely unclear^{17–19,28}. Across nearly two billion laboratory measurements spanning outpatient and intensive care settings, we find that purely personalized reference intervals consistently overcall abnormalities and are poorly associated with adverse outcomes. NORMA balances purely personalized and population intervals by training a transformer to predict what a patient’s next laboratory value is expected to fall if they remain physiologically stable, and defining the reference interval from this conditional distribution. In doing so, NORMA_{RI} inherits the sensitivity of personalized interpretation while maintaining the specificity of population-level definitions of normal variation.

In eICU, Per_{RI} flagged the majority of laboratory measurements as abnormal, and similarly elevated abnormality rates were observed in CHS and INSPIRE compared to population-based intervals, indicating that a purely personalized framework would likely generate alerts for a substantial fraction of tests across critical care, outpatient, and perioperative settings. At the scale of a national health system processing millions of tests annually, this could translate to an enormous burden of false alarms, unnecessary follow-up testing, and potential patient anxiety^{29,32,33}. By contrast, NORMA_{RI} moderated this alert burden, flagging fewer measurements than Per_{RI} while yielding abnormalities that were more strongly enriched for adverse clinical outcomes.

This benefit was not apparent from aggregate discrimination alone. Across individual analytes and outcomes, NORMA_{RI} and Pop_{RI} achieved broadly similar concordance indices, whereas Per_{RI} consistently performed worst. This similarity between NORMA_{RI} and Pop_{RI} might suggest that personalization adds little value; however, the value of NORMA_{RI} is seen in individuals that Pop_{RI} labels normal but NORMA_{RI} flags as abnormal. In this reclassified subset, NORMA_{RI} achieved substantially higher precision and specificity than Per_{RI} .

By anchoring individualized intervals to population-level expectations, NORMA reduces the risk that chronic or subclinical disease is absorbed into a patient’s estimated baseline, a limitation of purely personalized approaches such as Per_{RI} . This limitation is not unique to the Gaussian mixture approach used for Per_{RI} . Even with Bayesian updating or hierarchical extensions, a population prior converges toward the patient’s observed distribution as measurements accumulate, eventually washing out the prior and reproducing the same overcalling behavior^{28,34}. However, such methods are also nondeterministic, computationally expensive, introduce prior-strength and group-structure hyperparameters, and lack native handling of time or irregular

sampling. NORMA avoids this by conditioning on a healthy state at every prediction, regardless of how many measurements are available, which is why its interval width stabilizes rather than continuing to narrow. Furthermore, it does not require explicit pre-filtering of trajectories for stability; it handles irregular measurement spacing natively and its inputs are limited to data already present in most electronic health records. In a clinical deployment, NORMA could provide more "personalized" assessment for patients even without sufficient measurement history.

NORMA builds on a parallel line of work that scales sequence modelling over longitudinal health records into clinical foundation models. Early efforts demonstrated that deep learning over raw EHR streams could match or exceed task-specific models for mortality, readmission, and laboratory forecasting^{35–38}. More recent transformer-based foundation models, trained on event streams from millions of patient timelines, have demonstrated that longitudinal records can support zero- or few- shot forecasting of diagnoses, procedures, and disease progression at increasing scale, with prediction horizons extending years into the future and architectures expanding to incorporate multimodal clinical data^{39–42}. The same paradigm is now extending beyond structured EHRs to continuous physiological streams such as continuous glucose monitoring and other wearables^{43,44}.

These models demonstrate that longitudinal patient representations can support broad outcome prediction across clinical domains. While NORMA learns from patient trajectories, it is distinct in the prediction task. Rather than learning a general representation of trajectories, such as APOLLO, to predict diagnoses, procedures, or downstream clinical outcomes, it models continuous biomarker distributions and estimates the range of values expected for a given patient under a specified future laboratory state⁴². NORMA therefore draws on counterfactual prediction and sequence-editing frameworks, such as CLEF, which ask how a trajectory would change under an imposed condition or intervention^{45–47}. Here, the imposed condition is a future normal laboratory state, allowing NORMA to estimate an individualized healthy-state reference interval rather than simply forecast the next observed measurement. The clinical outcomes evaluated here, including all-cause mortality, acute kidney injury, type 2 diabetes, and chronic kidney disease, were used only for downstream validation rather than model training. This disease-agnostic design suggests that deviations from an individual's expected healthy trajectory may carry prognostic relevance across diverse disease states without requiring outcome-specific retraining. As longitudinal biomarker monitoring expands further into precision medicine, the challenge of reconciling population-derived and individualized reference standards will broaden, and conditional prediction anchored to a healthy reference state may generalize across these settings^{43,44,47,48}.

Our study had several limitations. First, NORMA's current scope is limited by the analytes and clinical inputs used for training. NORMA was trained and evaluated on 30 common blood tests and conditioned only on age, sex, and laboratory trajectories; it does not currently incorporate comorbidities, medications, or other clinical context, which could further refine expected trajectories and abnormality thresholds⁴⁹. Performance was also not uniform across analytes. MPV showed the lowest forecasting accuracy, consistent with its having the smallest training sample size and weakest forecasting accuracy among the 30 analytes. Second, some modeling assumptions may affect calibration. In CHS, we modeled predictive distributions as Gaussian, which may not fully capture skewed or heavy-tailed analytes. However, we implemented quantile regression in eICU, and results were consistent across both parameterizations, suggesting that NORMA's clinical value does not depend on a strict Gaussian assumption. Broader exploration of output distributions may further improve calibration for skewed or heavy-tailed analytes. Third, generalizability and clinical impact remain to be established prospectively. The CHS analysis was conducted within a single national health system; eICU, although geographically diverse, represents only the ICU segment of hospital care; and INSPIRE reflects perioperative care at a single academic medical center. Performance may differ in emergency departments, primary care clinics, non-surgical inpatient settings, or populations with different demographics and laboratory utilization patterns. Finally, this study evaluated retrospective associations between abnormality flags and

clinical outcomes. Prospective trials are needed to determine whether NORMA_{RI}-augmented interpretation accelerates time to diagnosis, alters downstream testing, and influences clinician behavior. Such evaluation may further benefit from cohort-specific fine-tuning, for example on Clalit data before deployment, which could improve local performance at the cost of generalizability.

Individualized laboratory interpretation can be made more precise and more clinically useful by combining patient-specific trajectories with population-level expectations and uncertainty-aware prediction. By validating across longitudinal outpatient and acute settings, outcome horizons, and patient populations, we show that this framework generalizes beyond a single clinical context. NORMA provides a framework for integrating personalized laboratory interpretation into routine practice, balancing the sensitivity of personalized intervals with the specificity of population-anchored prediction.

Methods

Study Populations

We used five cohorts spanning model development and external validation. Model development used MIMIC-IV and EHRSHOT. External validation used three independent cohorts representing distinct clinical settings and time horizons: Clalit Health Services (CHS), a national longitudinal health system cohort from Israel; the eICU Collaborative Research Database (eICU-CRD), a multicenter critical care cohort from the United States; and INSPIRE, a perioperative cohort from Seoul National University Hospital in South Korea. Across these cohorts, we evaluated three reference interval frameworks: population-based reference intervals (PopRI), personalized reference intervals (Per_{RI}), and NORMA-derived reference intervals (NORMA_{RI}). No development data were used in external validation.

MIMIC-IV and EHRSHOT. MIMIC-IV contributed 179,601 patients from a tertiary-care hospital system and was used for acute-care, short-horizon model development. EHRSHOT contributed 5,676 patients from longitudinal electronic health records and was used for intermediate-horizon model development. Together, these cohorts provided the development data used for NORMA training, validation, and held-out testing.

Clalit Health Services. The CHS cohort comprises nationwide longitudinal electronic healthcare data from Israel's largest health maintenance organization⁵⁰. We included adults aged 18–99 years with repeated outpatient laboratory testing between 2000 and 2024. For each analyte, eligibility required at least five outpatient measurements spaced at least 90 days apart prior to January 1, 2015 (baseline period) and at least one measurement between January 1, 2015 and January 1, 2016 (classification period). We excluded hospitalizations, emergency department visits, and urgent encounters during baseline ascertainment. The first measurement during the classification period served as the index value and defined the index date. We assessed eligibility independently per analyte, allowing individuals to contribute to multiple analyses.

We followed individuals for up to ten years from each analyte-specific index date. For those without events, we censored follow-up at the earliest of health plan disenrollment or death. Patients remaining enrolled without a recorded death were censored at the last available date in the dataset (January 1, 2024). Primary outcomes included all-cause mortality and incident diagnoses of type 2 diabetes (T2D) and chronic kidney disease (CKD). We ascertained outcomes using ICD-9/10 codes supplemented by laboratory criteria. We defined T2D as HbA1c $\geq 6.5\%$ or fasting glucose ≥ 126 mg/dL on two occasions; CKD as an estimated glomerular filtration rate < 60 mL/min/1.73 m² on two measurements at least 90 days apart.

eICU Collaborative Research Database. The eICU Collaborative Research Database (eICU-CRD v2.0) is a multicenter critical care cohort comprising more than 200,859 ICU admissions from 208 hospitals across

the United States⁵¹. We included adults aged 18–99 years with repeated laboratory measurements during their ICU stay. We applied a baseline filter requiring at least five measurements. After filtering, 98,432 unique patients contributed over 20 million measurements across 29 analytes.

For each patient–analyte sequence, we used the first 75% of measurements as the baseline to estimate Per_{RI} and $NORMA_{RI}$, and we used the remaining measurements as index values for classification. Primary outcomes included in-hospital mortality, acute kidney injury (AKI), sepsis, and prolonged ICU length of stay (>7 days). Outcomes were ascertained using ICD-9 diagnosis codes and structured clinical documentation.

INSPIRE. The INSPIRE dataset is a publicly available perioperative research cohort comprising approximately 130,000 surgical cases from a single academic medical center (Seoul National University Hospital) in South Korea between 2011 and 2020⁵². We included adults aged 20–90 years with repeated laboratory measurements during the perioperative period. We applied a baseline filter requiring at least five measurements. After filtering, 51,159 unique patients contributed over 10 million measurements across 19 analytes.

For each patient–analyte sequence, we used the first 75% of measurements as the baseline to estimate Per_{RI} and $NORMA_{RI}$, and we used the remaining measurements as index values for classification. Primary outcomes included in-hospital mortality, prolonged length of stay, unplanned ICU admission, and perioperative infection. Outcomes were ascertained using ICD-10-CM diagnosis codes and structured clinical documentation.

Laboratory Measurements

We selected 30 routinely measured blood analytes from common clinical panels based on clinical ubiquity and sufficient repeated measurement across development and validation datasets. We grouped analytes into four standard clinical panels: complete blood count (hematocrit, hemoglobin, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, mean corpuscular volume, mean platelet volume, platelet count, red blood cell count, red cell distribution width, white blood cell count); comprehensive metabolic panel (sodium, potassium, chloride, calcium, bicarbonate, blood urea nitrogen, creatinine, glucose, aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase, total bilirubin, direct bilirubin, total protein, albumin); lipid panel (total cholesterol, low-density lipoprotein cholesterol, high-density lipoprotein cholesterol, triglycerides); glycemic control (glycated hemoglobin).

We mapped all laboratory results to Logical Observation Identifiers Names and Codes (LOINC)⁵³. We retained only positive values, standardized units to canonical formats, and excluded analyte-specific outliers exceeding ± 3 standard deviations from the analyte median. When duplicate results shared identical timestamps, we averaged values within each analyte.

Reference Interval Frameworks

Population Reference Intervals. We defined Pop_{RI} using externally specified laboratory reference ranges and clinical decision thresholds rather than estimating cohort-specific intervals. For each analyte, we used the adult reference range reported by the American Board of Internal Medicine Laboratory Test Reference Ranges⁵⁴. Where sex-specific reference intervals were provided, classifications were made using the patient's sex-specific range. Supplementary Table 1 provides the reference interval, sex stratification, and canonical unit used for each analyte.

Personalized Reference Intervals. We estimated Per_{RI} following Foy *et al.* For each patient–analyte pair, we fit Gaussian mixture models with one to three components to baseline measurements and selected model order using the Akaike information criterion¹⁶. From the selected model, we chose the mixture component that explained the largest number of that patient's baseline measurements and treated it as the individual's

physiological setpoint. We defined the Per_{RI} as the mean of this component ± 2 standard deviations.

NORMA (Normal Outcome Range Modeling with Attention)

NORMA is a conditional, decoder-only transformer that models the distribution of a patient's next laboratory value given their longitudinal measurement history and a query specifying a future health state and prediction horizon. We trained the model on 3.4 million longitudinal sequences from MIMIC-IV and EHRSHOT, requiring at least three repeated measurements per patient without constraints on sampling intervals or clinical setting^{55,56}. Data were split at the patient level into training (70%), validation (10%), and test (20%) sets to prevent leakage across partitions. Each sequence was tokenized, padded, and masked to handle variable-length histories. A weighted sampling scheme was used during training to balance representation across 30 laboratory analytes. Alternative input encodings with justifications are summarized (Supplementary Table 3).

Input representation. For a given patient-biomarker pair, we denote the ordered sequence of observed laboratory values as $x = (x_1, \dots, x_T)$, with corresponding measurement times $t = (t_1, \dots, t_T)$ and clinical states $s = (s_1, \dots, s_T)$, where each s_i indicates the laboratory state relative to the population reference interval (low, normal, or high).

The model processes these inputs as a sequence of three token types: a context token, history tokens, and a query token. The context token z_c encodes static patient covariates: sex g , age a , and laboratory test code c , as a sum of learned embeddings:

$$z_c = e_g(g) + e_a(a) + e_c(c)$$

Each history token z_i^{hist} represents one prior measurement and combines three components:

$$z_i^{hist} = f_v(\tilde{x}_i) + e_s(s_i) + f_\tau(\Delta t_i)$$

where f_v is a linear projection of the input value $\tilde{x}_i$, e_s is a learned state embedding, and f_τ encodes the inter-measurement interval $\Delta t_i = t_i - t_{i-1}$.

The query token, z_{query} , specifies the context at the time of prediction:

$$z_{query} = e_s(s_{T+1}) + f_h(t_{T+1} - t_T)$$

where s_{T+1} denotes the future health state and $t_{T+1} - t_T$ the prediction horizon. The full sequence is processed using a causally masked Transformer decoder to obtain the predictive distribution of the next observed laboratory value by $P(x_{T+1}|x, t, s, g, a, c, s^*, t^*)$.

Training Objective. Given the input sequence, the model is optimized to predict the conditional distribution of the next observed value x_{T+1} . We trained NORMA under two output parameterizations.

Under the Gaussian parameterization, the model jointly estimates a predicted mean μ and log-variance $\log \sigma^2$:

$$L = \frac{1}{2}[\log \sigma^2 + (x_{T+1} - \mu)^2 / \sigma^2]$$

Under the quantile parameterization, the model directly predicts fixed quantiles $\tau \in \{0.025, 0.25, 0.50, 0.75, 0.975\}$ using a pinball loss without distributional assumptions:

$$L_\tau = \tau \times \max(0, x_{T+1} - q_\tau) + (1 - \tau) \times \max(0, q_\tau - x_{T+1})$$

where q_τ is the predicted τ -th quantile. Both parameterizations share the same core architecture and were trained on the same development data.

NORMA reference intervals. We define the NORMA reference interval (NORMA_{RI}) as the 95% prediction interval obtained by conditioning on a future normal laboratory state. Under the Gaussian parameterization, this corresponds to $[\mu - 1.96\sigma, \mu + 1.96\sigma]$; under the quantile parameterization, it corresponds to $[\hat{q}_{0.025}, \hat{q}_{0.975}]$.

Evaluation. We first evaluated forecasting performance for next-step prediction. For point accuracy, we used the 50th percentile (median) prediction and compared it against three baselines: a person-specific historical mean model, last observation carried forward ("Last"), and an autoregressive integrated moving average model (ARIMA). Performance was quantified using mean absolute error (MAE), mean absolute percentage error (MAPE), and coefficient of determination (R^2).

Sensitivity Analysis. To assess how NORMA captures uncertainty across clinical contexts, we conducted one-at-a-time sensitivity sweeps over five input features: patient age, sex, number of prior measurements, prediction horizon, and within-sequence variability. For each of 30 laboratory tests, we constructed a synthetic baseline sequence: 10 measurements spaced 90 days apart, each set to the midpoint of the sex-specific population reference interval, for a 50-year-old male with a 30-day prediction horizon. We then varied one feature at a time while holding all others at baseline values. Age was swept from 20 to 80 years in 5-year increments; history length from 2 to 300 measurements; and prediction horizon from 7 days to 10 years. For within-sequence variability, we parameterized the standard deviation as a multiplier of one-tenth of the reference range width (0.0 to 3.0) and, at each noise level, drew 30 independent histories from $N(\text{midpoint}, \sigma)$ to capture sampling variability. For each configuration, we recorded the predicted median (50th quantile), the 90% prediction interval width, and the percent change in interval width relative to baseline, enabling direct comparison of sensitivity magnitude across biomarkers.

NORMA configuration. For CHS validation, we applied NORMA with the default configuration: Time2Vec temporal encoding, ternary health-state indicators, raw-value laboratory inputs, raw age projected linearly, and the Gaussian output parameterization. For eICU validation, we used log-delta-t temporal encoding with periodic components, ternary health-state indicators, within-sequence normalization of laboratory values, binned age embeddings in decade-wide bins, a dedicated context token for patient covariates, and the quantile output parameterization. INSPIRE used the same configuration as eICU. All cohort-specific NORMA configurations are summarized in Supplementary Table 3.

Statistical Analysis

Mortality association analysis. We assessed the association between laboratory values and mortality using two complementary approaches. First, we grouped index-period measurements into quintiles of raw analyte values within each analyte and estimated mortality rates per quintile. Second, to evaluate whether deviation from a patient's personal baseline predicts mortality, we computed a deviation score for each index measurement as the absolute z-score from the patient's baseline mean ($|\text{value} - \mu_{\text{baseline}}| / \sigma_{\text{baseline}}$), where μ_{baseline} and σ_{baseline} are the mean and standard deviation of the patient's baseline measurements. We grouped deviation scores into deciles and estimated mortality rates within each decile. For both analyses, we computed 95% confidence intervals using Wilson score intervals. Analytes with fewer than 20 observations were excluded.

Abnormality classification. We classified each index measurement as normal or abnormal under Pop_{RI} , Per_{RI} , and NORMA_{RI} . To avoid incorporating pre-existing pathology into Per_{RI} thresholds, we excluded patients whose Per_{RI} mean fell outside the Pop_{RI} . Measurements classified as Pop_{RI} -abnormal were also classified as abnormal using Per_{RI} and NORMA_{RI} .

Clinical Risk Stratification and Prediction. We evaluated whether abnormal classifications under each framework identified individuals at elevated risk of future events. For each analyte and framework (Pop_{RI} , Per_{RI} , NORMA_{RI}), we defined a binary indicator of abnormality for values outside the corresponding interval.

For each analyte–outcome pair, we calculated positive predictive value (probability of the event given abnormal classification), sensitivity (probability of abnormal classification among those with the event), and specificity (probability of normal classification among those without the event). To isolate differences between interval paradigms, we restricted analyses to measurements within the Pop_{RI} , because values outside the Pop_{RI} necessarily fall outside Per_{RI} and NORMA_{RI} .

We estimated associations using Cox proportional hazards models with a single binary abnormality indicator adjusted for age and sex. We split data into training (60 percent) and test (40 percent) sets stratified by event status and approximate event time in one-year bins. We evaluated performance at 1, 3, 5, and 10 years using time-dependent receiver operating characteristic curves and quantified uncertainty with 95% confidence intervals from 1,000 bootstrap resamples. We applied Benjamini-Hochberg false discovery rate correction to account for multiple comparisons across outcomes and analytes.

Ethics Approval

The use of Clalit Health Services data in this study was approved by the Clalit Health Services Institutional Review Board (Helsinki Committee). MIMIC-IV, eICU-CRD, and INSPIRE were accessed under the PhysioNet Credentialed Health Data Use Agreement; EHRSHOT was accessed under the Stanford Research Use Agreement. All datasets were de-identified prior to release.

Data Availability

This study utilizes two publicly available datasets for the development of NORMA: EHRSHOT and MIMIC-IV, both accessible to qualified researchers under their respective data-use agreements. The eICU Collaborative Research Database is also publicly available via PhysioNet. Clalit Health Services data are not publicly available.

Code Availability

The full source code for the NORMA model, including data processing, training, and evaluation pipelines, is publicly available at <https://github.com/aashnapshah/NORMA>. The interactive web-based user interface for individualized laboratory interpretation is available at <https://norma-tpy0.onrender.com/>.

Acknowledgments

We gratefully acknowledge support from the Ivan and Francesca Berkowitz Family Living Laboratory Collaboration at Harvard Medical School and Clalit Research Institute, NIEHS R01ES032470, and NIDDK R01DK137993.

References

1. Danielle B Freedman. Towards better test utilization - strategies to improve physician ordering and their impact on patient outcomes. *EJIFCC*, 26(1):15–30, January 2015.
2. G H Guyatt, A D Oxman, M Ali, A Willan, W McIlroy, and C Patterson. Laboratory diagnosis of iron-deficiency anemia: an overview. *J. Gen. Intern. Med.*, 7(2):145–153, March 1992.
3. P Newsome, R Cramb, S Davison, J Dillon, M Foulerton, E Godfrey, Richard Hall, Ulrike Harrower, M Hudson, A Langford, A Mackie, R Mitchell-Thain, K Sennett, N Sheron, J Verne, Martine Walmsley, and A Yeoman. Guidelines on the management of abnormal liver blood tests. *Gut*, 67:6–19, November 2017.
4. Silvio E Inzucchi. Clinical practice. diagnosis of diabetes. *N. Engl. J. Med.*, 367(6):542–550, August 2012.
5. Kenneth A Sikaris. Enhancing the clinical value of medical laboratory testing. *Clin. Biochem. Rev.*, 38(3): 107–114, November 2017.
6. Ferruccio Ceriotti, Rolf Hinzmann, and Mauro Panteghini. Reference intervals: the way forward. *Ann. Clin. Biochem.*, 46(Pt 1):8–17, January 2009.
7. Richard C Friedberg, Rhona Souers, Elizabeth A Wagar, Ana K Stankovic, Paul N Valenstein, and College of American Pathologists. The origin of reference intervals: A college of american pathologists Q-probes study of “normal ranges” used in 163 clinical laboratories. *Arch. Pathol. Lab. Med.*, 131(3):348–357, March 2007.
8. Joris L J M Müskens, Rudolf Bertijn Kool, Simone A van Dulmen, and Gert P Westert. Overuse of diagnostic testing in healthcare: a systematic review. *BMJ Qual. Saf.*, 31(1):54–63, January 2022.
9. Brian A Ference, Henry N Ginsberg, Ian Graham, Kausik K Ray, Chris J Packard, Eric Bruckert, Robert A Hegele, Ronald M Krauss, Frederick J Raal, Heribert Schunkert, Gerald F Watts, Jan Borén, Sergio Fazio, Jay D Horton, Luis Masana, Stephen J Nicholls, Børge G Nordestgaard, Bart van de Sluis, Marja-Riitta Taskinen, Lale Tokgözoğlu, Ulf Landmesser, Ulrich Laufs, Olov Wiklund, Jane K Stock, M John Chapman, and Alberico L Catapano. Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. evidence from genetic, epidemiologic, and clinical studies. a consensus statement from the european atherosclerosis society consensus panel. *Eur. Heart J.*, 38(32):2459–2472, August 2017.
10. Thore Buerge, Jakob Steinfeldt, Greg Ruyoga, Maik Pietzner, Daniele Bizzarri, Dina Vojinovic, Julius Upmeyer Zu Belzen, Lukas Look, Paul Kittner, Lara Christmann, Noah Hollmann, Henrik Strangalies, Jana M Braunger, Benjamin Wild, Scott T Chiesa, Joachim Spranger, Fabian Klostermann, Erik B van den Akker, Stella Trompet, Simon P Mooijaart, Naveed Sattar, J Wouter Jukema, Birgit Lavrijssen, Maryam Kavousi, Mohsen Ghanbari, Mohammad A Ikram, Eline Slagboom, Mika Kivimäki, Claudia Langenberg, John Deanfield, Roland Eils, and Ulf Landmesser. Metabolomic profiles predict individual multidisease outcomes. *Nat. Med.*, 28(11):2309–2320, November 2022.
11. Edoardo G Giannini, Roberto Testa, and Vincenzo Savarino. Liver enzyme alteration: a guide for clinicians. *CMAJ*, 172(3):367–379, February 2005.
12. M C Walters and H T Abelson. Interpretation of the complete blood count. *Pediatr. Clin. North Am.*, 43(3): 599–622, June 1996.
13. James C Boyd. Defining laboratory reference values and decision limits: populations, intervals, and interpretations. *Asian J. Androl.*, 12(1):83–90, January 2010.
14. Arjun K Manrai, Chirag J Patel, and John P A Ioannidis. In the era of precision medicine and big data, who is normal? *JAMA*, 319(19):1981–1982, May 2018.
15. D J Nazir, R S Roberts, S A Hill, and M J McQueen. Monthly intra-individual variation in lipids over a

- 1-year period in 22 normal subjects. *Clin. Biochem.*, 32(5):381–389, July 1999.
16. Brody H Foy, Rachel Petherbridge, Maxwell T Roth, Cindy Zhang, Daniel C De Souza, Christopher Mow, Hasmukh R Patel, Chhaya H Patel, Samantha N Ho, Evie Lam, Camille E Powe, Robert P Hasserjian, Konrad J Karczewski, Veronica Tozzo, and John M Higgins. Haematological setpoints are a stable and patient-specific deep phenotype. *Nature*, 637(8045):430–438, January 2025.
 17. Shuo Wang, Min Zhao, Zihan Su, and Runqing Mu. Annual biological variation and personalized reference intervals of clinical chemistry and hematology analytes. *Clin. Chem. Lab. Med.*, 60(4):606–617, March 2022.
 18. Abdurrahman Coskun, Sverre Sandberg, Ibrahim Unsal, Fulya G Yavuz, Coskun Cavusoglu, Mustafa Serteser, Meltem Kilercik, and Aasne K Aarsand. Personalized reference intervals - statistical approaches and considerations. *Clin. Chem. Lab. Med.*, 60(4):629–635, March 2022.
 19. Abdurrahman Coşkun, Sverre Sandberg, Ibrahim Unsal, Coskun Cavusoglu, Mustafa Serteser, Meltem Kilercik, and Aasne K Aarsand. Personalized reference intervals in laboratory medicine: A new model based on within-subject biological variation. *Clin. Chem.*, 67(2):374–384, January 2021.
 20. A E Obstfeld, K Patel, J C Boyd, J Drees, D T Holmes, J P Ioannidis, and A K Manrai. Data mining approaches to reference interval studies. *Clinical Chemistry*, 67(9):1175–1181, 2021.
 21. Mary E Lacy, Gregory A Wellenius, Anne E Sumner, Adolfo Correa, Mercedes R Carnethon, Robert I Liem, James G Wilson, David B Sacks, David R Jacobs, Jr, April P Carson, Xi Luo, Annie Gjelsvik, Alexander P Reiner, Rakhi P Naik, Simin Liu, Solomon K Musani, Charles B Eaton, and Wen-Chih Wu. Association of sickle cell trait with hemoglobin A1c in african americans. *JAMA*, 317(5):507–515, February 2017.
 22. Kenneth R Feingold. Guidelines for the management of high blood cholesterol. In *Endotext [Internet]*. MDText. com, Inc., 2025.
 23. O Yaw Addo, Emma X Yu, Anne M Williams, Melissa Fox Young, Andrea J Sharma, Zuguo Mei, Nicholas J Kassebaum, Maria Elena D Jefferds, and Parminder S Suchdev. Evaluation of hemoglobin cutoff levels to define anemia among healthy individuals. *JAMA Netw. Open*, 4(8):e2119123, August 2021.
 24. D H Rushton, R Dover, A W Sainsbury, M J Norris, J J Gilkes, and I D Ramsay. Why should women have lower reference limits for haemoglobin and ferritin concentrations than men? *BMJ*, 322(7298):1355–1357, June 2001.
 25. James A Diao, Yixuan He, Rohan Khazanchi, Max Jordan Nguemeni Tiako, Jonathan I Witonsky, Emma Pierson, Pranav Rajpurkar, Jennifer R Elhawary, Luke Melas-Kyriazi, Albert Yen, Alicia R Martin, Sean Levy, Chirag J Patel, Maha Farhat, Luisa N Borrell, Michael H Cho, Edwin K Silverman, Esteban G Burchard, and Arjun K Manrai. Implications of race adjustment in lung-function equations. *N. Engl. J. Med.*, 390(22):2083–2097, June 2024.
 26. Darshali A Vyas, Leo G Eisenstein, and David S Jones. Hidden in plain sight—reconsidering the use of race correction in clinical algorithms. *New England Journal of Medicine*, 383(9):874–882, 2020.
 27. Aashna P Shah, James A Diao, Emma Pierson, Chirag J Patel, and Arjun K Manrai. Disentangling proxies of demographic adjustments in clinical equations. *arXiv [q-bio.QM]*, November 2025.
 28. Nabihah Tayob and Ziding Feng. Personalized statistical learning algorithms to improve the early detection of cancer using longitudinal biomarkers. *Cancer Biomark.*, 33(2):199–210, 2022.
 29. Isaac S Kohane, Daniel R Masys, and Russ B Altman. The incidentalome: a threat to genomic medicine. *JAMA*, 296(2):212–215, July 2006.
 30. Christina Koch, Katherine Roberts, Christopher Petruccelli, and Daniel J Morgan. The frequency of unnecessary testing in hospitalized patients. *Am. J. Med.*, 131(5):500–503, May 2018.


31. Henrik L Jørgensen and Bent S Lind. Blood tests - too much of a good thing. *Scand. J. Prim. Health Care*, 40(2):165–166, June 2022.
32. Christopher Naugler and Irene Ma. More than half of abnormal results from laboratory tests ordered by family physicians could be false-positive. *Can. Fam. Physician*, 64(3):202–203, March 2018.
33. Tony Badrick, Joe M El-Khoury, and Elvar Theodorsson. Laboratory reference intervals - history and modern approaches for improved utility. *Scand. J. Clin. Lab. Invest.*, 85(4):229–241, June 2025.
34. Davood Roshan, John Ferguson, Charles R Pedlar, Andrew Simpkin, William Wyns, Frank Sullivan, and John Newell. A comparison of methods to generate adaptive reference ranges in longitudinal monitoring. *PLoS One*, 16(2):e0247338, February 2021.
35. Alvin Rajkomar, Eyal Oren, Kai Chen, Andrew M Dai, Nissan Hajaj, Michaela Hardt, Peter J Liu, Xiaobing Liu, Jake Marcus, Mimi Sun, Patrik Sundberg, Hector Yee, Kun Zhang, Yi Zhang, Gerardo Flores, Gavin E Duggan, Jamie Irvine, Quoc Le, Kurt Litsch, Alexander Mossin, Justin Tansuwan, De Wang, James Wexler, Jimbo Wilson, Dana Ludwig, Samuel L Volchenboum, Katherine Chou, Michael Pearson, Srinivasan Madabushi, Nigam H Shah, Atul J Butte, Michael D Howell, Claire Cui, Greg S Corrado, and Jeffrey Dean. Scalable and accurate deep learning with electronic health records. *NPJ Digit. Med.*, 1(1):18, May 2018.
36. Matthew B A McDermott, Bret Nestor, Peniel Argaw, and Isaac Kohane. Event stream GPT: A data pre-processing and modeling library for generative, pre-trained transformers over continuous-time sequences of complex events. *arXiv [cs.LG]*, June 2023.
37. Zhichao Yang, Avijit Mitra, Weisong Liu, Dan Berlowitz, and Hong Yu. TransformEHR: transformer-based encoder-decoder generative model to enhance prediction of disease outcomes using electronic health records. *Nat. Commun.*, 14(1):7857, November 2023.
38. Lavender Yao Jiang, Xujin Chris Liu, Nima Pour Nejatian, Mustafa Nasir-Moin, Duo Wang, Anas Abidin, Kevin Eaton, Howard Antony Riina, Ilya Laufer, Paawan Punjabi, Madeline Miceli, Nora C Kim, Cordelia Orillac, Zane Schnurman, Christopher Livia, Hannah Weiss, David Kurland, Sean Neifert, Yosef Dastagirzada, Douglas Kondziolka, Alexander T M Cheung, Grace Yang, Ming Cao, Mona Flores, Anthony B Costa, Yindalon Aphinyanaphongs, Kyunghyun Cho, and Eric Karl Oermann. Health system-scale language models are all-purpose prediction engines. *Nature*, 619(7969):357–362, July 2023.
39. Artem Shmatko, Alexander Wolfgang Jung, Kumar Gaurav, Søren Brunak, Laust Hvas Mortensen, Ewan Birney, Tom Fitzgerald, and Moritz Gerstung. Learning the natural history of human disease with generative transformers. *Nature*, 647(8088):248–256, November 2025.
40. Shane Waxler, Paul Blazek, Davis White, Daniel Sneider, Kevin Chung, Mani Nagarathnam, Patrick Williams, Hank Voeller, Karen Wong, Matthew Swanhorst, Sheng Zhang, Naoto Usuyama, Cliff Wong, Tristan Naumann, Hoifung Poon, Andrew Loza, Daniella Meeker, Seth Hain, and Rahul Shah. Generative medical event models improve with scale. *arXiv [cs.LG]*, November 2025.
41. Pawel Renc, Yugang Jia, Anthony E Samir, Jaroslaw Was, Quanzheng Li, David W Bates, and Arkadiusz Sitek. Zero shot health trajectory prediction using transformer. *NPJ Digit. Med.*, 7(1):256, September 2024.
42. Andrew Zhang, Tong Ding, Sophia J Wagner, Caiwei Tian, Ming Y Lu, Rowland Pettit, Joshua E Lewis, Alexandre Misrahi, Dandan Mo, Long Phi Le, and Faisal Mahmood. A multimodal and temporal foundation model for virtual patient representations at healthcare system scale. *arXiv [cs.LG]*, April 2026.
43. Guy Lutsker, Gal Sapir, Smadar Shilo, Jordi Merino, Anastasia Godneva, Jerry R Greenfield, Dorit Samocha-Bonet, Raja Dhir, Francisco Gude, Shie Mannor, Eli Meirom, Eric P Xing, Gal Chechik, Hagai Rossman, and Eran Segal. A foundation model for continuous glucose monitoring data. *Nature*, 650

(8103):978–986, February 2026.


44. Ahmed A Metwally, A Ali Heydari, Daniel McDuff, Alexandru Solot, Zeinab Esmaeilpour, Anthony Z Faranesh, Menglian Zhou, Girish Narayanswamy, Maxwell A Xu, Xin Liu, Yuzhe Yang, David B Savage, Mark Malhotra, Conor Heneghan, Shwetak Patel, Cathy Speed, and Javier L Prieto. Insulin resistance prediction from wearables and routine blood biomarkers. *Nature*, March 2026.
45. Valentyn Melnychuk, Dennis Frauen, and Stefan Feuerriegel. Causal transformer for estimating counterfactual outcomes. *arXiv [cs.LG]*, April 2022.
46. Michelle M Li, Kevin Li, Yasha Ektefaie, Ying Jin, Yepeng Huang, Shvat Messica, Tianxi Cai, and Marinka Zitnik. Controllable sequence editing for biological and clinical trajectories. *arXiv [cs.LG]*, February 2025.
47. Shakson Isaac, Yentl Collin, and Chirag Patel. SSM-CGM: Interpretable state-space forecasting model of continuous glucose monitoring for personalized diabetes management. *arXiv [cs.LG]*, October 2025.
48. Martin W McIntosh, Nicole Urban, and Beth Karlan. Generating longitudinal screening algorithms using novel biomarkers for disease. *Cancer Epidemiol. Biomarkers Prev.*, 11(2):159–166, February 2002.
49. Ruth Johnson, Uri Gottlieb, Galit Shaham, Lihi Eisen, Jacob Waxman, Stav Devons-Sberro, Curtis R Ginder, Peter Hong, Raheel Sayeed, Xiaorui Su, Ben Y Reis, Ran D Balicer, Noa Dagan, and Marinka Zitnik. ClinVec: Unified embeddings of clinical codes enable knowledge-grounded AI in medicine. *medRxiv*, May 2025.
50. Ran D Balicer, Efrat Shadmi, Nicky Lieberman, Sari Greenberg-Dotan, Margalit Goldfracht, Liora Jana, Arnon D Cohen, Sigal Regev-Rosenberg, and Orit Jacobson. Reducing health disparities: strategy planning and implementation in israel's largest health care organization. *Health Serv. Res.*, 46(4): 1281–1299, August 2011.
51. Tom J Pollard, Alistair E W Johnson, Jesse D Raffa, Leo A Celi, Roger G Mark, and Omar Badawi. The eICU collaborative research database, a freely available multi-center database for critical care research. *Sci. Data*, 5(1):180178, September 2018.
52. Leerang Lim, Hyeonhoon Lee, Chul-Woo Jung, Dayeon Sim, Xavier Borrat, Tom J Pollard, Leo A Celi, Roger G Mark, Simon T Vistisen, and Hyung-Chul Lee. INSPIRE, a publicly available research dataset for perioperative medicine. *Sci. Data*, 11(1):655, June 2024.
53. Clement J McDonald, Stanley M Huff, Jeffrey G Suico, Gilbert Hill, Dennis Leavelle, Raymond Aller, Arden Forrey, Kathy Mercer, Georges DeMoor, John Hook, Warren Williams, James Case, and Pat Maloney. LOINC, a universal standard for identifying laboratory observations: a 5-year update. *Clin. Chem.*, 49(4): 624–633, April 2003.
54. American Board of Internal Medicine. ABIM laboratory test reference ranges. Technical report, January 2025.
55. Alistair Johnson, Lucas Bulgarelli, Tom Pollard, Brian Gow, Benjamin Moody, Steven Horng, Leo Anthony Celi, and Roger Mark. MIMIC-IV, October 2024.
56. Michael Wornow, Rahul Thapa, Ethan Steinberg, Jason A Fries, and Nigam H Shah. EHRSHOT: An EHR benchmark for few-shot evaluation of foundation models. *arXiv [cs.LG]*, July 2023.

Figures


a Contextual Sequence Modeling




b Training and Validation Cohorts




c Reference Interval Paradigms



≥20 years | 37M sequences | 30 labs



d Clinical Utility of NORMA_{RI}



e Early Risk Stratification






Fig. 1 | NORMA framework overview. **a)** NORMA conditions on laboratory history, time, sex, age, and population reference state to predict the distribution of the next healthy laboratory value. **b)** Study design: training on EHRSHOT and MIMIC-IV; external validation on Clalit Health Services, the eICU Collaborative Research Database, and the INSPIRE cohort. Across all cohorts: 20 years, 37 million sequences, and 30 analytes spanning four clinical panels. **c)** Comparison of Pop_{RI}, Per_{RI}, and NORMA_{RI}. **d)** Illustrative example: a glucose value classified as normal by Pop_{RI} but flagged abnormal by Per_{RI} and NORMA_{RI}, with corresponding change in specificity, sensitivity, and precision for type 2 diabetes classification. **e)** Schematic of early abnormality detection by NORMA_{RI} relative to Pop_{RI} and Per_{RI} over time. Created with BioRender.com.

a Cohort Selection



b Intra-Patient vs Inter-Patient Variation



c Mortality Association






Fig. 2 | Within-person biological variability and mortality association. **a)** Cohort Selection. Cohort selection and inclusion criteria for Clalit Health Services, the eICU Collaborative Research Database, and the INSPIRE cohort. **b)** Intra-Patient vs Inter-Patient Variation. Within-person versus between-person coefficient of variation for 30 analytes, with the individuality index shown below. The dashed line indicates an individuality index of 0.6. **c)** Mortality Association. Observed mortality rate by quintile of z-score from the patient baseline mean in Clalit Health Services, the eICU Collaborative Research Database, and the INSPIRE cohort. Shaded bands denote 95% confidence intervals.


a NORMA Architecture



b Forecasting Performance



c Per-Analyte Performance



d Sensitivity Analysis




Fig. 3 | NORMA architecture, forecasting accuracy, and sensitivity analysis. **a)** NORMA Architecture. Model architecture: demographic context, laboratory history, and query tokens are jointly embedded and processed by a causal transformer decoder to produce quantile predictions for the next laboratory value. **b)** Forecasting Performance. Held-out test-set forecasting accuracy (mean absolute error, mean absolute percentage error, and coefficient of determination) for NORMA (quantile) versus ARIMA, population mean, and last-value-carried-forward baselines. Error bars denote 95% bootstrap confidence intervals. **c)** Per-Analyte Performance. Per-analyte coefficient of determination grouped by clinical panel. **d)** Sensitivity Analysis. Sensitivity of prediction interval width and midpoint deviation (each as a percentage of Pop_{RI} width) to history length, forecast horizon, and within-person variability.




Fig. 4 | Reclassification performance and early risk stratification in Clalit Health Services. a) Reclassification Rate. Reclassification rate by analyte among Pop_{RI}-normal tests. **b)** Hazard Ratios. Cox hazard ratios for all-cause mortality (adjusted for age and sex); analytes with $p < 0.05$ shown. **c)** True Positive Rate vs False Positive Rate. True positive rate versus false positive rate for clinical outcome prediction across anemia, chronic kidney disease, all-cause mortality, and type 2 diabetes, comparing Per_{RI} and NORMA_{RI}. **d)** Performance Improvement with NORMA. Change in precision, sensitivity, specificity, and balanced accuracy (NORMA_{RI} versus Per_{RI}); inner label shows the largest gain. **e)** Lead Time. Median lead time from NORMA_{RI} flag to first Pop_{RI} flag, by analyte (months). Error bars denote the interquartile range.




Fig. 5 | Reclassification performance and early risk stratification in the eICU Collaborative Research Database. a) Reclassification Rate. Reclassification rate by analyte among Pop_{RI}-normal tests. **b)** Hazard Ratios. Cox hazard ratios for in-hospital mortality (adjusted for age and sex); analytes with $p < 0.05$ shown. **c)** True Positive Rate vs False Positive Rate. True positive rate versus false positive rate for clinical outcome prediction across in-hospital mortality, acute kidney injury, sepsis, and prolonged ICU stay, comparing Per_{RI} and NORMA_{RI}. **d)** Performance Improvement with NORMA. Change in precision, sensitivity, specificity, and balanced accuracy (NORMA_{RI} versus Per_{RI}); inner label shows the largest gain. **e)** Lead Time. Median lead time from NORMA_{RI} flag to first Pop_{RI} flag, by analyte (hours). Error bars denote the interquartile range.




Fig. 6 | Reclassification performance and early risk stratification in the INSPIRE cohort. a) Reclassification Rate. Reclassification rate by analyte among Pop_{RI}-normal tests. **b)** Hazard Ratios. Cox hazard ratios for in-hospital mortality (adjusted for age and sex); analytes with $p < 0.05$ shown. **c)** True Positive Rate vs False Positive Rate. True positive rate versus false positive rate for clinical outcome prediction across in-hospital mortality, perioperative infection, prolonged hospital stay, and unplanned ICU admission, comparing Per_{RI} and NORMA_{RI}. **d)** Performance Improvement with NORMA. Change in precision, sensitivity, specificity, and balanced accuracy (NORMA_{RI} versus Per_{RI}); inner label shows the largest gain. **e)** Lead Time. Median lead time from NORMA_{RI} flag to first Pop_{RI} flag, by analyte (hours). Error bars denote the interquartile range.

Supplementary Figures


a Forecasting Performance




b Per-Analyte Performance




c Sensitivity Analysis




Supplementary Fig. 1 | NORMA forecasting performance with Gaussian loss. a) Forecasting Performance. Held-out test-set accuracy (mean absolute error, mean absolute percentage error, and coefficient of determination) for the Gaussian variant versus ARIMA, population mean, and last-value-carried baselines. Error bars denote 95% bootstrap confidence intervals. **b)** Per-Analyte Performance. Per-analyte coefficient of determination grouped by clinical panel. **c)** Sensitivity Analysis. Sensitivity of Gaussian-head prediction interval width and midpoint deviation (each as a percentage of Pop_{RI} width) to history length, forecast horizon, and within-person variability.




Supplementary Fig. 2 | Per-analyte forecasting performance. NORMA using quantile and Gaussian loss, compared to ARIMA, the best performing baseline. Color intensity encodes mean absolute error (MAE), mean absolute percentage error (MAPE), and coefficient of determination (R^2) on the held-out test set.



Supplementary Fig. 3 | Association of absolute deviation from the personalized baseline with observed mortality. Each panel shows the observed mortality rate by standardized absolute deviation from the personalized baseline, binned by decile, for Clalit Health Services, the eICU Collaborative Research Database, and the INSPIRE cohort.




Supplementary Fig. 4 | Age-dependent NORMA_{RI} midpoints. Each panel shows individual midpoint predictions with a polynomial smooth overlaid, for Clalit Health Services, the eICU Collaborative Research Database, and the INSPIRE cohort.




Supplementary Fig. 5 | Abnormality prevalence by analyte and reference interval method. Grouped bars show the percentage of measurements classified as abnormal by Pop_{RI}, Per_{RI}, and NORMA_{RI}, for the eICU Collaborative Research Database, Clalit Health Services, and the INSPIRE cohort.

a Hazard Ratios Across All Outcomes




b Concordance Index Comparison




Supplementary Fig. 6 | Proportional hazards analysis in Clalit Health Services. a) Hazard Ratios Across All Outcomes. Cox hazard ratios for chronic kidney disease, all-cause mortality, type 2 diabetes, and anemia, comparing Pop_{RI} , Per_{RI} , and $NORMA_{RI}$ abnormality flags; only analytes with $p < 0.05$ are shown. **b)** Concordance Index Comparison. Concordance index for the top 10 analytes ranked by $NORMA_{RI}$ concordance per outcome.

a Hazard Ratios Across All Outcomes




b Concordance Index Comparison




Supplementary Fig. 7 | Proportional hazards analysis in the eICU Collaborative Research Database. a) Hazard Ratios Across All Outcomes. Cox hazard ratios for acute kidney injury, in-hospital mortality, prolonged ICU stay, and sepsis, comparing Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags; only analytes with $p < 0.05$ are shown. **b) Concordance Index Comparison.** Concordance index for the top 10 analytes ranked by NORMA_{RI} concordance per outcome.

a Hazard Ratios Across All Outcomes




b Concordance Index Comparison



Supplementary Fig. 8 | Proportional hazards analysis in the INSPIRE cohort. a) Hazard Ratios Across All Outcomes. Cox hazard ratios for in-hospital mortality, perioperative infection, prolonged hospital stay, and unplanned ICU admission, comparing Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags; only analytes with $p < 0.05$ are shown. **b)** Concordance Index Comparison. Concordance index for the top 10 analytes ranked by NORMA_{RI} concordance per outcome.



Supplementary Fig. 9 | Clinical outcome prediction performance in Clalit Health Services. Each point shows Per_{RI} and $NORMA_{RI}$ values for precision, sensitivity, specificity, and balanced accuracy across outcomes. Connected pairs show the direction and magnitude of change between methods for each analyte. Analytes with fewer than 100 measurements are omitted.



Supplementary Fig. 10 | Clinical outcome prediction performance in the eICU Collaborative Research Database. Each point shows Per_{RI} and $NORMA_{RI}$ values for precision, sensitivity, specificity, and balanced accuracy across outcomes. Connected pairs show the direction and magnitude of change between methods for each analyte. Analytes with fewer than 100 measurements are omitted.



Supplementary Fig. 11 | Clinical outcome prediction performance in the INSPIRE cohort. Each point shows Per_{RI} and $NORMA_{RI}$ values for precision, sensitivity, specificity, and balanced accuracy across outcomes. Connected pairs show the direction and magnitude of change between methods for each analyte. Analytes with fewer than 100 measurements are omitted.

Supplementary Tables

Supplementary Table 1 | Analyte reference information. Full analyte name, abbreviation, unit of measurement, and conventional Pop_{RI} for all 30 laboratory analytes included in the study. Pop_{RI} were obtained from the American Board of Internal Medicine Laboratory Test Reference Ranges (January 2025).

Analyte	Full Name	Unit	Population RI
A1C	Hemoglobin A1c	%	4.0–5.6
ALB	Albumin	g/dL	3.5–5.5
ALP	Alkaline Phosphatase	U/L	30–120
ALT	Alanine Aminotransferase	U/L	0–35
AST	Aspartate Aminotransferase	U/L	0–35
BUN	Blood Urea Nitrogen	mg/dL	8–20
CA	Calcium	mg/dL	8.6–10.2
CL	Chloride	mEq/L	98–106
CO2	Bicarbonate	mEq/L	23–28
CRE	Creatinine	mg/dL	F: 0.5–1.1, M: 0.7–1.3
DBIL	Direct Bilirubin	mg/dL	0.1–0.3
GLU	Glucose	mg/dL	70–99
HCT	Hematocrit	%	F: 37–47, M: 42–50
HDL	HDL Cholesterol	mg/dL	40–100
HGB	Hemoglobin	g/dL	F: 12.0–16.0, M: 14.0–18.0
K	Potassium	mEq/L	3.5–5.0
LDL	LDL Cholesterol	mg/dL	<130
MCH	Mean Corpuscular Hemoglobin	pg	28–32
MCHC	MCH Concentration	g/dL	33–36
MCV	Mean Corpuscular Volume	fL	80–98
MPV	Mean Platelet Volume	fL	7.5–12.5
NA	Sodium	mEq/L	136–145
PLT	Platelet Count	10 ³ /μL	150–450
RBC	Red Blood Cell Count	10 ⁶ /μL	F: 4.0–5.2, M: 4.5–5.9
RDW	Red Cell Distribution Width	%	9–14.5
TBIL	Total Bilirubin	mg/dL	0.3–1.0
TC	Total Cholesterol	mg/dL	100–200
TGL	Triglycerides	mg/dL	<150
TP	Total Protein	g/dL	6.0–8.0
WBC	White Blood Cell Count	10 ³ /μL	4.5–11.0

Supplementary Table 2 | Cohort characteristics. Patient counts, demographics (age, sex, follow-up duration), and per-analyte measurement counts and mean values ($\pm$ standard deviation) across all five cohorts: EHRSHOT, MIMIC-IV, Clalit Health Services, the eICU Collaborative Research Database, and the INSPIRE cohort.

	EHRSHOT		MIMIC-IV		eICU-CRD		CHS		INSPIRE	
<i>Patients</i>	5,676		179,601		98,432		1,450,862		51,159	
<i>Age</i>	54.9 $\pm$ 17.2		58.2 $\pm$ 19.0		63.3 $\pm$ 16.1		55.1 $\pm$ 18.6		57.6 $\pm$ 14.9	
<i>Sex</i>	52% F / 48% M		54% F / 46% M		46% F / 54% M		63% F / 37% M		52% F / 48% M	
<i>Time</i>	2.49 [0.28, 7.20] yr		1.01 [0.06, 4.08] yr		0.02 [0.01, 0.03] yr		20.24 [17.51, 21.47] yr		0.48 [0.17, 0.84] yr	
Analyte	N	Value	N	Value	N	Value	N	Value	N	Value
A1C	1,745	6.3 $\pm$ 1.2	39,996	6.6 $\pm$ 1.3	—	—	336,493	7.1 $\pm$ 1.3	391	6.7 $\pm$ 1.0
ALB	4,282	3.3 $\pm$ 0.8	55,220	3.8 $\pm$ 0.8	19,075	2.6 $\pm$ 0.7	790,253	4.1 $\pm$ 0.5	30,728	3.4 $\pm$ 0.6
ALP	4,206	112.5 $\pm$ 58.8	78,930	107.0 $\pm$ 58.4	13,582	97.0 $\pm$ 43.2	1,099,044	82.1 $\pm$ 34.3	27,797	73.4 $\pm$ 29.1
ALT	4,219	37.8 $\pm$ 25.2	92,599	30.2 $\pm$ 21.9	12,425	33.8 $\pm$ 23.4	1,233,902	21.9 $\pm$ 17.3	26,144	21.0 $\pm$ 12.5
AST	4,220	31.9 $\pm$ 19.5	90,639	32.7 $\pm$ 21.4	12,704	37.1 $\pm$ 25.6	1,236,781	23.6 $\pm$ 16.3	26,374	24.2 $\pm$ 10.0
BUN	5,180	23.0 $\pm$ 15.2	153,329	22.2 $\pm$ 14.3	57,540	25.3 $\pm$ 15.5	1,251,453	17.4 $\pm$ 8.3	29,912	16.1 $\pm$ 8.0
CA	5,162	8.8 $\pm$ 0.7	121,478	8.8 $\pm$ 0.7	57,067	8.3 $\pm$ 0.7	909,036	9.3 $\pm$ 0.5	29,788	8.5 $\pm$ 0.7
CL	5,172	102.4 $\pm$ 5.0	148,304	101.8 $\pm$ 5.2	59,172	103.8 $\pm$ 6.2	31,171	103.1 $\pm$ 3.9	30,952	102.5 $\pm$ 4.8
CO2	5,187	25.9 $\pm$ 3.8	145,386	25.6 $\pm$ 4.1	70,208	25.0 $\pm$ 4.7	299	31.4 $\pm$ 12.1	6,247	24.8 $\pm$ 3.8
CRE	5,161	1.1 $\pm$ 0.5	156,044	1.1 $\pm$ 0.6	54,820	1.1 $\pm$ 0.6	1,310,465	0.9 $\pm$ 0.4	42,255	0.8 $\pm$ 0.2
DBIL	1,429	0.3 $\pm$ 0.3	6,635	0.7 $\pm$ 0.9	1,147	0.6 $\pm$ 0.5	—	—	—	—
GLU	5,150	124.9 $\pm$ 39.9	147,195	118.3 $\pm$ 40.4	90,958	139.6 $\pm$ 44.3	1,351,168	110.0 $\pm$ 36.1	26,168	143.8 $\pm$ 48.4
HCT	5,470	32.2 $\pm$ 6.8	164,903	33.6 $\pm$ 6.8	60,110	30.1 $\pm$ 5.9	1,407,912	39.1 $\pm$ 5.2	39,739	32.9 $\pm$ 5.8
HDL	1,549	53.6 $\pm$ 17.5	49,011	55.8 $\pm$ 17.7	11	26.0 $\pm$ 14.3	1,002,204	48.0 $\pm$ 15.9	—	—
HGB	5,369	10.7 $\pm$ 2.3	162,168	11.0 $\pm$ 2.3	60,482	9.9 $\pm$ 2.0	1,408,118	12.8 $\pm$ 1.8	36,187	11.1 $\pm$ 2.0
K	5,236	4.1 $\pm$ 0.5	150,333	4.2 $\pm$ 0.5	66,569	4.0 $\pm$ 0.6	1,182,544	4.5 $\pm$ 0.5	36,363	4.0 $\pm$ 0.5
LDL	821	94.9 $\pm$ 34.9	48,688	104.2 $\pm$ 37.8	3	92.1 $\pm$ 56.2	1,187,844	105.1 $\pm$ 33.6	—	—
MCH	5,294	30.4 $\pm$ 2.7	161,671	29.7 $\pm$ 2.8	48,432	29.7 $\pm$ 2.1	1,250,759	28.8 $\pm$ 2.3	—	—
MCHC	5,317	33.2 $\pm$ 1.4	161,806	32.6 $\pm$ 1.6	51,710	32.8 $\pm$ 1.3	1,407,568	32.8 $\pm$ 1.3	—	—
MCV	5,304	91.5 $\pm$ 6.9	161,661	91.2 $\pm$ 7.3	50,841	90.4 $\pm$ 5.7	1,400,648	87.7 $\pm$ 6.1	—	—
MPV	371	9.6 $\pm$ 1.6	—	—	37,083	9.7 $\pm$ 1.4	1,304,857	9.5 $\pm$ 1.4	—	—
NA	5,193	137.4 $\pm$ 4.2	149,303	138.7 $\pm$ 4.2	62,419	138.6 $\pm$ 5.0	1,187,537	139.9 $\pm$ 3.0	36,127	137.4 $\pm$ 3.5
PLT	5,320	198.0 $\pm$ 111.4	162,308	229.4 $\pm$ 113.3	53,371	199.0 $\pm$ 98.4	1,407,716	243.4 $\pm$ 73.6	34,006	206.0 $\pm$ 92.3
RBC	5,312	3.5 $\pm$ 0.8	161,825	3.7 $\pm$ 0.8	53,584	3.4 $\pm$ 0.7	1,404,794	4.5 $\pm$ 0.6	—	—
RDW	5,315	16.1 $\pm$ 2.9	161,640	15.1 $\pm$ 2.4	48,715	15.8 $\pm$ 2.0	1,403,339	14.1 $\pm$ 1.5	—	—
TBIL	4,169	0.6 $\pm$ 0.4	73,002	0.7 $\pm$ 0.6	12,738	0.8 $\pm$ 0.5	815,774	0.6 $\pm$ 0.4	28,571	0.8 $\pm$ 0.4
TC	1,635	173.5 $\pm$ 45.0	49,762	187.5 $\pm$ 44.6	35	108.3 $\pm$ 44.0	1,260,730	181.7 $\pm$ 40.7	—	—
TGL	4	165.7 $\pm$ 100.8	48,445	140.4 $\pm$ 77.5	312	144.9 $\pm$ 68.0	1,244,518	133.5 $\pm$ 65.3	—	—
TP	4,213	6.6 $\pm$ 1.0	18,673	7.0 $\pm$ 0.7	15,134	5.8 $\pm$ 1.0	791,481	7.1 $\pm$ 0.6	28,603	6.1 $\pm$ 0.9
WBC	5,345	7.8 $\pm$ 4.6	164,291	7.7 $\pm$ 4.0	52,141	10.6 $\pm$ 4.6	1,409,027	7.5 $\pm$ 3.7	34,273	8.2 $\pm$ 3.5

Supplementary Table 3 | NORMA architectural and training design choices. Summary of configurable components including temporal encoding, health state encoding, age and laboratory value representations, output parameterization, and context token usage.

Component	Design Choice	Description
Time embedding	Time2Vec	Encodes the number of days elapsed since the first measurement using learned linear and sinusoidal functions. Captures long-term temporal trends and potential periodic patterns.
	Delta- t	Encodes the log-compressed time difference between consecutive measurements. Explicitly represents irregular sampling and emphasizes how much time has passed since the last value.
Health state encoding	Binary	Encodes each lab value as either within or outside the population reference range; does not distinguish high vs. low and can struggle with J-shaped biomarkers.
	Ternary	Encodes whether the lab value is low, normal, or high relative to the reference range; handles asymmetric risk patterns; however, can increase overfitting risk.
Age encoding	Raw	Uses age directly as a numeric input; assumes linear effects.
	Binned	Discretizes age into groups (e.g., decades) and maps each bin to a learned embedding vector; captures nonlinear and threshold effects.
Lab value encoding	Raw	Uses observed biomarker values directly as model input; preserves absolute clinical scale and keeps preprocessing minimal.
	Within-sequence normalization	Applies per-sequence z-scoring with a learnable scale and shift, then denormalizes at output; stabilizes optimization and improves cross-analyte sharing.
Training objective	Gaussian NLL	Minimizes the negative log-likelihood of a normal distribution by predicting mean and log-variance; aligns with conventional reference interval assumptions but may underperform for skewed or heavy-tailed analytes.
	Quantile (pinball) loss	Minimizes pinball loss for fixed quantiles (2.5, 25, 50, 75, 97.5%); distribution-free and robust to skewness or heteroscedasticity.

Supplementary Table 4 | Within-person and between-person biological variability. Per-analyte within-person coefficient of variation, between-person coefficient of variation, and individuality index for the eICU Collaborative Research Database and Clalit Health Services. Values are reported as median [95% confidence interval].

Analyte	eICU-CRD			CHS			INSPIRE		
	CV _{intra}	CV _{inter}	II	CV _{intra}	CV _{inter}	II	CV _{intra}	CV _{inter}	II
A1C	—	—	—	8.6 [8.5, 8.6]	16.5 [16.5, 16.6]	0.5 [0.5, 0.5]	4.8 [4.4, 5.3]	13.0 [11.8, 13.8]	0.4 [0.3, 0.4]
ALB	8.9 [8.8, 8.9]	22.6 [22.4, 22.7]	0.4 [0.4, 0.4]	6.7 [6.7, 6.8]	18.9 [18.3, 19.4]	0.4 [0.3, 0.4]	8.4 [8.3, 8.4]	13.4 [13.3, 13.5]	0.6 [0.6, 0.6]
ALP	10.6 [10.5, 10.7]	42.1 [41.8, 42.4]	0.2 [0.2, 0.3]	17.8 [17.7, 17.8]	30.1 [30.0, 30.2]	0.6 [0.6, 0.6]	10.9 [10.7, 11.0]	34.0 [33.7, 34.4]	0.3 [0.3, 0.3]
ALT	15.5 [15.3, 15.7]	65.3 [64.8, 65.8]	0.2 [0.2, 0.2]	34.2 [34.2, 34.3]	50.2 [50.0, 50.5]	0.7 [0.7, 0.7]	19.2 [19.0, 19.4]	49.8 [49.3, 50.3]	0.4 [0.4, 0.4]
AST	18.3 [18.1, 18.5]	63.4 [62.9, 63.8]	0.3 [0.3, 0.3]	24.7 [24.6, 24.7]	35.1 [34.9, 35.4]	0.7 [0.7, 0.7]	14.4 [14.3, 14.6]	32.6 [32.2, 32.9]	0.4 [0.4, 0.5]
BUN	16.5 [16.4, 16.6]	58.7 [58.5, 58.9]	0.3 [0.3, 0.3]	20.5 [20.4, 20.5]	36.2 [36.2, 36.3]	0.6 [0.6, 0.6]	15.4 [15.2, 15.5]	42.0 [41.5, 42.4]	0.4 [0.4, 0.4]
CA	3.8 [3.7, 3.8]	7.2 [7.1, 7.2]	0.5 [0.5, 0.5]	3.5 [3.5, 3.5]	3.7 [3.7, 3.7]	0.9 [0.9, 0.9]	4.4 [4.3, 4.4]	5.6 [5.5, 5.6]	0.8 [0.8, 0.8]
CL	2.0 [2.0, 2.0]	5.1 [5.0, 5.1]	0.4 [0.4, 0.4]	2.4 [2.3, 2.4]	2.3 [2.2, 2.3]	1.0 [1.0, 1.1]	1.9 [1.8, 1.9]	3.3 [3.3, 3.3]	0.6 [0.6, 0.6]
CO2	7.1 [7.1, 7.2]	15.9 [15.9, 16.0]	0.5 [0.5, 0.5]	17.7 [17.3, 18.2]	27.0 [25.9, 28.1]	0.7 [0.6, 0.7]	5.5 [5.4, 5.6]	10.2 [10.1, 10.4]	0.5 [0.5, 0.6]
CRE	13.5 [13.4, 13.5]	47.7 [47.5, 47.9]	0.3 [0.3, 0.3]	12.2 [12.2, 12.2]	34.7 [34.4, 34.9]	0.3 [0.3, 0.4]	11.0 [11.0, 11.1]	25.3 [25.2, 25.5]	0.4 [0.4, 0.4]
DBIL	—	82.8 [81.0, 84.7]	—	—	—	—	—	—	—
GLU	14.8 [14.8, 14.9]	24.8 [24.7, 25.0]	0.6 [0.6, 0.6]	13.7 [13.7, 13.7]	24.0 [23.9, 24.0]	0.6 [0.6, 0.6]	12.8 [12.7, 12.9]	23.7 [23.5, 23.9]	0.5 [0.5, 0.6]
HCT	6.2 [6.2, 6.3]	18.4 [18.3, 18.5]	0.3 [0.3, 0.3]	6.3 [6.3, 6.4]	9.6 [9.6, 9.6]	0.7 [0.7, 0.7]	6.8 [6.7, 6.8]	14.1 [14.0, 14.2]	0.5 [0.5, 0.5]
HDL	7.6 [6.0, 9.5]	39.7 [33.9, 44.8]	0.2 [0.2, 0.2]	24.4 [24.4, 24.5]	30.1 [30.1, 30.2]	0.8 [0.8, 0.8]	—	—	—
HGB	6.7 [6.7, 6.7]	19.1 [19.0, 19.1]	0.3 [0.3, 0.3]	6.3 [6.3, 6.3]	10.5 [10.5, 10.5]	0.6 [0.6, 0.6]	6.9 [6.8, 6.9]	14.6 [14.5, 14.7]	0.5 [0.5, 0.5]
K	7.6 [7.6, 7.6]	10.5 [10.4, 10.5]	0.7 [0.7, 0.7]	7.1 [7.1, 7.1]	6.5 [6.4, 6.5]	1.1 [1.1, 1.1]	7.5 [7.5, 7.6]	8.6 [8.6, 8.7]	0.9 [0.9, 0.9]
LDL	8.6 [6.0, 11.5]	45.6 [38.4, 52.0]	0.2 [0.1, 0.3]	20.6 [20.5, 20.6]	23.7 [23.7, 23.8]	0.9 [0.9, 0.9]	—	—	—
MCH	1.1 [1.1, 1.1]	6.9 [6.9, 7.0]	0.1 [0.1, 0.1]	3.6 [3.6, 3.6]	7.1 [7.1, 7.1]	0.5 [0.5, 0.5]	—	—	—
MCHC	1.3 [1.3, 1.3]	3.6 [3.6, 3.7]	0.3 [0.3, 0.3]	2.9 [2.9, 2.9]	2.7 [2.7, 2.7]	1.1 [1.1, 1.1]	—	—	—
MCV	1.0 [1.0, 1.0]	6.2 [6.2, 6.3]	0.2 [0.2, 0.2]	3.2 [3.2, 3.2]	6.1 [6.1, 6.1]	0.5 [0.5, 0.5]	—	—	—
MPV	2.9 [2.9, 2.9]	12.8 [12.8, 12.9]	0.2 [0.2, 0.2]	10.3 [10.3, 10.3]	10.8 [10.8, 10.8]	0.9 [0.9, 1.0]	—	—	—
NA	1.3 [1.3, 1.3]	2.9 [2.9, 2.9]	0.4 [0.4, 0.4]	1.5 [1.5, 1.5]	1.2 [1.2, 1.2]	1.2 [1.2, 1.2]	1.0 [1.0, 1.0]	1.7 [1.7, 1.7]	0.6 [0.6, 0.6]
PLT	11.8 [11.7, 11.9]	41.6 [41.4, 41.8]	0.3 [0.3, 0.3]	14.1 [14.1, 14.1]	23.8 [23.7, 23.8]	0.6 [0.6, 0.6]	12.0 [11.9, 12.2]	33.5 [33.2, 33.8]	0.4 [0.3, 0.4]
RBC	7.2 [7.2, 7.3]	18.6 [18.5, 18.7]	0.4 [0.4, 0.4]	5.9 [5.9, 5.9]	10.1 [10.1, 10.1]	0.6 [0.6, 0.6]	—	—	—
RDW	2.1 [2.0, 2.1]	12.6 [12.6, 12.7]	0.2 [0.2, 0.2]	5.7 [5.7, 5.7]	7.1 [7.1, 7.1]	0.8 [0.8, 0.8]	—	—	—
TBIL	21.9 [21.8, 22.1]	58.2 [57.7, 58.6]	0.4 [0.4, 0.4]	26.0 [25.9, 26.0]	42.8 [42.6, 43.0]	0.6 [0.6, 0.6]	21.7 [21.6, 21.9]	44.6 [44.2, 45.0]	0.5 [0.5, 0.5]
TC	7.6 [6.6, 8.8]	38.4 [34.9, 41.4]	0.2 [0.2, 0.2]	13.9 [13.9, 14.0]	16.6 [16.6, 16.6]	0.8 [0.8, 0.8]	—	—	—
TGL	16.0 [15.3, 16.6]	43.2 [41.8, 44.6]	0.4 [0.3, 0.4]	—	—	—	—	—	—
TP	6.5 [6.4, 6.5]	14.3 [14.2, 14.4]	0.5 [0.5, 0.5]	4.8 [4.8, 4.8]	5.6 [5.5, 5.6]	0.9 [0.9, 0.9]	7.5 [7.4, 7.5]	10.9 [10.8, 11.0]	0.7 [0.7, 0.7]
WBC	15.6 [15.5, 15.7]	37.5 [37.3, 37.6]	0.4 [0.4, 0.4]	19.3 [19.2, 19.3]	—	0.0 [0.0, 0.7]	18.3 [18.2, 18.5]	31.8 [31.5, 32.0]	0.6 [0.6, 0.6]

Supplementary Table 5 | NORMA forecasting performance. Mean absolute error (MAE), mean absolute percentage error (MAPE), and coefficient of determination (R^2) for NORMA Gaussian, NORMA quantile, ARIMA, population mean, and last-value-carried-forward, reported on training, validation, and held-out test sets. Values are reported as median [interquartile range] across analytes.

Metric		NORMA Gaussian	NORMA Quantile	ARIMA	Mean	Last
MAE	Train	6.0 [3.2, 11.1]	5.7 [2.9, 10.1]	6.8 [2.5, 10.2]	8.4 [4.5, 15.0]	7.4 [3.7, 11.0]
	Val	6.0 [2.6, 10.7]	5.8 [2.9, 8.7]	6.8 [3.6, 10.0]	8.3 [4.9, 13.5]	7.3 [3.8, 10.3]
	Test	6.0 [2.6, 10.2]	5.9 [3.6, 9.1]	6.7 [4.3, 11.9]	8.4 [4.5, 13.6]	7.3 [3.9, 10.7]
MAPE	Train	10.9 [8.8, 14.0]	11.9 [8.6, 14.7]	15.9 [11.3, 20.1]	19.4 [14.8, 25.4]	14.8 [10.9, 19.6]
	Val	11.1 [7.9, 13.2]	12.6 [8.7, 15.6]	15.7 [12.1, 21.5]	19.7 [14.0, 27.4]	15.2 [10.6, 20.1]
	Test	11.1 [8.5, 14.3]	12.3 [7.9, 17.3]	16.9 [11.3, 23.3]	19.7 [13.5, 26.7]	15.2 [10.6, 21.1]
R2	Train	0.76 [0.72, 0.80]	0.70 [0.66, 0.75]	0.62 [0.57, 0.70]	0.46 [0.41, 0.49]	0.56 [0.47, 0.62]
	Val	0.75 [0.69, 0.78]	0.69 [0.66, 0.73]	0.62 [0.53, 0.67]	0.46 [0.43, 0.49]	0.55 [0.49, 0.62]
	Test	0.75 [0.70, 0.78]	0.68 [0.64, 0.72]	0.58 [0.51, 0.64]	0.45 [0.41, 0.50]	0.54 [0.49, 0.61]

Supplementary Table 6 | Sensitivity of NORMA prediction intervals to patient features. Median and interquartile range of prediction interval width change (as a percentage of Pop_{RI} width) and midpoint shift (as a percentage of Pop_{RI} width) in response to history length, forecast horizon, and within-person variability, for both the quantile and Gaussian parameterizations.

Model	Feature	CI Width Change (%)		Midpoint Shift (%)	
		Median	IQR	Median	IQR
Quantile	History Length	4.8	[1.0, 14.4]	0.0	[0.0, 0.0]
Quantile	Prediction Horizon	1.6	[0.1, 11.1]	0.0	[0.0, 0.0]
Quantile	Within-Person Variability	115.2	[111.4, 120.5]	0.7	[0.3, 1.7]
Gaussian	History Length	28.3	[20.6, 41.8]	12.8	[5.6, 20.6]
Gaussian	Prediction Horizon	7.2	[3.0, 9.9]	1.6	[0.8, 3.2]
Gaussian	Within-Person Variability	7.2	[3.7, 12.4]	1.3	[0.3, 3.6]

Supplementary Table 7 | Abnormality prevalence and reclassification rates. Overall percentage of measurements classified as abnormal by Pop_{RI}, Per_{RI}, and NORMA_{RI}, and the reclassification rate (the proportion of Pop_{RI}-normal tests flagged abnormal by Per_{RI} or NORMA_{RI}) in the eICU Collaborative Research Database and Clalit Health Services.

Dataset	N	Pop _{RI} (%)	Per _{RI} (%)	NORMA _{RI} (%)	Per _{RI} RR (%)	NORMA _{RI} RR (%)
eICU-CRD	29	50.2	68.1	55.8	37.3	9.2
CHS	31	29.6	46.8	39.1	27.2	12.0
INSPIRE	19	35.1	57.0	42.5	34.3	14.4

Supplementary Table 8 | Per-analyte reclassification. Among tests with a Per_{RI} setpoint within Pop_{RI} , the number per 1,000 flagged abnormal by Pop_{RI} , Per_{RI} , and $NORMA_{RI}$ for each analyte, in the eICU Collaborative Research Database and Clalit Health Services.

Analyte	CHS			eICU-CRD			INSPIRE		
	Pop_{RI}	Per_{RI}	$NORMA_{RI}$	Pop_{RI}	Per_{RI}	$NORMA_{RI}$	Pop_{RI}	Per_{RI}	$NORMA_{RI}$
A1C	240	315	282	—	—	—	278	472	333
ALB	146	284	244	466	551	504	121	284	151
ALP	119	435	218	224	535	294	123	534	225
ALT	136	414	210	252	531	327	225	492	326
AST	71	383	142	186	441	240	45	396	118
BUN	235	458	267	360	546	400	254	479	285
CA	140	290	199	245	378	300	194	345	220
CL	229	415	282	322	461	360	274	430	310
CO2	545	602	562	428	512	467	475	577	562
CRE	141	256	191	231	333	266	123	234	160
DBIL	—	—	—	136	215	172	—	—	—
GLU	326	517	350	541	586	556	619	696	719
HCT	322	497	377	535	677	566	304	517	351
HDL	157	445	217	—	1,000	—	—	—	—
HGB	279	424	331	518	643	555	282	478	315
K	122	250	186	148	281	200	145	331	229
LDL	118	414	195	—	800	—	—	—	—
MCH	184	363	290	83	308	177	—	—	—
MCHC	400	498	439	289	452	339	—	—	—
MCV	94	375	266	48	372	141	—	—	—
MPV	76	314	880	44	320	616	—	—	—
NA	111	361	150	246	418	300	160	409	230
PLT	75	412	125	122	524	310	88	540	312
RBC	240	486	291	475	570	509	—	—	—
RDW	234	378	304	248	456	312	—	—	—
TBIL	119	220	489	149	293	200	72	266	138
TC	184	443	240	286	743	400	—	—	—
TGL	—	—	—	212	519	285	—	—	—
TP	102	273	220	277	427	329	120	310	150
WBC	146	377	228	256	435	298	230	395	266

Supplementary Table 9 | In-hospital mortality prediction performance in the eICU Collaborative Research Database. Per-analyte precision, sensitivity, specificity, and balanced accuracy for Per_{RI} and NORMA_{RI} abnormality flags, restricted to measurements classified as normal by Pop_{RI}. Analytes with fewer than 100 measurements are omitted (—).

Analyte	N	Events	Per _{RI}				NORMA _{RI}			
			Precision	Sensitivity	Specificity	Accuracy	Precision	Sensitivity	Specificity	Accuracy
A1C	—	—	—	—	—	—	—	—	—	—
ALB	891	49	0.05	0.20	0.75	0.48	0.08	0.18	0.88	0.53
ALP	8,843	1,143	0.13	0.60	0.42	0.51	0.16	0.21	0.83	0.52
ALT	7,391	913	0.12	0.56	0.43	0.50	0.12	0.20	0.80	0.50
AST	8,035	793	0.10	0.53	0.48	0.50	0.11	0.17	0.85	0.51
BUN	19,881	1,058	0.06	0.48	0.54	0.51	0.06	0.15	0.87	0.51
CA	16,083	1,214	0.08	0.33	0.67	0.50	0.08	0.17	0.83	0.50
CL	26,608	2,149	0.08	0.37	0.62	0.49	0.10	0.16	0.88	0.52
CO2	29,891	2,393	0.06	0.23	0.69	0.46	0.08	0.16	0.83	0.50
CRE	27,847	1,883	0.08	0.31	0.75	0.53	0.10	0.15	0.90	0.53
DBIL	289	49	0.06	0.06	0.80	0.43	0.04	0.02	0.90	0.46
GLU	9,534	696	0.04	0.15	0.74	0.45	0.08	0.10	0.91	0.51
HCT	2,919	254	0.10	0.49	0.58	0.53	0.13	0.18	0.89	0.53
HDL	—	—	—	—	—	—	—	—	—	—
HGB	3,383	264	0.09	0.42	0.64	0.53	0.12	0.20	0.87	0.54
K	57,801	5,680	0.11	0.38	0.66	0.52	0.12	0.20	0.83	0.52
LDL	—	—	—	—	—	—	—	—	—	—
MCH	32,416	3,272	0.11	0.49	0.54	0.51	0.12	0.29	0.76	0.53
MCHC	22,846	1,998	0.09	0.41	0.61	0.51	0.10	0.18	0.85	0.51
MCV	43,792	4,335	0.11	0.61	0.43	0.52	0.14	0.30	0.79	0.54
MPV	34,479	3,564	0.12	0.59	0.48	0.53	0.11	0.91	0.13	0.52
NA	40,769	3,828	0.10	0.44	0.57	0.50	0.10	0.65	0.37	0.51
PLT	33,546	2,580	0.07	0.65	0.31	0.48	0.07	0.37	0.59	0.48
RBC	4,595	355	0.10	0.34	0.73	0.54	0.10	0.15	0.88	0.52
RDW	16,275	889	0.06	0.49	0.57	0.53	0.06	0.17	0.84	0.51
TBIL	8,985	1,041	0.14	0.35	0.70	0.53	0.15	0.16	0.88	0.52
TC	11	2	0.29	1.00	0.44	0.72	0.67	1.00	0.89	0.94
TGL	176	34	0.20	0.56	0.46	0.51	0.22	0.18	0.85	0.51
TP	4,859	486	0.11	0.38	0.67	0.52	0.11	0.16	0.86	0.51
WBC	25,827	1,714	0.07	0.42	0.58	0.50	0.07	0.15	0.87	0.51

Supplementary Table 10 | Acute kidney injury prediction performance in the eICU Collaborative Research Database. Per-analyte precision, sensitivity, specificity, and balanced accuracy for Per_{RI} and NORMA_{RI} abnormality flags, restricted to measurements classified as normal by Pop_{RI}. Analytes with fewer than 100 measurements are omitted (—).

Analyte	N	Events	Per _{RI}				NORMA _{RI}			
			Precision	Sensitivity	Specificity	Accuracy	Precision	Sensitivity	Specificity	Accuracy
A1C	—	—	—	—	—	—	—	—	—	—
ALB	891	130	0.14	0.23	0.75	0.49	0.17	0.15	0.88	0.51
ALP	8,843	1,712	0.19	0.58	0.41	0.50	0.21	0.18	0.83	0.51
ALT	7,391	1,408	0.17	0.51	0.42	0.47	0.18	0.19	0.80	0.49
AST	8,035	1,384	0.17	0.50	0.47	0.49	0.16	0.14	0.85	0.49
BUN	19,881	1,164	0.05	0.38	0.53	0.46	0.05	0.12	0.87	0.49
CA	16,083	2,079	0.13	0.34	0.67	0.51	0.15	0.20	0.83	0.52
CL	26,608	3,839	0.13	0.34	0.62	0.48	0.16	0.14	0.88	0.51
CO2	29,891	3,601	0.09	0.23	0.68	0.45	0.11	0.16	0.83	0.49
CRE	27,847	2,062	0.07	0.23	0.74	0.48	0.05	0.06	0.89	0.48
DBIL	289	78	0.36	0.23	0.85	0.54	0.33	0.10	0.92	0.51
GLU	9,534	1,257	0.11	0.20	0.74	0.47	0.15	0.10	0.92	0.51
HCT	2,919	266	0.10	0.47	0.58	0.53	0.10	0.13	0.89	0.51
HDL	—	—	—	—	—	—	—	—	—	—
HGB	3,383	255	0.09	0.43	0.64	0.54	0.08	0.14	0.87	0.50
K	57,801	8,390	0.14	0.34	0.66	0.50	0.17	0.20	0.83	0.52
LDL	—	—	—	—	—	—	—	—	—	—
MCH	32,416	4,932	0.16	0.48	0.54	0.51	0.17	0.28	0.76	0.52
MCHC	22,846	3,339	0.14	0.37	0.61	0.49	0.14	0.15	0.84	0.50
MCV	43,792	6,744	0.15	0.56	0.43	0.49	0.16	0.23	0.78	0.50
MPV	34,479	5,045	0.15	0.53	0.47	0.50	0.15	0.88	0.13	0.51
NA	40,769	6,154	0.14	0.40	0.56	0.48	0.15	0.61	0.37	0.49
PLT	33,546	4,508	0.13	0.65	0.30	0.48	0.13	0.39	0.59	0.49
RBC	4,595	429	0.11	0.31	0.73	0.52	0.11	0.14	0.88	0.51
RDW	16,275	1,681	0.10	0.42	0.57	0.50	0.11	0.17	0.84	0.51
TBIL	8,985	1,660	0.19	0.31	0.70	0.51	0.20	0.13	0.88	0.51
TC	11	3	0.29	0.67	0.38	0.52	0.33	0.33	0.75	0.54
TGL	176	42	0.21	0.48	0.43	0.46	0.22	0.14	0.84	0.49
TP	4,859	864	0.19	0.36	0.67	0.52	0.21	0.17	0.87	0.52
WBC	25,827	3,491	0.12	0.39	0.57	0.48	0.14	0.13	0.87	0.50

Supplementary Table 11 | Sepsis prediction performance in the eICU Collaborative Research Database. Per-analyte precision, sensitivity, specificity, and balanced accuracy for Per_{RI} and NORMA_{RI} abnormality flags, restricted to measurements classified as normal by Pop_{RI}. Analytes with fewer than 100 measurements are omitted (—).

Analyte	N	Events	Per _{RI}				NORMA _{RI}			
			Precision	Sensitivity	Specificity	Accuracy	Precision	Sensitivity	Specificity	Accuracy
A1C	—	—	—	—	—	—	—	—	—	—
ALB	891	87	0.08	0.20	0.75	0.47	0.13	0.16	0.88	0.52
ALP	8,843	2,106	0.23	0.57	0.41	0.49	0.23	0.16	0.83	0.50
ALT	7,391	1,925	0.25	0.54	0.42	0.48	0.25	0.19	0.80	0.49
AST	8,035	2,036	0.24	0.49	0.47	0.48	0.23	0.14	0.85	0.49
BUN	19,881	2,759	0.12	0.40	0.53	0.47	0.12	0.11	0.87	0.49
CA	16,083	2,034	0.13	0.34	0.67	0.51	0.14	0.19	0.83	0.51
CL	26,608	4,164	0.14	0.34	0.62	0.48	0.17	0.13	0.87	0.50
CO2	29,891	4,190	0.11	0.25	0.68	0.47	0.12	0.14	0.83	0.49
CRE	27,847	4,045	0.14	0.25	0.74	0.50	0.13	0.10	0.90	0.49
DBIL	289	72	0.32	0.22	0.84	0.53	0.33	0.11	0.93	0.52
GLU	9,534	1,557	0.13	0.20	0.74	0.47	0.18	0.10	0.92	0.51
HCT	2,919	332	0.11	0.40	0.57	0.49	0.12	0.12	0.89	0.50
HDL	—	—	—	—	—	—	—	—	—	—
HGB	3,383	363	0.10	0.35	0.63	0.49	0.12	0.15	0.87	0.51
K	57,801	9,268	0.16	0.34	0.65	0.49	0.17	0.18	0.83	0.51
LDL	—	—	—	—	—	—	—	—	—	—
MCH	32,416	5,623	0.17	0.47	0.53	0.50	0.19	0.27	0.76	0.51
MCHC	22,846	3,540	0.15	0.36	0.61	0.49	0.15	0.15	0.84	0.50
MCV	43,792	7,707	0.17	0.57	0.43	0.50	0.19	0.23	0.79	0.51
MPV	34,479	5,542	0.16	0.52	0.47	0.50	0.16	0.88	0.13	0.50
NA	40,769	6,847	0.15	0.40	0.56	0.48	0.17	0.62	0.37	0.50
PLT	33,546	5,676	0.17	0.67	0.31	0.49	0.17	0.41	0.59	0.50
RBC	4,595	594	0.13	0.27	0.73	0.50	0.10	0.09	0.88	0.48
RDW	16,275	2,064	0.12	0.41	0.57	0.49	0.12	0.16	0.84	0.50
TBIL	8,985	2,286	0.26	0.30	0.70	0.50	0.25	0.12	0.88	0.50
TC	11	2	0.29	1.00	0.44	0.72	0.00	0.00	0.67	0.33
TGL	176	56	0.35	0.61	0.48	0.55	0.30	0.14	0.84	0.49
TP	4,859	1,118	0.24	0.35	0.67	0.51	0.26	0.16	0.87	0.51
WBC	25,827	3,531	0.12	0.36	0.57	0.47	0.12	0.11	0.87	0.49

Supplementary Table 12 | Prolonged ICU stay prediction performance in the eICU Collaborative Research Database. Per-analyte precision, sensitivity, specificity, and balanced accuracy for Per_{RI} and NORMA_{RI} abnormality flags, restricted to measurements classified as normal by Pop_{RI}. Analytes with fewer than 100 measurements are omitted (—).

Analyte	N	Events	Per _{RI}				NORMA _{RI}			
			Precision	Sensitivity	Specificity	Accuracy	Precision	Sensitivity	Specificity	Accuracy
A1C	—	—	—	—	—	—	—	—	—	—
ALB	891	158	0.17	0.24	0.75	0.50	0.23	0.16	0.88	0.52
ALP	8,843	2,882	0.32	0.57	0.41	0.49	0.35	0.19	0.84	0.51
ALT	7,391	2,412	0.32	0.55	0.42	0.48	0.33	0.20	0.80	0.50
AST	8,035	2,561	0.30	0.48	0.46	0.47	0.31	0.14	0.85	0.50
BUN	19,881	3,352	0.15	0.41	0.53	0.47	0.19	0.15	0.87	0.51
CA	16,083	2,941	0.20	0.35	0.68	0.51	0.23	0.21	0.84	0.53
CL	26,608	5,167	0.15	0.28	0.60	0.44	0.23	0.15	0.88	0.52
CO2	29,891	5,495	0.12	0.20	0.67	0.43	0.23	0.20	0.84	0.52
CRE	27,847	5,026	0.19	0.27	0.75	0.51	0.19	0.11	0.90	0.50
DBIL	289	137	0.56	0.20	0.85	0.53	0.50	0.09	0.92	0.50
GLU	9,534	718	0.06	0.18	0.75	0.46	0.13	0.15	0.92	0.54
HCT	2,919	467	0.16	0.41	0.57	0.49	0.17	0.12	0.89	0.51
HDL	—	—	—	—	—	—	—	—	—	—
HGB	3,383	486	0.13	0.33	0.63	0.48	0.15	0.14	0.87	0.50
K	57,801	11,154	0.20	0.35	0.66	0.50	0.26	0.23	0.84	0.54
LDL	—	—	—	—	—	—	—	—	—	—
MCH	32,416	7,342	0.25	0.51	0.55	0.53	0.28	0.31	0.77	0.54
MCHC	22,846	4,454	0.18	0.35	0.60	0.48	0.19	0.15	0.84	0.50
MCV	43,792	9,765	0.22	0.57	0.43	0.50	0.27	0.27	0.80	0.53
MPV	34,479	7,956	0.24	0.55	0.48	0.51	0.24	0.90	0.14	0.52
NA	40,769	8,509	0.17	0.36	0.55	0.45	0.21	0.64	0.37	0.51
PLT	33,546	7,568	0.22	0.67	0.30	0.49	0.27	0.48	0.62	0.55
RBC	4,595	778	0.18	0.29	0.73	0.51	0.19	0.14	0.89	0.51
RDW	16,275	2,917	0.19	0.45	0.57	0.51	0.20	0.18	0.84	0.51
TBIL	8,985	2,983	0.35	0.32	0.71	0.51	0.35	0.13	0.88	0.51
TC	11	1	0.00	0.00	0.30	0.15	0.33	1.00	0.80	0.90
TGL	176	117	0.66	0.54	0.44	0.49	0.59	0.14	0.81	0.47
TP	4,859	1,495	0.33	0.36	0.67	0.52	0.37	0.17	0.87	0.52
WBC	25,827	5,040	0.17	0.38	0.57	0.47	0.20	0.13	0.87	0.50

Supplementary Table 13 | In-hospital mortality prediction performance in the INSPIRE cohort. Per-analyte precision, sensitivity, specificity, and balanced accuracy for Per_{RI} and NORMA_{RI} abnormality flags, restricted to measurements classified as normal by Pop_{RI}. Analytes with fewer than 100 measurements are omitted (—).

Analyte	N	Events	Per _{RI}				NORMA _{RI}			
			Precision	Sensitivity	Specificity	Accuracy	Precision	Sensitivity	Specificity	Accuracy
A1C	26	0	0.00	—	0.73	—	0.00	—	0.92	—
ALB	15,833	62	0.00	0.19	0.81	0.50	0.01	0.07	0.96	0.52
ALP	23,817	387	0.02	0.50	0.53	0.51	0.02	0.17	0.89	0.53
ALT	17,229	383	0.02	0.31	0.65	0.48	0.02	0.14	0.87	0.51
AST	24,951	513	0.03	0.45	0.63	0.54	0.03	0.12	0.92	0.52
BUN	18,100	225	0.01	0.32	0.69	0.51	0.03	0.10	0.96	0.53
CA	12,367	197	0.02	0.28	0.82	0.55	0.07	0.12	0.97	0.55
CL	19,435	310	0.01	0.20	0.77	0.48	0.04	0.12	0.95	0.54
CO2	3,345	188	0.04	0.21	0.68	0.44	0.08	0.35	0.76	0.55
CRE	32,702	355	0.03	0.35	0.89	0.62	0.07	0.18	0.97	0.58
DBIL	—	—	—	—	—	—	—	—	—	—
GLU	3,293	58	0.02	0.31	0.78	0.54	0.02	0.28	0.75	0.51
HCT	9,823	45	0.01	0.40	0.69	0.55	0.01	0.20	0.94	0.57
HDL	—	—	—	—	—	—	—	—	—	—
HGB	11,389	61	0.01	0.43	0.73	0.58	0.01	0.07	0.96	0.51
K	31,429	580	0.03	0.39	0.76	0.57	0.07	0.30	0.92	0.61
LDL	—	—	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—	—	—
NA	26,515	367	0.02	0.50	0.69	0.60	0.07	0.33	0.94	0.63
PLT	27,257	260	0.01	0.43	0.49	0.46	0.01	0.28	0.75	0.52
RBC	—	—	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—	—	—
TBIL	23,415	397	0.03	0.38	0.79	0.59	0.02	0.09	0.93	0.51
TC	—	—	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—	—	—
TP	16,931	178	0.01	0.21	0.78	0.49	0.01	0.03	0.96	0.50
WBC	23,905	281	0.01	0.23	0.78	0.51	0.02	0.09	0.95	0.52

Supplementary Table 14 | Perioperative infection prediction performance in the INSPIRE cohort. Per-analyte precision, sensitivity, specificity, and balanced accuracy for Per_{RI} and NORMA_{RI} abnormality flags, restricted to measurements classified as normal by Pop_{RI}. Analytes with fewer than 100 measurements are omitted (—).

Analyte	N	Events	Per _{RI}				NORMA _{RI}			
			Precision	Sensitivity	Specificity	Accuracy	Precision	Sensitivity	Specificity	Accuracy
A1C	26	2	0.14	0.50	0.75	0.62	0.00	0.00	0.92	0.46
ALB	15,833	437	0.03	0.17	0.81	0.49	0.03	0.04	0.96	0.50
ALP	23,817	1,174	0.04	0.37	0.52	0.44	0.04	0.10	0.89	0.49
ALT	17,229	1,005	0.05	0.29	0.64	0.46	0.05	0.12	0.87	0.49
AST	24,951	1,439	0.05	0.33	0.62	0.48	0.06	0.09	0.92	0.50
BUN	18,100	736	0.03	0.23	0.68	0.46	0.02	0.02	0.96	0.49
CA	12,367	481	0.03	0.15	0.81	0.48	0.06	0.04	0.97	0.51
CL	19,435	854	0.03	0.14	0.76	0.45	0.04	0.04	0.95	0.50
CO2	3,345	421	0.09	0.22	0.67	0.44	0.12	0.24	0.75	0.50
CRE	32,702	1,124	0.04	0.14	0.89	0.51	0.05	0.04	0.97	0.51
DBIL	—	—	—	—	—	—	—	—	—	—
GLU	3,293	161	0.04	0.19	0.77	0.48	0.06	0.30	0.75	0.53
HCT	9,823	201	0.02	0.31	0.69	0.50	0.03	0.08	0.94	0.51
HDL	—	—	—	—	—	—	—	—	—	—
HGB	11,389	277	0.02	0.24	0.73	0.49	0.03	0.05	0.96	0.51
K	31,429	1,374	0.04	0.22	0.76	0.49	0.07	0.13	0.92	0.53
LDL	—	—	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—	—	—
NA	26,515	975	0.03	0.28	0.69	0.49	0.07	0.12	0.94	0.53
PLT	27,257	1,137	0.03	0.36	0.49	0.42	0.03	0.20	0.75	0.47
RBC	—	—	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—	—	—
TBIL	23,415	1,292	0.07	0.25	0.79	0.52	0.05	0.07	0.93	0.50
TC	—	—	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—	—	—
TP	16,931	763	0.04	0.19	0.78	0.48	0.08	0.06	0.97	0.51
WBC	23,905	1,043	0.03	0.17	0.78	0.48	0.04	0.04	0.95	0.50

Supplementary Table 15 | Prolonged hospital stay prediction performance in the INSPIRE cohort. Per-analyte precision, sensitivity, specificity, and balanced accuracy for Per_{RI} and NORMA_{RI} abnormality flags, restricted to measurements classified as normal by Pop_{RI}. Analytes with fewer than 100 measurements are omitted (—).

Analyte	N	Events	Per _{RI}				NORMA _{RI}			
			Precision	Sensitivity	Specificity	Accuracy	Precision	Sensitivity	Specificity	Accuracy
A1C	26	19	0.71	0.26	0.71	0.49	1.00	0.10	1.00	0.55
ALB	15,833	10,599	0.68	0.19	0.81	0.50	0.66	0.03	0.96	0.50
ALP	23,817	18,548	0.78	0.48	0.53	0.50	0.70	0.10	0.85	0.47
ALT	17,229	13,384	0.76	0.34	0.61	0.48	0.74	0.13	0.84	0.48
AST	24,951	19,532	0.76	0.36	0.59	0.48	0.72	0.07	0.90	0.48
BUN	18,100	13,840	0.74	0.30	0.65	0.48	0.70	0.04	0.95	0.49
CA	12,367	8,496	0.67	0.18	0.81	0.49	0.65	0.03	0.97	0.50
CL	19,435	15,298	0.74	0.22	0.72	0.47	0.73	0.04	0.94	0.49
CO2	3,345	2,885	0.83	0.31	0.62	0.46	0.89	0.25	0.80	0.53
CRE	32,702	22,812	0.70	0.12	0.89	0.50	0.67	0.03	0.97	0.50
DBIL	—	—	—	—	—	—	—	—	—	—
GLU	3,293	2,352	0.67	0.21	0.74	0.48	0.74	0.26	0.78	0.52
HCT	9,823	5,503	0.52	0.29	0.66	0.47	0.49	0.06	0.93	0.49
HDL	—	—	—	—	—	—	—	—	—	—
HGB	11,389	6,945	0.60	0.27	0.72	0.49	0.53	0.04	0.95	0.49
K	31,429	23,052	0.71	0.24	0.73	0.48	0.72	0.08	0.92	0.50
LDL	—	—	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—	—	—
NA	26,515	18,858	0.69	0.30	0.67	0.48	0.68	0.06	0.93	0.49
PLT	27,257	19,942	0.71	0.50	0.46	0.48	0.68	0.23	0.71	0.47
RBC	—	—	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—	—	—
TBIL	23,415	17,682	0.75	0.21	0.78	0.50	0.70	0.07	0.91	0.49
TC	—	—	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—	—	—
TP	16,931	12,395	0.73	0.22	0.78	0.50	0.74	0.04	0.97	0.50
WBC	23,905	17,474	0.69	0.21	0.75	0.48	0.76	0.05	0.96	0.50

Supplementary Table 16 | Unplanned ICU admission prediction performance in the INSPIRE cohort. Per-analyte precision, sensitivity, specificity, and balanced accuracy for Per_{RI} and NORMA_{RI} abnormality flags, restricted to measurements classified as normal by Pop_{RI}. Analytes with fewer than 100 measurements are omitted (—).

Analyte	N	Events	Per _{RI}				NORMA _{RI}			
			Precision	Sensitivity	Specificity	Accuracy	Precision	Sensitivity	Specificity	Accuracy
A1C	26	15	0.43	0.20	0.64	0.42	1.00	0.13	1.00	0.57
ALB	15,833	4,067	0.22	0.16	0.80	0.48	0.20	0.03	0.96	0.49
ALP	23,817	7,129	0.28	0.45	0.52	0.48	0.26	0.10	0.88	0.49
ALT	17,229	5,695	0.31	0.34	0.64	0.49	0.35	0.14	0.87	0.51
AST	24,951	7,856	0.27	0.32	0.60	0.46	0.25	0.06	0.92	0.49
BUN	18,100	5,436	0.29	0.30	0.68	0.49	0.33	0.04	0.96	0.50
CA	12,367	3,418	0.23	0.15	0.81	0.48	0.20	0.02	0.97	0.49
CL	19,435	5,038	0.19	0.17	0.74	0.46	0.34	0.06	0.96	0.51
CO2	3,345	1,870	0.51	0.29	0.65	0.47	0.57	0.25	0.76	0.51
CRE	32,702	6,625	0.21	0.12	0.89	0.50	0.19	0.03	0.97	0.50
DBIL	—	—	—	—	—	—	—	—	—	—
GLU	3,293	622	0.14	0.16	0.76	0.46	0.22	0.29	0.76	0.52
HCT	9,823	1,745	0.15	0.27	0.68	0.47	0.25	0.09	0.94	0.52
HDL	—	—	—	—	—	—	—	—	—	—
HGB	11,389	2,448	0.21	0.27	0.73	0.50	0.24	0.05	0.96	0.50
K	31,429	8,519	0.25	0.22	0.75	0.48	0.32	0.09	0.93	0.51
LDL	—	—	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—	—	—
NA	26,515	6,473	0.20	0.26	0.68	0.47	0.32	0.09	0.94	0.51
PLT	27,257	6,866	0.20	0.41	0.46	0.44	0.22	0.21	0.74	0.48
RBC	—	—	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—	—	—
TBIL	23,415	6,578	0.28	0.21	0.79	0.50	0.27	0.07	0.93	0.50
TC	—	—	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—	—	—
TP	16,931	4,750	0.23	0.18	0.76	0.47	0.22	0.03	0.96	0.49
WBC	23,905	6,052	0.19	0.17	0.76	0.47	0.23	0.04	0.95	0.50

Supplementary Table 17 | Positive predictive value across outcomes in Clalit Health Services. Measurements classified as normal by Pop_{RI} are excluded. Per 100 patients flagged abnormal by Per_{RI} or NORMA_{RI}, the number who experienced each outcome (anemia, chronic kidney disease, all-cause mortality, and type 2 diabetes), by analyte. Analytes with fewer than 100 measurements are omitted (—).

Analyte	Anemia		Chronic Kidney Disease		Mortality		Type 2 Diabetes	
	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}
A1C	21	30	39	48	23	34	70	78
ALB	21	25	44	51	26	33	78	81
ALP	19	24	40	47	25	30	74	79
ALT	19	22	40	43	24	27	74	78
AST	19	23	40	45	26	29	74	79
BUN	14	20	26	34	18	24	73	80
CA	21	27	45	53	28	35	77	82
CL	30	38	52	62	36	46	93	97
CO2	100	100	75	100	50	80	100	100
CRE	16	24	34	57	21	32	75	84
DBIL	—	—	—	—	—	—	—	—
GLU	16	23	30	38	18	26	53	64
HCT	10	12	28	34	15	19	70	75
HDL	17	21	38	42	22	26	71	74
HGB	9	12	28	33	15	19	70	76
K	21	25	41	48	27	34	78	83
LDL	20	24	43	50	26	30	79	86
MCH	16	18	38	41	23	27	71	74
MCHC	14	17	30	34	18	22	73	77
MCV	17	20	37	40	23	26	73	75
MPV	19	18	37	36	24	24	74	73
NA	19	26	41	51	26	34	76	84
PLT	18	25	35	45	23	32	73	79
RBC	13	18	30	36	17	22	71	78
RDW	14	15	32	33	18	19	71	72
TBIL	24	22	46	44	30	29	79	77
TC	21	26	43	50	27	31	80	87
TGL	—	—	—	—	—	—	—	—
TP	23	27	47	54	30	37	78	81
WBC	18	22	37	41	23	27	73	78
Overall	21	25	39	46	24	30	75	80

Supplementary Table 18 | Positive predictive value across outcomes in the eICU Collaborative Research Database. Measurements classified as normal by Pop_{RI} are excluded. Per 100 patients flagged abnormal by Per_{RI} or NORMA_{RI}, the number who experienced each outcome (acute kidney injury, in-hospital mortality, prolonged ICU stay, and sepsis), by analyte. Analytes with fewer than 100 measurements are omitted (—).

Analyte	Acute Kidney Injury		Mortality		Prolonged LOS (>7d)		Sepsis	
	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}
A1C	—	—	—	—	—	—	—	—
ALB	14	17	5	8	17	23	8	13
ALP	19	21	13	16	32	35	23	23
ALT	17	18	12	12	32	33	25	25
AST	16	16	10	11	30	31	24	23
BUN	5	5	6	6	15	19	12	12
CA	13	15	8	8	20	23	13	14
CL	13	16	8	10	15	23	14	16
CO2	9	11	6	8	12	23	11	12
CRE	7	5	8	10	19	19	14	13
DBIL	36	33	6	4	56	50	32	33
GLU	11	16	4	8	6	13	13	18
HCT	10	10	10	13	16	17	11	12
HDL	—	—	—	—	—	—	—	—
HGB	9	8	9	12	13	15	10	12
K	14	17	11	12	20	26	16	17
LDL	—	—	—	—	—	—	—	—
MCH	16	17	11	12	25	28	18	19
MCHC	14	14	9	10	18	19	15	15
MCV	15	16	11	14	22	27	18	19
MPV	15	15	12	11	24	24	16	16
NA	14	15	10	10	17	21	15	17
PLT	13	13	7	7	22	27	16	17
RBC	11	11	10	10	18	19	13	10
RDW	10	11	6	6	19	20	12	12
TBIL	19	20	14	15	35	35	26	25
TC	29	33	29	67	0	33	29	0
TGL	21	22	20	22	66	59	35	30
TP	19	22	11	11	33	37	24	26
WBC	12	14	7	8	17	20	12	12
Overall	15	16	10	13	23	27	18	17

Supplementary Table 19 | Positive predictive value across outcomes in the INSPIRE cohort. Measurements classified as normal by Pop_{RI} are excluded. Per 100 patients flagged abnormal by Per_{RI} or NORMA_{RI}, the number who experienced each outcome (in-hospital mortality, perioperative infection, prolonged hospital stay, and unplanned ICU admission), by analyte. Analytes with fewer than 100 measurements are omitted (—).

Analyte	Mortality		Perioperative Infection		Prolonged LOS (>7d)		Unplanned ICU Admission	
	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}	Per _{RI}	NORMA _{RI}
A1C	0	0	14	0	71	100	43	100
ALB	0	1	2	3	68	66	22	20
ALP	2	2	4	4	78	70	28	26
ALT	2	2	5	5	76	74	31	35
AST	2	3	5	6	76	72	27	25
BUN	1	3	3	2	74	70	29	33
CA	2	7	3	6	67	65	23	20
CL	1	4	3	4	74	73	19	34
CO2	4	8	9	12	83	89	51	57
CRE	3	7	4	5	70	67	21	19
DBIL	—	—	—	—	—	—	—	—
GLU	2	2	4	6	67	74	14	22
HCT	1	1	2	2	52	49	15	25
HDL	—	—	—	—	—	—	—	—
HGB	1	1	2	3	60	52	21	24
K	3	7	4	7	71	72	25	32
LDL	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—
NA	2	7	3	7	69	68	20	32
PLT	1	1	3	3	71	68	20	22
RBC	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—
TBIL	3	2	7	5	75	70	28	27
TC	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—
TP	1	1	4	8	73	74	22	22
WBC	1	2	3	4	69	76	19	23
Overall	2	3	4	5	71	71	25	31

Supplementary Table 20 | Lead time for NORMA_{RI} abnormality detection relative to Pop_{RI}. Per analyte and cohort: number of tests flagged by NORMA_{RI} alone (NORMA_{RI}-only) and the subset later confirmed by Pop_{RI} (later population), with median lead time and interquartile range (hours for eICU-CRD and INSPIRE, months for CHS).

Analyte	eICU-CRD				CHS				INSPIRE			
	NORMA-only	Later Pop _{RI}	Median (h)	IQR (h)	NORMA-only	Later Pop _{RI}	Median (mo)	IQR (mo)	NORMA-only	Later Pop _{RI}	Median (h)	IQR (h)
A1C	—	—	—	—	14,227	5,064	12.4	5.7–27.5	24	0	—	—
ALB	2,185	865	30.0	23.1–73.9	807,619	436,471	7.9	1.2–29.9	1,493	29	23.5	13.3–25.7
ALP	4,362	757	49.6	24.2–120.2	1,196,400	543,146	15.5	3.1–50.6	3,442	201	60.4	24.7–102.3
ALT	6,749	1,235	47.6	23.9–99.4	1,199,702	703,509	11.9	2.4–41.0	3,210	187	47.6	19.2–109.0
AST	7,720	1,317	47.4	23.9–95.5	1,195,620	551,419	10.0	1.6–38.0	2,204	29	16.6	8.2–24.0
BUN	25,578	7,039	28.0	23.1–71.3	568,660	402,995	7.6	1.1–29.2	1,983	409	24.9	13.0–55.2
CA	45,543	13,222	25.1	19.6–69.2	574,875	356,694	6.4	0.8–29.0	1,419	342	39.8	16.3–77.0
CL	24,313	8,477	34.7	23.0–72.8	44,929	36,351	4.9	0.5–22.7	3,124	932	24.0	11.4–54.2
CO2	52,641	26,365	25.6	19.3–64.5	96	80	2.9	0.3–8.7	3,787	2,418	8.4	3.5–19.2
CRE	40,464	8,744	26.0	21.6–67.7	1,070,918	687,844	7.5	1.4–27.1	2,321	826	43.2	16.1–121.8
DBIL	425	72	25.2	20.8–49.3	—	—	—	—	—	—	—	—
GLU	20,333	17,921	5.6	3.2–12.1	358,479	248,307	5.5	0.9–20.6	14,216	8,944	4.6	2.5–8.7
HCT	2,406	998	24.2	17.7–48.1	1,083,585	762,204	8.5	1.8–30.2	2,240	484	10.4	3.4–21.5
HDL	2	0	—	—	515,071	120,646	21.9	7.3–53.3	—	—	—	—
HGB	4,408	1,526	23.7	14.3–47.1	1,065,707	585,178	16.1	3.4–44.3	1,576	301	14.8	5.7–32.6
K	62,211	16,970	40.9	23.0–78.6	1,337,595	881,518	4.5	0.4–25.8	5,722	2,611	15.7	5.8–37.2
LDL	3	2	35.1	29.2–41.0	764,300	172,816	17.6	6.4–44.5	—	—	—	—
MCH	40,943	5,361	45.5	23.4–93.1	2,300,580	1,420,584	11.2	2.8–37.1	—	—	—	—
MCHC	38,761	11,733	25.1	22.5–49.9	840,255	776,047	7.6	1.6–29.9	—	—	—	—
MCV	19,132	1,789	48.2	23.7–120.7	4,464,421	1,941,232	18.8	4.1–62.8	—	—	—	—
MPV	101,702	4,805	37.5	23.2–72.5	20,300,446	8,370,452	20.0	4.8–53.3	—	—	—	—
NA	115,529	31,882	26.6	21.1–69.9	805,446	506,896	8.0	0.9–34.7	4,825	1,308	29.8	11.3–75.0
PLT	46,110	5,482	49.7	24.0–120.8	1,356,487	660,368	8.5	1.1–40.3	8,493	277	23.3	8.9–64.1
RBC	6,592	2,129	23.9	16.5–47.6	1,202,775	769,663	8.9	1.8–32.9	—	—	—	—
RDW	18,706	3,308	26.3	22.8–70.0	1,580,050	933,310	16.2	3.1–52.3	—	—	—	—
TBIL	11,136	1,721	47.4	23.8–95.5	2,966,137	1,268,891	26.4	6.0–68.5	2,742	71	23.8	9.7–83.0
TC	28	7	70.3	42.0–120.9	675,432	330,110	14.8	5.0–39.6	—	—	—	—
TGL	317	38	60.0	25.4–145.5	—	—	—	—	—	—	—	—
TP	11,311	2,592	34.8	22.9–74.1	962,840	530,403	6.9	0.9–28.6	1,227	19	13.0	11.9–24.0
WBC	31,253	8,381	35.3	23.3–72.7	2,183,898	1,402,700	7.4	1.0–33.0	2,389	348	26.0	13.8–57.4
Overall	740,863	184,738	34.8	23.1–72.6	51,436,550	25,404,898	8.7	1.7–33.8	66,437	19,735	23.7	11.4–54.7

Supplementary Table 21 | Proportional hazards analysis for all-cause mortality in Clalit Health Services. Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	9517 (3118)	6345 (2079)	0.84 [0.65, 1.09]	0.689	0.58 [0.54, 0.63]	0.699	0.82 [0.65, 1.03]	0.688
ALB	13612 (4221)	9076 (2815)	4.15 [3.83, 4.49]	0.785	1.21 [1.10, 1.33]	0.740	3.11 [2.80, 3.46]	0.757
ALP	20262 (5615)	13508 (3743)	2.05 [1.94, 2.17]	0.760	0.96 [0.87, 1.05]	0.741	2.01 [1.86, 2.18]	0.749
ALT	25254 (6667)	16836 (4445)	2.08 [1.95, 2.22]	0.761	0.94 [0.86, 1.03]	0.749	1.35 [1.27, 1.43]	0.752
AST	25296 (6662)	16864 (4442)	1.85 [1.76, 1.94]	0.758	1.02 [0.94, 1.12]	0.744	1.55 [1.47, 1.65]	0.750
BUN	26980 (7187)	17987 (4792)	2.82 [2.53, 3.15]	0.747	0.81 [0.75, 0.88]	0.739	2.58 [2.31, 2.87]	0.746
CA	15946 (4688)	10631 (3125)	1.93 [1.81, 2.07]	0.754	1.08 [0.99, 1.17]	0.740	1.88 [1.73, 2.05]	0.749
CL	846 (307)	564 (205)	3.57 [1.67, 7.61]	0.767	1.90 [0.97, 3.71]	0.760	4.10 [1.82, 9.27]	0.755
CO2	—	—	—	—	—	—	—	—
CRE	29331 (7624)	19555 (5083)	3.00 [2.82, 3.18]	0.770	1.04 [0.97, 1.12]	0.744	3.30 [3.07, 3.54]	0.768
DBIL	—	—	—	—	—	—	—	—
GLU	28928 (7343)	19286 (4895)	1.60 [1.39, 1.85]	0.749	0.77 [0.70, 0.84]	0.748	1.61 [1.38, 1.86]	0.749
HCT	31026 (7576)	20685 (5051)	3.60 [3.15, 4.12]	0.774	0.79 [0.73, 0.86]	0.765	3.36 [2.86, 3.94]	0.771
HDL	20800 (5296)	13867 (3530)	1.68 [1.56, 1.81]	0.733	0.73 [0.67, 0.81]	0.726	1.72 [1.57, 1.89]	0.730
HGB	31020 (7578)	20680 (5052)	3.70 [3.32, 4.12]	0.779	0.83 [0.77, 0.90]	0.764	3.72 [3.24, 4.26]	0.772
K	25819 (7121)	17213 (4748)	1.63 [1.53, 1.73]	0.748	1.07 [0.99, 1.15]	0.742	1.69 [1.56, 1.83]	0.746
LDL	23706 (6093)	15804 (4062)	0.81 [0.77, 0.85]	0.735	0.74 [0.69, 0.79]	0.736	0.87 [0.82, 0.92]	0.734
MCH	28147 (6984)	18765 (4656)	2.04 [1.89, 2.20]	0.767	0.98 [0.91, 1.07]	0.760	2.18 [1.90, 2.50]	0.763
MCHC	31009 (7580)	20673 (5053)	1.20 [0.89, 1.61]	0.767	0.96 [0.90, 1.03]	0.767	0.99 [0.76, 1.28]	0.767
MCV	30848 (7546)	20566 (5031)	1.70 [1.62, 1.78]	0.774	0.90 [0.83, 0.97]	0.762	1.95 [1.77, 2.16]	0.766
MPV	28679 (7094)	19120 (4729)	1.28 [1.22, 1.35]	0.765	0.93 [0.87, 0.99]	0.762	0.10 [0.04, 0.23]	0.762
NA	25977 (7133)	17318 (4756)	2.17 [2.04, 2.31]	0.753	1.03 [0.95, 1.12]	0.738	2.02 [1.88, 2.16]	0.750
PLT	30976 (7564)	20652 (5043)	1.98 [1.89, 2.08]	0.779	0.92 [0.83, 1.01]	0.765	1.88 [1.77, 1.98]	0.775
RBC	30946 (7556)	20631 (5038)	—	—	0.78 [0.71, 0.85]	0.764	—	—
RDW	30874 (7528)	20583 (5018)	3.09 [2.82, 3.39]	0.774	0.90 [0.83, 0.97]	0.762	2.23 [1.96, 2.52]	0.765
TBIL	13506 (3888)	9005 (2593)	1.61 [1.50, 1.72]	0.744	1.13 [1.04, 1.22]	0.737	1.09 [0.98, 1.21]	0.738
TC	25805 (6534)	17204 (4357)	0.93 [0.87, 0.99]	0.739	0.80 [0.75, 0.86]	0.739	0.96 [0.90, 1.03]	0.739
TGL	—	—	—	—	—	—	—	—
TP	13638 (4217)	9092 (2812)	2.23 [2.08, 2.38]	0.756	1.07 [0.98, 1.16]	0.736	2.73 [2.46, 3.04]	0.745
WBC	31053 (7583)	20702 (5055)	1.86 [1.75, 1.97]	0.773	0.98 [0.89, 1.07]	0.763	1.90 [1.75, 2.07]	0.768

Supplementary Table 22 | Proportional hazards analysis for in-hospital mortality in the eICU Collaborative Research Database. Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	—	—	—	—	—	—	—	—
ALB	11151 (1560)	7435 (1040)	1.04 [0.75, 1.43]	0.544	1.01 [0.65, 1.57]	0.544	1.14 [0.78, 1.65]	0.544
ALP	7966 (1118)	5311 (746)	0.97 [0.86, 1.09]	0.534	1.09 [0.93, 1.28]	0.534	1.12 [0.99, 1.26]	0.535
ALT	7288 (976)	4859 (650)	0.98 [0.86, 1.11]	0.538	1.05 [0.89, 1.24]	0.539	0.99 [0.87, 1.13]	0.537
AST	7446 (912)	4964 (608)	1.68 [1.47, 1.93]	0.571	1.82 [1.52, 2.19]	0.562	1.61 [1.40, 1.86]	0.575
BUN	33379 (3378)	22253 (2252)	1.74 [1.58, 1.92]	0.573	1.84 [1.59, 2.14]	0.559	1.77 [1.59, 1.97]	0.571
CA	33080 (3474)	22054 (2316)	1.81 [1.64, 1.98]	0.577	1.68 [1.48, 1.91]	0.564	1.75 [1.58, 1.94]	0.571
CL	34320 (3560)	22881 (2374)	1.52 [1.40, 1.65]	0.572	1.69 [1.51, 1.89]	0.568	1.58 [1.44, 1.72]	0.574
CO2	40679 (4792)	27120 (3194)	1.94 [1.76, 2.13]	0.586	1.82 [1.60, 2.08]	0.566	1.90 [1.71, 2.11]	0.576
CRE	31723 (3152)	21149 (2102)	1.89 [1.74, 2.05]	0.602	1.97 [1.78, 2.17]	0.599	1.98 [1.81, 2.16]	0.604
DBIL	684 (122)	456 (82)	1.23 [0.81, 1.87]	0.546	1.07 [0.69, 1.65]	0.512	1.14 [0.75, 1.75]	0.537
GLU	52615 (5182)	35077 (3455)	2.14 [1.73, 2.65]	0.565	1.65 [1.26, 2.15]	0.563	2.19 [1.75, 2.74]	0.565
HCT	34809 (3553)	23206 (2368)	1.11 [0.89, 1.38]	0.558	1.45 [0.98, 2.15]	0.556	1.31 [1.01, 1.71]	0.558
HDL	—	—	—	—	—	—	—	—
HGB	35018 (3622)	23346 (2415)	1.01 [0.82, 1.23]	0.561	1.11 [0.83, 1.50]	0.561	1.07 [0.85, 1.35]	0.561
K	38603 (3943)	25736 (2628)	1.34 [1.26, 1.43]	0.563	1.31 [1.22, 1.41]	0.563	1.36 [1.27, 1.45]	0.564
LDL	—	—	—	—	—	—	—	—
MCH	28067 (2889)	18712 (1926)	0.99 [0.92, 1.07]	0.558	0.95 [0.88, 1.03]	0.558	0.99 [0.92, 1.06]	0.558
MCHC	29987 (3063)	19992 (2042)	0.85 [0.78, 0.92]	0.563	0.87 [0.78, 0.97]	0.558	0.89 [0.82, 0.98]	0.561
MCV	29488 (3017)	19659 (2011)	1.26 [1.16, 1.37]	0.559	1.35 [1.25, 1.47]	0.564	1.35 [1.25, 1.45]	0.569
MPV	21532 (2191)	14355 (1461)	0.95 [0.84, 1.07]	0.568	1.17 [1.08, 1.28]	0.575	1.20 [1.02, 1.40]	0.571
NA	36226 (3839)	24152 (2559)	1.22 [1.14, 1.31]	0.551	1.37 [1.25, 1.50]	0.557	1.22 [1.11, 1.35]	0.552
PLT	30955 (3214)	20638 (2142)	2.50 [2.32, 2.68]	0.654	1.77 [1.58, 1.99]	0.570	1.67 [1.54, 1.82]	0.579
RBC	31071 (3221)	20714 (2148)	1.05 [0.88, 1.25]	0.562	1.07 [0.85, 1.34]	0.561	1.10 [0.90, 1.33]	0.561
RDW	28255 (2852)	18838 (1901)	1.85 [1.65, 2.07]	0.601	2.23 [1.89, 2.63]	0.589	1.91 [1.68, 2.16]	0.595
TBIL	7465 (930)	4978 (620)	1.39 [1.22, 1.59]	0.552	1.28 [1.12, 1.47]	0.550	1.38 [1.22, 1.58]	0.552
TC	—	—	—	—	—	—	—	—
TGL	186 (37)	124 (24)	1.09 [0.54, 2.18]	0.594	1.03 [0.43, 2.48]	0.590	1.22 [0.59, 2.49]	0.598
TP	8881 (1307)	5921 (871)	1.67 [1.44, 1.93]	0.576	1.68 [1.37, 2.06]	0.554	1.66 [1.42, 1.95]	0.562
WBC	30250 (2917)	20167 (1945)	2.39 [2.17, 2.63]	0.619	2.48 [2.16, 2.85]	0.598	2.41 [2.17, 2.67]	0.617

Supplementary Table 23 | Proportional hazards analysis for acute kidney injury in the eICU Collaborative Research Database. Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	—	—	—	—	—	—	—	—
ALB	11121 (2490)	7414 (1660)	1.28 [1.01, 1.63]	0.516	1.45 [1.03, 2.03]	0.515	1.36 [1.04, 1.79]	0.515
ALP	7949 (1702)	5300 (1134)	1.25 [1.14, 1.37]	0.544	1.12 [0.99, 1.26]	0.516	1.24 [1.12, 1.36]	0.534
ALT	7267 (1472)	4846 (982)	1.08 [0.97, 1.19]	0.504	0.94 [0.82, 1.07]	0.505	1.06 [0.95, 1.18]	0.504
AST	7429 (1464)	4954 (976)	1.19 [1.08, 1.32]	0.529	1.07 [0.95, 1.21]	0.504	1.14 [1.03, 1.27]	0.522
BUN	33279 (4964)	22187 (3310)	1.89 [1.76, 2.04]	0.566	1.63 [1.47, 1.82]	0.535	1.73 [1.60, 1.87]	0.555
CA	32980 (5407)	21987 (3604)	1.43 [1.34, 1.53]	0.541	1.40 [1.28, 1.52]	0.533	1.39 [1.29, 1.49]	0.539
CL	34213 (5471)	22809 (3647)	1.43 [1.34, 1.52]	0.540	1.24 [1.14, 1.34]	0.525	1.43 [1.34, 1.52]	0.538
CO2	40566 (6164)	27044 (4109)	1.40 [1.31, 1.50]	0.534	1.35 [1.23, 1.49]	0.524	1.33 [1.24, 1.44]	0.528
CRE	31638 (4454)	21092 (2969)	1.77 [1.66, 1.89]	0.576	1.55 [1.44, 1.66]	0.555	1.59 [1.49, 1.69]	0.560
DBIL	682 (211)	456 (141)	1.32 [0.97, 1.79]	0.520	1.32 [0.94, 1.86]	0.520	1.40 [1.01, 1.93]	0.523
GLU	52476 (6866)	34984 (4577)	1.06 [0.94, 1.19]	0.519	0.96 [0.82, 1.13]	0.518	1.04 [0.92, 1.17]	0.519
HCT	34716 (5015)	23144 (3343)	1.91 [1.55, 2.36]	0.524	2.59 [1.78, 3.79]	0.521	1.97 [1.56, 2.49]	0.524
HDL	—	—	—	—	—	—	—	—
HGB	34923 (5009)	23282 (3339)	1.97 [1.62, 2.40]	0.527	2.18 [1.61, 2.96]	0.524	2.01 [1.61, 2.50]	0.526
K	38496 (5647)	25664 (3764)	1.40 [1.33, 1.48]	0.557	1.33 [1.26, 1.41]	0.544	1.39 [1.32, 1.47]	0.554
LDL	—	—	—	—	—	—	—	—
MCH	27981 (4313)	18654 (2875)	1.12 [1.05, 1.19]	0.528	1.08 [1.01, 1.16]	0.527	1.14 [1.07, 1.21]	0.528
MCHC	29898 (4621)	19933 (3080)	1.17 [1.10, 1.25]	0.531	1.14 [1.05, 1.25]	0.529	1.13 [1.05, 1.21]	0.530
MCV	29400 (4528)	19601 (3018)	1.12 [1.04, 1.21]	0.523	1.02 [0.96, 1.08]	0.522	1.06 [1.00, 1.13]	0.523
MPV	21459 (3144)	14307 (2096)	1.30 [1.18, 1.43]	0.535	1.07 [1.00, 1.15]	0.523	1.10 [0.98, 1.24]	0.526
NA	36123 (5599)	24082 (3733)	1.13 [1.07, 1.19]	0.527	1.06 [0.99, 1.13]	0.528	1.01 [0.94, 1.08]	0.525
PLT	30866 (4735)	20578 (3156)	1.38 [1.30, 1.46]	0.543	1.03 [0.95, 1.12]	0.517	1.15 [1.08, 1.22]	0.524
RBC	30981 (4736)	20654 (3158)	1.75 [1.50, 2.04]	0.520	1.71 [1.40, 2.09]	0.521	1.78 [1.50, 2.11]	0.521
RDW	28174 (4340)	18783 (2894)	1.73 [1.60, 1.87]	0.571	1.65 [1.50, 1.83]	0.553	1.71 [1.58, 1.86]	0.568
TBIL	7447 (1482)	4966 (988)	1.15 [1.04, 1.28]	0.526	1.11 [1.00, 1.23]	0.527	1.12 [1.01, 1.24]	0.532
TC	20 (4)	14 (2)	0.52 [0.07, 3.77]	0.640	—	0.600	0.89 [0.08, 10.23]	0.640
TGL	186 (54)	124 (36)	1.18 [0.67, 2.09]	0.551	0.97 [0.47, 2.01]	0.515	1.02 [0.58, 1.82]	0.525
TP	8860 (1890)	5908 (1260)	1.28 [1.15, 1.43]	0.517	1.34 [1.15, 1.55]	0.510	1.31 [1.16, 1.48]	0.518
WBC	30162 (4532)	20108 (3021)	1.27 [1.19, 1.35]	0.537	1.17 [1.08, 1.26]	0.524	1.28 [1.20, 1.36]	0.531

Supplementary Table 24 | Proportional hazards analysis for sepsis in the eICU Collaborative Research Database.
Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	—	—	—	—	—	—	—	—
ALB	11106 (2754)	7405 (1836)	2.27 [1.72, 3.00]	0.519	2.42 [1.64, 3.56]	0.518	2.24 [1.66, 3.02]	0.518
ALP	7938 (2020)	5292 (1347)	1.17 [1.07, 1.28]	0.544	0.98 [0.88, 1.10]	0.503	1.12 [1.02, 1.22]	0.532
ALT	7256 (1877)	4838 (1252)	1.01 [0.92, 1.11]	0.510	1.01 [0.90, 1.13]	0.510	0.99 [0.90, 1.09]	0.512
AST	7415 (1904)	4944 (1270)	1.08 [0.98, 1.18]	0.502	0.96 [0.87, 1.07]	0.517	1.08 [0.98, 1.18]	0.495
BUN	33240 (5690)	22161 (3793)	1.22 [1.15, 1.30]	0.533	1.08 [0.99, 1.17]	0.527	1.16 [1.09, 1.24]	0.527
CA	32941 (5802)	21962 (3868)	1.53 [1.44, 1.63]	0.549	1.56 [1.43, 1.71]	0.538	1.51 [1.40, 1.62]	0.544
CL	34176 (5909)	22785 (3939)	1.34 [1.26, 1.41]	0.539	1.29 [1.20, 1.39]	0.528	1.32 [1.24, 1.40]	0.536
CO2	40521 (6798)	27015 (4532)	1.26 [1.18, 1.34]	0.540	1.17 [1.07, 1.27]	0.532	1.19 [1.12, 1.28]	0.534
CRE	31595 (5314)	21064 (3542)	1.28 [1.21, 1.35]	0.547	1.21 [1.14, 1.29]	0.536	1.21 [1.15, 1.28]	0.537
DBIL	682 (186)	456 (125)	0.98 [0.72, 1.33]	0.513	1.01 [0.72, 1.41]	0.515	1.09 [0.79, 1.50]	0.522
GLU	52429 (7603)	34953 (5068)	0.98 [0.88, 1.09]	0.527	0.93 [0.80, 1.08]	0.527	0.96 [0.86, 1.08]	0.527
HCT	34671 (5744)	23114 (3829)	1.50 [1.26, 1.79]	0.528	1.75 [1.31, 2.34]	0.527	1.47 [1.22, 1.78]	0.527
HDL	—	—	—	—	—	—	—	—
HGB	34876 (5787)	23252 (3858)	1.50 [1.28, 1.76]	0.531	1.53 [1.22, 1.93]	0.527	1.50 [1.26, 1.79]	0.529
K	38461 (6288)	25642 (4192)	1.48 [1.40, 1.55]	0.560	1.32 [1.25, 1.40]	0.541	1.41 [1.34, 1.48]	0.551
LDL	—	—	—	—	—	—	—	—
MCH	27943 (5010)	18630 (3340)	1.20 [1.14, 1.27]	0.533	1.11 [1.04, 1.18]	0.525	1.19 [1.13, 1.26]	0.533
MCHC	29860 (5363)	19908 (3575)	1.32 [1.24, 1.41]	0.547	1.30 [1.20, 1.41]	0.529	1.31 [1.23, 1.40]	0.541
MCV	29362 (5220)	19575 (3480)	1.24 [1.16, 1.33]	0.529	1.12 [1.05, 1.18]	0.510	1.22 [1.15, 1.29]	0.520
MPV	21432 (3423)	14289 (2282)	1.27 [1.16, 1.39]	0.520	1.05 [0.98, 1.12]	0.513	1.04 [0.93, 1.16]	0.514
NA	36085 (6030)	24058 (4020)	1.05 [1.00, 1.11]	0.519	1.00 [0.93, 1.06]	0.519	0.98 [0.92, 1.05]	0.519
PLT	30826 (5403)	20552 (3603)	1.16 [1.10, 1.23]	0.532	1.03 [0.95, 1.11]	0.526	1.07 [1.01, 1.13]	0.530
RBC	30939 (5515)	20626 (3677)	1.38 [1.22, 1.57]	0.522	1.47 [1.24, 1.74]	0.518	1.36 [1.19, 1.56]	0.519
RDW	28135 (5022)	18758 (3349)	1.83 [1.70, 1.96]	0.557	1.67 [1.52, 1.83]	0.535	1.77 [1.64, 1.91]	0.545
TBIL	7437 (1878)	4959 (1252)	1.02 [0.93, 1.12]	0.517	0.98 [0.89, 1.08]	0.511	1.01 [0.92, 1.11]	0.515
TC	20 (6)	14 (5)	0.71 [0.13, 3.89]	0.425	—	0.375	0.76 [0.08, 7.71]	0.425
TGL	186 (63)	124 (42)	0.95 [0.56, 1.59]	0.409	1.28 [0.63, 2.63]	0.439	0.89 [0.53, 1.50]	0.405
TP	8848 (2278)	5899 (1518)	1.07 [0.97, 1.18]	0.529	1.08 [0.95, 1.22]	0.527	1.11 [1.00, 1.24]	0.531
WBC	30127 (5170)	20086 (3447)	1.48 [1.39, 1.57]	0.545	1.28 [1.19, 1.38]	0.523	1.39 [1.31, 1.48]	0.536

Supplementary Table 25 | Proportional hazards analysis for prolonged ICU stay in the eICU Collaborative Research Database. Prolonged ICU stay is defined as ICU stay greater than 7 days. Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	—	—	—	—	—	—	—	—
ALB	11140 (3587)	7428 (2392)	0.75 [0.61, 0.93]	0.535	0.70 [0.51, 0.96]	0.536	0.77 [0.60, 0.97]	0.535
ALP	7956 (2821)	5305 (1881)	0.68 [0.63, 0.73]	0.551	0.84 [0.76, 0.92]	0.533	0.73 [0.68, 0.79]	0.548
ALT	7276 (2596)	4851 (1731)	0.82 [0.76, 0.88]	0.533	0.87 [0.78, 0.96]	0.520	0.85 [0.78, 0.92]	0.527
AST	7437 (2658)	4958 (1772)	0.86 [0.80, 0.93]	0.545	0.98 [0.89, 1.07]	0.534	0.87 [0.81, 0.94]	0.545
BUN	33336 (7040)	22225 (4694)	0.66 [0.62, 0.71]	0.550	0.76 [0.70, 0.83]	0.535	0.67 [0.63, 0.72]	0.548
CA	33039 (6996)	22026 (4664)	0.79 [0.75, 0.84]	0.541	0.77 [0.71, 0.83]	0.537	0.75 [0.71, 0.80]	0.544
CL	34278 (7202)	22853 (4801)	0.74 [0.70, 0.78]	0.552	0.84 [0.79, 0.90]	0.538	0.75 [0.71, 0.80]	0.548
CO2	40636 (7193)	27092 (4795)	0.63 [0.58, 0.68]	0.547	0.72 [0.66, 0.80]	0.540	0.61 [0.56, 0.66]	0.545
CRE	31684 (6787)	21123 (4525)	0.68 [0.65, 0.72]	0.566	0.66 [0.62, 0.70]	0.562	0.69 [0.66, 0.73]	0.566
DBIL	684 (328)	456 (218)	1.02 [0.81, 1.29]	0.512	0.97 [0.75, 1.25]	0.482	0.95 [0.75, 1.22]	0.461
GLU	52561 (7373)	35041 (4915)	0.44 [0.36, 0.53]	0.537	0.53 [0.41, 0.70]	0.535	0.47 [0.38, 0.57]	0.536
HCT	34765 (7128)	23178 (4753)	0.53 [0.46, 0.61]	0.541	0.64 [0.51, 0.81]	0.537	0.53 [0.45, 0.62]	0.539
HDL	—	—	—	—	—	—	—	—
HGB	34972 (7128)	23316 (4752)	0.50 [0.43, 0.58]	0.543	0.51 [0.42, 0.62]	0.537	0.52 [0.44, 0.61]	0.542
K	38559 (7280)	25707 (4854)	0.77 [0.73, 0.80]	0.553	0.81 [0.77, 0.85]	0.543	0.76 [0.73, 0.80]	0.554
LDL	—	—	—	—	—	—	—	—
MCH	28026 (6451)	18684 (4301)	0.92 [0.87, 0.97]	0.537	0.86 [0.81, 0.91]	0.544	0.83 [0.79, 0.87]	0.541
MCHC	29944 (6867)	19964 (4578)	0.67 [0.63, 0.71]	0.559	0.67 [0.62, 0.72]	0.552	0.65 [0.61, 0.69]	0.559
MCV	29445 (6783)	19631 (4523)	0.88 [0.83, 0.94]	0.539	0.97 [0.92, 1.02]	0.540	0.83 [0.80, 0.88]	0.546
MPV	21492 (4995)	14328 (3330)	0.85 [0.78, 0.92]	0.535	0.98 [0.92, 1.04]	0.529	0.93 [0.84, 1.03]	0.532
NA	36186 (7220)	24125 (4814)	0.74 [0.71, 0.78]	0.560	0.80 [0.75, 0.85]	0.543	0.76 [0.71, 0.82]	0.549
PLT	30913 (7030)	20609 (4686)	0.90 [0.86, 0.94]	0.534	1.22 [1.14, 1.30]	0.542	0.96 [0.91, 1.01]	0.537
RBC	31027 (7105)	20686 (4737)	0.55 [0.49, 0.62]	0.549	0.53 [0.45, 0.63]	0.541	0.54 [0.47, 0.62]	0.547
RDW	28216 (6496)	18812 (4331)	0.67 [0.63, 0.71]	0.557	0.70 [0.65, 0.76]	0.546	0.67 [0.63, 0.72]	0.554
TBIL	7456 (2547)	4972 (1699)	0.86 [0.79, 0.93]	0.537	0.90 [0.83, 0.98]	0.542	0.87 [0.81, 0.95]	0.534
TC	20 (8)	14 (5)	0.67 [0.11, 4.11]	0.600	—	—	0.20 [0.00, 12.22]	0.500
TGL	186 (134)	124 (90)	0.82 [0.57, 1.18]	0.527	0.74 [0.46, 1.19]	0.533	0.91 [0.63, 1.31]	0.528
TP	8871 (3098)	5914 (2065)	1.03 [0.95, 1.12]	0.524	0.97 [0.87, 1.09]	0.527	0.98 [0.90, 1.07]	0.527
WBC	30209 (7003)	20140 (4668)	0.75 [0.71, 0.79]	0.557	0.79 [0.74, 0.85]	0.545	0.74 [0.69, 0.78]	0.552

Supplementary Table 26 | Proportional hazards analysis for in-hospital mortality in the INSPIRE cohort. Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	233 (13)	156 (8)	0.13 [0.01, 2.06]	0.446	0.13 [0.01, 2.06]	0.446	—	0.482
ALB	18427 (470)	12285 (314)	6.58 [4.90, 8.83]	0.805	4.28 [3.06, 6.00]	0.719	5.87 [4.30, 8.02]	0.785
ALP	16668 (437)	11113 (292)	2.10 [1.72, 2.56]	0.745	2.40 [1.89, 3.04]	0.727	2.43 [1.98, 2.98]	0.754
ALT	15676 (441)	10452 (294)	1.66 [1.37, 2.02]	0.632	1.48 [1.20, 1.82]	0.636	1.51 [1.24, 1.83]	0.637
AST	15814 (415)	10544 (277)	2.17 [1.76, 2.68]	0.742	1.99 [1.62, 2.45]	0.713	2.21 [1.81, 2.70]	0.732
BUN	17938 (447)	11960 (298)	2.40 [1.90, 3.02]	0.724	2.21 [1.71, 2.86]	0.670	2.41 [1.91, 3.06]	0.707
CA	17863 (456)	11909 (304)	3.08 [2.48, 3.83]	0.764	2.89 [2.25, 3.71]	0.713	3.24 [2.58, 4.07]	0.776
CL	18561 (442)	12374 (294)	2.79 [2.21, 3.53]	0.791	2.61 [2.02, 3.39]	0.718	2.81 [2.20, 3.60]	0.773
CO2	3742 (331)	2495 (221)	3.71 [2.58, 5.34]	0.643	3.25 [2.10, 5.03]	0.616	3.20 [2.15, 4.76]	0.627
CRE	24427 (421)	16285 (281)	2.63 [2.12, 3.26]	0.731	2.47 [1.95, 3.13]	0.745	2.70 [2.16, 3.38]	0.750
DBIL	—	—	—	—	—	—	—	—
GLU	15690 (419)	10461 (279)	1.87 [1.41, 2.49]	0.681	1.91 [1.37, 2.65]	0.658	2.07 [1.39, 3.09]	0.655
HCT	23831 (460)	15888 (307)	4.33 [3.01, 6.23]	0.730	3.96 [2.49, 6.29]	0.676	4.07 [2.76, 6.00]	0.724
HDL	—	—	—	—	—	—	—	—
HGB	21702 (466)	14469 (310)	4.38 [3.06, 6.29]	0.719	3.82 [2.48, 5.89]	0.683	4.31 [2.94, 6.31]	0.717
K	21804 (440)	14537 (294)	2.98 [2.43, 3.67]	0.821	2.61 [2.10, 3.24]	0.747	2.99 [2.42, 3.69]	0.806
LDL	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—
NA	21663 (455)	14442 (303)	2.37 [1.95, 2.88]	0.741	2.90 [2.31, 3.65]	0.738	2.64 [2.15, 3.24]	0.765
PLT	20394 (466)	13597 (310)	4.55 [3.66, 5.65]	0.846	3.75 [2.89, 4.86]	0.752	3.96 [3.15, 4.96]	0.816
RBC	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—
TBIL	17133 (424)	11423 (282)	3.36 [2.76, 4.10]	0.758	2.78 [2.25, 3.43]	0.761	2.87 [2.36, 3.51]	0.765
TC	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—
TP	17152 (470)	11436 (313)	4.44 [3.57, 5.52]	0.822	3.20 [2.54, 4.04]	0.733	3.93 [3.15, 4.90]	0.799
WBC	20554 (458)	13703 (305)	3.80 [3.03, 4.78]	0.755	3.42 [2.68, 4.37]	0.751	3.83 [3.03, 4.84]	0.799

Supplementary Table 27 | Proportional hazards analysis for perioperative infection in the INSPIRE cohort. Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	226 (32)	152 (21)	0.82 [0.35, 1.97]	0.559	1.30 [0.43, 4.00]	0.572	1.80 [0.58, 5.56]	0.572
ALB	18112 (691)	12076 (461)	1.22 [1.05, 1.43]	0.624	1.14 [0.97, 1.34]	0.626	1.15 [0.98, 1.34]	0.631
ALP	16373 (656)	10916 (438)	1.16 [0.98, 1.38]	0.593	1.12 [0.96, 1.30]	0.598	1.16 [0.99, 1.36]	0.590
ALT	15382 (665)	10255 (443)	1.03 [0.88, 1.21]	0.610	1.02 [0.87, 1.19]	0.610	1.08 [0.93, 1.26]	0.614
AST	15519 (665)	10347 (444)	1.45 [1.15, 1.84]	0.642	1.20 [1.02, 1.40]	0.625	1.26 [1.03, 1.54]	0.644
BUN	17637 (682)	11758 (455)	1.09 [0.94, 1.27]	0.557	1.06 [0.90, 1.24]	0.548	1.12 [0.96, 1.31]	0.556
CA	17550 (687)	11700 (458)	1.17 [1.00, 1.36]	0.582	1.21 [1.04, 1.42]	0.593	1.15 [0.99, 1.34]	0.585
CL	18256 (689)	12172 (460)	1.09 [0.94, 1.27]	0.603	1.03 [0.88, 1.20]	0.595	1.14 [0.98, 1.32]	0.605
CO2	3583 (413)	2390 (275)	1.00 [0.81, 1.22]	0.575	0.98 [0.78, 1.23]	0.577	1.02 [0.82, 1.27]	0.577
CRE	24107 (694)	16072 (462)	1.08 [0.92, 1.26]	0.573	0.97 [0.84, 1.13]	0.571	1.07 [0.91, 1.25]	0.571
DBIL	—	—	—	—	—	—	—	—
GLU	15420 (629)	10281 (419)	0.97 [0.81, 1.16]	0.547	1.00 [0.81, 1.22]	0.547	1.00 [0.79, 1.27]	0.547
HCT	23504 (722)	15670 (482)	1.09 [0.92, 1.28]	0.616	1.10 [0.90, 1.35]	0.612	1.05 [0.88, 1.26]	0.615
HDL	—	—	—	—	—	—	—	—
HGB	21375 (716)	14251 (478)	1.06 [0.90, 1.25]	0.602	1.00 [0.82, 1.20]	0.592	1.06 [0.89, 1.26]	0.601
K	21492 (704)	14329 (469)	1.04 [0.88, 1.24]	0.610	1.05 [0.90, 1.22]	0.609	1.02 [0.87, 1.19]	0.611
LDL	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—
NA	21349 (703)	14234 (469)	1.12 [0.95, 1.33]	0.604	1.03 [0.88, 1.19]	0.600	0.97 [0.83, 1.13]	0.593
PLT	20078 (706)	13386 (471)	1.06 [0.90, 1.26]	0.611	1.15 [0.99, 1.33]	0.603	1.22 [1.05, 1.41]	0.638
RBC	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—
TBIL	16829 (676)	11220 (450)	1.07 [0.87, 1.31]	0.576	1.10 [0.94, 1.29]	0.584	1.02 [0.85, 1.22]	0.578
TC	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—
TP	16841 (680)	11228 (454)	1.27 [1.08, 1.49]	0.601	1.41 [1.21, 1.64]	0.582	1.25 [1.08, 1.46]	0.607
WBC	20235 (707)	13491 (471)	1.18 [1.01, 1.37]	0.558	1.21 [1.05, 1.41]	0.564	1.26 [1.09, 1.47]	0.559

Supplementary Table 28 | Proportional hazards analysis for prolonged hospital stay in the INSPIRE cohort.
Prolonged hospital stay is defined as hospital stay greater than 7 days. Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	233 (164)	156 (109)	0.95 [0.62, 1.45]	0.511	0.82 [0.46, 1.45]	0.500	0.88 [0.52, 1.48]	0.523
ALB	18427 (14140)	12285 (9427)	0.70 [0.67, 0.73]	0.566	0.73 [0.71, 0.76]	0.568	0.66 [0.64, 0.69]	0.572
ALP	16668 (13163)	11113 (8776)	0.67 [0.64, 0.71]	0.558	1.10 [1.06, 1.13]	0.532	0.82 [0.79, 0.85]	0.538
ALT	15676 (12343)	10452 (8230)	0.85 [0.82, 0.88]	0.545	1.02 [0.99, 1.06]	0.533	0.92 [0.89, 0.95]	0.536
AST	15814 (12427)	10544 (8286)	0.71 [0.65, 0.77]	0.539	1.12 [1.08, 1.16]	0.539	0.95 [0.90, 1.00]	0.530
BUN	17938 (14031)	11960 (9355)	0.75 [0.72, 0.77]	0.553	0.96 [0.92, 0.99]	0.538	0.76 [0.74, 0.79]	0.551
CA	17863 (13403)	11909 (8936)	0.89 [0.86, 0.92]	0.539	0.95 [0.92, 0.98]	0.538	0.88 [0.85, 0.91]	0.540
CL	18561 (14572)	12374 (9714)	0.77 [0.74, 0.79]	0.552	0.97 [0.93, 1.00]	0.525	0.83 [0.81, 0.86]	0.540
CO2	3742 (3239)	2495 (2160)	0.83 [0.77, 0.89]	0.516	0.98 [0.91, 1.06]	0.507	0.85 [0.79, 0.92]	0.515
CRE	24427 (17097)	16285 (11399)	0.65 [0.63, 0.68]	0.580	0.71 [0.68, 0.73]	0.574	0.68 [0.66, 0.71]	0.578
DBIL	—	—	—	—	—	—	—	—
GLU	15690 (11167)	10461 (7446)	1.03 [0.99, 1.07]	0.536	1.07 [1.02, 1.11]	0.538	1.02 [0.97, 1.07]	0.536
HCT	23831 (16876)	15888 (11251)	0.81 [0.78, 0.83]	0.560	0.85 [0.82, 0.88]	0.555	0.80 [0.78, 0.83]	0.560
HDL	—	—	—	—	—	—	—	—
HGB	21702 (15789)	14469 (10527)	0.74 [0.71, 0.76]	0.566	0.75 [0.72, 0.78]	0.557	0.71 [0.69, 0.74]	0.568
K	21804 (15982)	14537 (10655)	0.73 [0.69, 0.76]	0.538	1.02 [0.99, 1.05]	0.533	0.85 [0.82, 0.89]	0.532
LDL	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—
NA	21663 (15923)	14442 (10615)	0.71 [0.68, 0.74]	0.546	0.95 [0.92, 0.98]	0.528	0.76 [0.73, 0.79]	0.545
PLT	20394 (15193)	13597 (10129)	0.69 [0.66, 0.73]	0.553	1.22 [1.18, 1.26]	0.553	1.04 [1.00, 1.07]	0.542
RBC	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—
TBIL	17133 (13177)	11423 (8786)	0.87 [0.82, 0.92]	0.534	0.95 [0.92, 0.99]	0.532	0.95 [0.91, 0.99]	0.532
TC	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—
TP	17152 (13544)	11436 (9031)	0.79 [0.76, 0.83]	0.542	0.96 [0.93, 0.99]	0.535	0.78 [0.75, 0.81]	0.545
WBC	20554 (15316)	13703 (10211)	0.80 [0.78, 0.83]	0.545	0.97 [0.94, 1.00]	0.537	0.84 [0.81, 0.87]	0.543

Supplementary Table 29 | Proportional hazards analysis for unplanned ICU admission in the INSPIRE cohort.
Per-analyte hazard ratios (with 95% confidence intervals) and concordance indices for Pop_{RI}, Per_{RI}, and NORMA_{RI} abnormality flags, adjusted for age and sex.

Analyte	N train (events)	N test (events)	Pop _{RI}		Per _{RI}		NORMA _{RI}	
			HR [95% CI]	C	HR [95% CI]	C	HR [95% CI]	C
A1C	233 (105)	156 (70)	1.09 [0.68, 1.74]	0.495	1.00 [0.59, 1.69]	0.499	1.17 [0.66, 2.07]	0.528
ALB	18427 (5221)	12285 (3481)	0.85 [0.80, 0.91]	0.541	0.90 [0.86, 0.96]	0.542	0.94 [0.89, 1.00]	0.539
ALP	16668 (5096)	11113 (3397)	1.07 [0.99, 1.15]	0.535	0.96 [0.91, 1.01]	0.539	1.01 [0.95, 1.08]	0.536
ALT	15676 (5066)	10452 (3378)	0.95 [0.90, 1.02]	0.535	0.94 [0.88, 0.99]	0.537	1.01 [0.96, 1.07]	0.534
AST	15814 (5026)	10544 (3351)	1.14 [1.01, 1.28]	0.530	0.84 [0.79, 0.89]	0.540	0.90 [0.82, 0.98]	0.532
BUN	17938 (5376)	11960 (3585)	1.00 [0.94, 1.06]	0.523	0.94 [0.89, 0.99]	0.524	1.01 [0.96, 1.07]	0.523
CA	17863 (5265)	11909 (3510)	0.85 [0.80, 0.90]	0.534	0.87 [0.82, 0.91]	0.536	0.84 [0.79, 0.89]	0.536
CL	18561 (5536)	12374 (3691)	1.29 [1.22, 1.37]	0.530	1.10 [1.04, 1.15]	0.521	1.42 [1.34, 1.49]	0.544
CO2	3742 (2059)	2495 (1373)	0.93 [0.86, 1.02]	0.521	0.93 [0.84, 1.02]	0.520	0.90 [0.82, 0.99]	0.518
CRE	24427 (5246)	16285 (3498)	1.26 [1.18, 1.35]	0.563	1.19 [1.12, 1.26]	0.562	1.22 [1.15, 1.30]	0.562
DBIL	—	—	—	—	—	—	—	—
GLU	15690 (4539)	10461 (3026)	1.03 [0.97, 1.10]	0.529	0.93 [0.87, 1.00]	0.531	1.26 [1.15, 1.37]	0.529
HCT	23831 (6037)	15888 (4025)	0.94 [0.89, 0.99]	0.529	0.98 [0.92, 1.04]	0.529	1.05 [0.99, 1.10]	0.528
HDL	—	—	—	—	—	—	—	—
HGB	21702 (5792)	14469 (3862)	1.01 [0.96, 1.07]	0.532	1.10 [1.04, 1.17]	0.532	1.05 [0.99, 1.11]	0.533
K	21804 (5871)	14537 (3915)	1.11 [1.02, 1.20]	0.520	0.93 [0.88, 0.98]	0.522	1.13 [1.05, 1.20]	0.518
LDL	—	—	—	—	—	—	—	—
MCH	—	—	—	—	—	—	—	—
MCHC	—	—	—	—	—	—	—	—
MCV	—	—	—	—	—	—	—	—
MPV	—	—	—	—	—	—	—	—
NA	21663 (5875)	14442 (3916)	1.31 [1.22, 1.40]	0.532	0.98 [0.93, 1.03]	0.511	1.37 [1.29, 1.45]	0.539
PLT	20394 (5724)	13597 (3817)	1.35 [1.25, 1.45]	0.537	0.83 [0.79, 0.87]	0.540	0.99 [0.93, 1.04]	0.531
RBC	—	—	—	—	—	—	—	—
RDW	—	—	—	—	—	—	—	—
TBIL	17133 (5043)	11423 (3363)	1.11 [1.02, 1.21]	0.534	1.07 [1.01, 1.14]	0.535	1.09 [1.02, 1.17]	0.535
TC	—	—	—	—	—	—	—	—
TGL	—	—	—	—	—	—	—	—
TP	17152 (5154)	11436 (3437)	0.87 [0.81, 0.94]	0.539	0.87 [0.82, 0.92]	0.542	0.89 [0.83, 0.95]	0.538
WBC	20554 (5663)	13703 (3775)	1.16 [1.10, 1.23]	0.540	1.00 [0.95, 1.05]	0.537	1.15 [1.09, 1.22]	0.541