

Default Contagion, Matrix Approximation, and Control in Sparse Financial Networks

Aoxin Zhang*

Yingzhe Wang†

May 26, 2026

Abstract

We study systemic default contagion in sparse financial networks and develop a framework for deciding when aggregate exposure matrices are reliable and when node-level network information changes tail risk and control design. The first contribution is a multi-population McKean–Vlasov foundation for distance-to-default dynamics with common noise, bounded state-dependent killing, loss feedback, sparse weighted exposures, and regulatory intervention, including quantitative convergence, propagation of chaos, stability in contagion matrices, controlled well-posedness, a two-population HJB characterization, and a steep-killing bridge to absorbing-boundary contagion. The second contribution is a set of computable matrix-approximation diagnostics: finite-grid bounds driven by row-exposure dispersion and square-edge spread, constructive tail-loss gaps for networks sharing the same aggregate matrix, and a spectral-radius criterion for local cascade onset. The third contribution is an information-value theory for control, showing that node-level graph pressure has strictly positive value when within-type pressure variation interacts with nonsaturated marginal killing reduction. Matched sparse-graph and matrix experiments, common-noise tests, HJB feedback diagnostics, fixed-budget control comparisons, and EBA/Pillar 3-calibrated synthetic networks validate the framework. The main conclusion is that finite-type matrices are effective in regular-mixing regimes, whereas concentrated sparse exposures generate tail-risk and intervention effects that require local-pressure diagnostics and network-aware control.

Keywords. systemic risk, financial networks, McKean–Vlasov equations, default contagion, sparse graphs, common noise, optimal control, matrix approximation

MSC 2020. 91G40, 91G80, 60H30, 60K35, 93E20, 49L20, 05C82

1 Introduction

Systemic default risk is generated by an interaction between balance-sheet fragility and network structure. The clearing-network model of Eisenberg and Noe [1] provides the standard limited-liability benchmark, while Allen and Gale [2], Gai and Kapadia [3], Acemoglu, Ozdaglar, and Tahbaz-Salehi [4], Elliott, Golub, and Jackson [5], and Glasserman and Young [6] show that diversification, connectivity, and shock size have non-monotone implications for cascade risk. A parallel large-population literature models default contagion through interacting particles and McKean–Vlasov equations. Hambly, Ledger, and Søjmark [7], Hambly and Søjmark [8], and Hambly

*School of Mathematical Sciences, Beijing Normal University, Beijing, China. Email: 202211130012@mail.bnu.edu.cn. ORCID: <https://orcid.org/0009-0006-3647-9545>.

†School of Mathematical Sciences, Beijing Normal University, Beijing, China. Corresponding author. Email: wangyz@bnu.edu.cn. ORCID: <https://orcid.org/0000-0003-2113-3307>.

and Jettkant [9] show how distance-to-default dynamics, loss feedback, state-dependent killing, and control can be represented through low-dimensional distributional dynamics.

The practical tension is that regulatory stress tests often need aggregate and interpretable objects, while real interbank networks remain sparse, heterogeneous, and locally concentrated. A finite-type contagion matrix is attractive because it summarizes cross-type exposures, supports spectral and row-exposure diagnostics, and leads to tractable control rules. Sparse financial networks, however, can transmit losses through hub nodes, core-periphery concentration, and local high-pressure neighborhoods that disappear under type averaging. This paper studies the boundary between these two representations by constructing a sparse graph contagion system and an aggregate matrix approximation from the same graph. The two systems share node types, initial states, kill intensities, loss feedback, and aggregate exposure sources, so their difference isolates the information loss caused by aggregating local network structure.

This boundary also matters for systemic-risk measurement and policy design. Market-based measures such as systemic expected shortfall and CoVaR quantify institution-level contributions to aggregate distress [10, 11]; clearing and network-contagion models explain how balance-sheet links propagate distress once a shock has entered the system. Our focus is the interface between these two viewpoints: we ask when a regulator can work with a low-dimensional exposure matrix, and when tail-risk measurement requires additional local graph information. This framing leads to a concrete information question. If two networks have the same type-level matrix but different within-block concentration, then any difference in tail loss, cascade probability, or intervention cost is attributable to information discarded by the matrix projection.

The paper therefore combines a regularized McKean–Vlasov limit with finite-network diagnostics. We show that the matrix–network boundary is theoretically characterizable through computable graph diagnostics and practically consequential for regulatory control design. The regularization through bounded state-dependent killing keeps the default mechanism close to absorbing-boundary contagion while leaving the controlled system compatible with dynamic programming. The sparse-network layer is finite and constructive: it compares matched graph and matrix systems under identical shocks and budgets, so the empirical design tests the same approximation boundary described by the theory.

Main contributions The first contribution is a multi-population McKean–Vlasov foundation for regularized default contagion with common noise, loss feedback, and control. We prove the large-system limit, a conditional $O(N_*^{-1/2})$ first-order convergence rate under explicit quadratic empirical-process and metric-stability inputs, same-type propagation of chaos, controlled well-posedness, and Lipschitz stability with respect to the contagion matrix. This foundation also supports two control-facing extensions: a two-population HJB characterization in a standard lifted comparison framework, and a steep-killing bridge to absorbing-boundary contagion under nonsticky-boundary, uniform-map-continuity, and local fixed-point-stability conditions. The result is a rigorous bounded-killing theory with a transparent interface to the absorbing-boundary and viscosity tools used in systemic-risk control.

The second contribution is a structural characterization of the matrix-approximation boundary. We derive a finite-grid absolute graph-to-matrix expected-loss bound controlled by two computable structural quantities: within-block row-exposure dispersion and square-edge exposure spread. A share-controlled corollary then translates the square-edge term into a maximum-edge-share diagnostic. We also prove a heterogeneous-network tail-gap construction, embed it in any larger finite type system, and connect local cascade onset to a spectral-radius threshold under an explicit low-loss second-order approximation. The numerical design then uses stochastic block, core-periphery,

heavy-tailed, Barabási–Albert, and Watts–Strogatz networks to show how the same aggregate matrix can underestimate sparse-graph tail loss when local exposure concentration is high.

The third contribution is a control and information-value analysis. We give an explicit variational sufficient condition under which node-level graph-pressure information has strictly positive control value near the matrix-approximation boundary, and we verify primitive pressure-monotonicity conditions for exponential killing in a nonsaturated local window. The rule-based experiments show where local pressure information gives low-cost cascade reduction, while the finite-dimensional $K = 2$ HJB loop turns the measure-valued Hamiltonian characterization into a computable feedback benchmark. The EBA Transparency Exercise and Pillar 3/EDAP calibration layer further tests whether the mechanism remains visible under external regulatory data constraints.

The rest of the paper is organized as follows. Section 2 positions the paper in the financial-network, McKean–Vlasov, and graph-limit literatures. Section 3 defines the finite-type and sparse-graph contagion systems. Sections 4–7 state the main theoretical results. Sections 8–14 present the numerical and calibration evidence, and Section 15 discusses implications and limitations.

2 Related Literature

2.1 Financial networks and default cascades

We now expand on the financial-network, interacting-particle, and graph-limit literatures introduced above, with emphasis on the technical distinction between aggregate exposure matrices and local sparse-network information.

Network models of systemic risk begin from the observation that a bank’s default losses depend on the payment, funding, liquidity, and asset-sale behavior of other institutions. The clearing-vector framework of Eisenberg and Noe [1] and the default-cost extension of Rogers and Veraart [12] provide benchmark models for mutual-liability networks. Liquidity and fire-sale channels are developed by Cifuentes, Ferrucci, and Shin [13], Gai, Haldane, and Kapadia [14], Caballero and Simsek [15], and May and Arinaminpathy [16]. Empirical and stylized interbank-network studies by Boss et al. [17], Nier et al. [18], Cont, Moussa, and Santos [19], Battiston et al. [20], Glasserman and Young [21], Caccioli et al. [22], Capponi, Chen, and Yao [23], and Gandy and Veraart [24] show that degree distributions, exposure concentration, centrality, and balance-sheet fragility jointly determine cascade paths. Empirical high-dimensional bank-network estimation, such as Demirer, Diebold, Liu, and Yilmaz [25], further reinforces the role of cross-bank connectedness in crisis periods.

The robust-yet-fragile mechanism is especially important for this paper. Allen and Gale [2], Gai and Kapadia [3], Acemoglu, Ozdaglar, and Tahbaz-Salehi [4], Elliott, Golub, and Jackson [5], Amini, Cont, and Minca [26], Gleeson [27], Hurd, Gleeson, and Melnik [28], and Glasserman and Young [6] all point to a common lesson: aggregate connectivity can stabilize small shocks while amplifying losses once local cascades start. Our sparse-graph experiments follow this line, but the object of comparison is deliberately narrower. We compare a sparse graph system with the finite-type matrix obtained from the same graph, so the measured gap is the price of type aggregation rather than a difference across unrelated network specifications.

2.2 Mean-field contagion, McKean–Vlasov models, and control

Mean-field contagion models describe the dynamic state of a large financial system through empirical measures and loss processes; the broader systemic-risk modeling background is surveyed in Fouque and Langsam [29]. Diffusion-based mechanisms for systemic risk and interbank stability are analyzed in Fouque and Ichiba [30] and Garnier, Papanicolaou, and Yang [31]. Hambly, Ledger, and Søjmark [7]

study McKean–Vlasov equations with positive feedback and blow-up, Hambly and Søjmark [8] derive an SPDE model for systemic risk with endogenous contagion, and Ledger and Søjmark [32] analyze uniqueness in the weak-feedback regime. Delarue et al. [33] provide a related particle-approximation theory for singular mean-field self-excitation, while Sznitman [34] and Budhiraja, Dupuis, and Fischer [35] give standard propagation-of-chaos and weak-convergence reference points for interacting-particle limits. Feinstein and Søjmark [36, 37] connect dynamic interbank default contagion with heterogeneous McKean–Vlasov limits, while Hambly and Jettkant [9] study state-dependent killing and central control. We use state-dependent killing as a regularized default mechanism because it keeps default probabilities state sensitive and gives a stable interface for finite-dimensional control diagnostics.

The control side draws on McKean–Vlasov control, infinite-dimensional HJB theory, numerical methods for McKean–Vlasov equations, and network mean-field methods. Huang, Malhamé, and Caines [38], Carmona, Fouque, and Sun [39], and Carmona and Delarue [40] provide the probabilistic background for mean-field control, mean-field games, and systemic-risk control; Fabbri, Gozzi, and Swiech [41] give dynamic programming and HJB theory in infinite dimensions; Crandall, Ishii, and Lions [42] provide the classical viscosity comparison framework; Cosso et al. [43] provide a comparison principle for Bellman equations on Wasserstein space, while Gangbo, Mészáros, Mou, and Zhang [44] give a complementary master-equation reference point under displacement monotonicity; and Chassagneux, Crisan, and Delarue [45] give numerical methods for McKean–Vlasov FBSDEs. Deep-BSDE methods such as Han, Jentzen, and E [46] provide an alternative high-dimensional PDE benchmark, but our finite-dimensional HJB loop is used as a transparent diagnostic rather than a black-box solver. Graph-limit and sparse-network perspectives enter through Lovász [47], Benjamini and Schramm [48], Borgs et al. [49], Delattre, Giacomini, and Luçon [50], Oliveira and Reis [51], Oliveira, Reis, and Stolerman [52], Parise and Ozdaglar [53], Caines and Huang [54], Carmona, Cooney, Graves, and Laurière [55], Aurell, Carmona, Dayanikli, and Laurière [56], Bayraktar, Chakraborty, and Wu [57], Lacker and Soret [58], and Bendel [59]. These works motivate our central question: when is a finite-type matrix a valid low-dimensional projection of sparse financial contagion, and when does node-level pressure information become necessary for tail-risk control?

3 Model

3.1 Finite-type state-dependent killing model

Consider a system of N financial institutions. Institution i has type $g(i) \in \{1, \dots, K\}$, and the state variable $X_i(t)$ represents its safety margin from the default region. The baseline kill intensity of type k is

$$\lambda_{0,k}(x) = \lambda_{\text{base},k} \exp\{\gamma_k(x_{b,k} - x)\}, \quad (1)$$

where $x_{b,k}$ is the risk boundary and γ_k is the state-sensitivity coefficient. Regulatory control a reduces the kill intensity in exponential form:

$$\lambda_k(x, a) = \lambda_{0,k}(x) \exp\{-\eta_k a\}, \quad (2)$$

and the kill probability over a discrete step Δt is

$$p_k(x, a; \Delta t) = 1 - \exp\{-\lambda_k(x, a)\Delta t\}. \quad (3)$$

This regularized killing mechanism preserves the state dependence of default risk and avoids the numerical instability of hard absorbing boundaries in finite-sample simulations.

The cumulative loss of type k and the aggregate loss are defined by

$$L_t^{k,N} = \frac{1}{N_k} \sum_{i:g(i)=k} \mathbf{1}_{\{\tau_i \leq t\}}, \quad L_t^N = \sum_{k=1}^K \pi_k L_t^{k,N}, \quad (4)$$

where $\pi_k = N_k/N$. The state dynamics of the finite-type matrix model are

$$dX_i^k(t) = \left[\beta_k + \kappa_k (m_t^{k,N} - X_i^k(t)) \right] dt + \sigma_k dW_i^k(t) + \sigma_0 dW^0(t) - \sum_{\ell=1}^K \Gamma_{k\ell} dL_t^{\ell,N}. \quad (5)$$

The matrix element $\Gamma_{k\ell} \geq 0$ denotes the average shock from newly realized losses in type ℓ to type k . The row sum $\sum_{\ell} \Gamma_{k\ell}$ measures the external exposure of the target type, the column sum $\sum_k \Gamma_{k\ell}$ measures the spillover impact of the source type, and the spectral radius $\rho(\Gamma)$ provides the aggregate feedback scale used later in the local threshold analysis.

3.2 Sparse graph local contagion and aggregate matrix

Let $A = (A_{ij})$ be a weighted directed adjacency matrix, where A_{ij} denotes the shock intensity from source node j to target node i when node j newly defaults. If $\Delta N_{j,t}$ denotes the new-default indicator of node j at time t , then the local graph pressure received by node i is

$$P_{i,t}^{\text{graph}} = \sum_{j=1}^N A_{ij} \Delta N_{j,t}. \quad (6)$$

At each discrete time step, nodes first undergo drift, diffusion, and state-dependent killing sampling. The surviving nodes then receive local graph-pressure shocks:

$$X_{i,t+\Delta t} = X_{i,t+\Delta t}^{\text{pre}} - P_{i,t}^{\text{graph}}. \quad (7)$$

To avoid notational ambiguity, we use distinct objects for losses, cascade events, and control costs in the graph system. Let

$$D_{i,t}^{\text{graph}} = \mathbf{1}_{\{\tau_i^{\text{graph}} \leq t\}}, \quad \Delta D_{i,t_m}^{\text{graph}} = D_{i,t_m}^{\text{graph}} - D_{i,t_{m-1}}^{\text{graph}},$$

where $0 = t_0 < t_1 < \dots < t_M = T$ is the numerical time grid. The type loss and aggregate loss of the graph system are defined by

$$L_t^{k,\text{graph}} = \frac{1}{N_k} \sum_{i:g(i)=k} D_{i,t}^{\text{graph}}, \quad L_t^{\text{graph}} = \sum_{k=1}^K \pi_k L_t^{k,\text{graph}} = \frac{1}{N} \sum_{i=1}^N D_{i,t}^{\text{graph}}. \quad (8)$$

Except for the size-distribution diagnostics specified separately in the external-data calibration section, all terminal losses reported in the main text are node default proportions as in (8). The cascade event is defined by

$$C^{\text{graph}} = \mathbf{1} \left\{ L_T^{\text{graph}} \geq \ell_{\text{cas}} \text{ or } \max_{1 \leq m \leq M} (L_{t_m}^{\text{graph}} - L_{t_{m-1}}^{\text{graph}}) \geq \delta_{\text{cas}} \right\}, \quad (9)$$

where ℓ_{cas} is the terminal system-loss threshold and δ_{cas} is the one-step new-loss threshold. If node-level controls $a_{i,t}$ are present, we denote the average control cost by

$$\mathcal{K}_T^{\text{graph}} = \frac{1}{N} \sum_{i=1}^N \int_0^T c_{g(i)}(a_{i,t}) dt \quad (10)$$

Thus C^{graph} always denotes a cascade event, not a control cost. The matrix approximation system defines L_t^{matrix} , C^{matrix} , and $\mathcal{K}_T^{\text{matrix}}$ in the same way, except that contagion shocks are generated by the type-average matrix $\hat{\Gamma}$.

We construct the type-aggregated matrix from the same sparse graph by

$$\hat{\Gamma}_{k\ell} = \frac{1}{n_k} \sum_{i:g(i)=k} \sum_{j:g(j)=\ell} A_{ij}. \quad (11)$$

This quantity represents the average weighted-edge exposure of a type k target node to type ℓ source nodes. Running (5) with $\hat{\Gamma}$ gives the finite-type matrix approximation of the same sparse graph. Since the graph system and the matrix system come from the same network, their difference isolates the information loss from aggregating local structure by type.

We define three matrix approximation gaps:

$$\text{Gap}_{\text{mean}} = \mathbb{E}[L_T^{\text{graph}}] - \mathbb{E}[L_T^{\text{matrix}}], \quad (12)$$

$$\text{Gap}_{\text{tail}} = Q_{0.95}(L_T^{\text{graph}}) - Q_{0.95}(L_T^{\text{matrix}}), \quad (13)$$

$$\text{Gap}_{\text{cascade}} = \mathbb{P}(C^{\text{graph}}) - \mathbb{P}(C^{\text{matrix}}), \quad (14)$$

where C denotes the cascade event in which terminal loss exceeds the threshold or one-step new loss exceeds the threshold. The tail gap is the main diagnostic, because local network structure can have a limited effect on mean loss while substantially changing tail paths.

The matrix-approximation theorem below uses two graph-level quantities that are computed directly from the adjacency matrix. The first is within-block row-exposure dispersion: nodes in the same target type should receive similar total exposure from each source type if the type matrix is to represent them accurately. The second is square-edge exposure spread: realized default-set fluctuations within a source type are small when exposure is distributed over many source nodes and large when a few edges dominate. In stochastic block networks these quantities are controlled by Bernoulli sampling and edge-weight moments. In core-periphery and heavy-tailed networks they become larger exactly when local concentration creates information that is lost under type aggregation.

3.3 Control objective and rule-based control

The continuous-time control problem of the central regulator can be written as

$$\inf_{a \in \mathcal{A}} \mathbb{E} \left[\Phi(L_T^1, \dots, L_T^K) + \int_0^T \sum_{k=1}^K \pi_k c_k(a_t^k) dt \right], \quad (15)$$

where Φ represents the terminal system-loss penalty, c_k is the control cost for type k , and $\mathcal{A}$ denotes type-level controls taking values in a compact control set; in the mean-field theory below these controls are common-noise adapted open-loop processes, while the finite sparse-graph experiments also compare node-level rule-based controls. The numerical part uses two complementary routes. The first route compares three interpretable control rules: the no-control rule serves as the benchmark; type-uniform control applies the same type-level control intensity to all nodes and represents a simple high-coverage regulatory intervention; graph-pressure threshold control triggers intervention according to node-level local pressure $P_{i,t}^{\text{graph}}$ or its smoothed version, and can therefore act on high-pressure nodes before local shocks are reflected in type-average losses. The second route solves a finite-dimensional projection HJB in the case $K = 2$, constructs feedback control consistent with the first-order condition in Theorem 6, and compares it with rule-based controls in the same forward particle system. The resulting control conclusions are organized as risk-cost frontiers and comparisons of information structures.

4 Mean-field limit and quantitative convergence

This section gives the mean-field foundation of the regularized default contagion model in the main text. We keep only theorem statements, proof sketches, and the explanations needed for the numerical design. Full technical proofs are given in Appendix [SM2](#).

Convention for the theoretical results The economic specification in (5) and in the kill-intensity equations above uses an exponential state-dependent kill intensity. All rigorous results in Sections 4–7 are stated for the corresponding bounded regularized killing specification: throughout these results we assume that there exists $\Lambda < \infty$ such that

$$0 \leq \lambda_k(x, a) \leq \Lambda, \quad |\lambda_k(x, a) - \lambda_k(y, a)| \leq L_\lambda |x - y|.$$

On any compact state-control domain $[x_{\min}, x_{\max}] \times [0, \bar{a}]$, the exponential specification satisfies these conditions with

$$\Lambda = \max_k \lambda_{\text{base},k} \exp\{\gamma_k(x_{b,k} - x_{\min})\},$$

as a direct consequence of the bounded-Lipschitz killing estimates in the appendix comparison-principle proof. In the finite-dimensional HJB computation, the state grid is compact and the cap can be chosen above the maximum exponential intensity attained on the grid. Proposition 4 and the appendix control proofs give the corresponding steep-killing bridge to absorbing-boundary contagion under nonsticky boundary crossing, absence of conditional atoms in the limiting default time, uniform steep-limit continuity of the driven default map, and local stability of the absorbing-boundary McKean–Vlasov fixed point.

Theorem 1 (multi-population McKean–Vlasov limit) Fix $T > 0$. Consider a system with K types of financial institutions, where type k has N_k particles and $N_* := \min_k N_k \rightarrow \infty$. Assume that initial states are independent and identically distributed within each type and have uniformly finite first moments, which are sufficient for the qualitative limit; the second-moment bound used for the quantitative rate is imposed separately in Theorem 2. Assume also that the drift b_k is Lipschitz in the state and measure variables, the kill intensity λ_k is bounded and Lipschitz, and the contagion matrix Γ is bounded. The type-level control $a_t = (a_t^1, \dots, a_t^K)$ takes values in a compact set and is progressively measurable with respect to the common-noise filtration $\mathbb{F}^{W^0}$; equivalently, after observing the common shock history, the regulator chooses one open-loop intensity for each type. Then the type-level surviving empirical measures and loss processes of the finite-particle system,

$$(\mu_t^{1,N}, \dots, \mu_t^{K,N}, L_t^{1,N}, \dots, L_t^{K,N}),$$

converge on $[0, T]$ to the multi-population McKean–Vlasov limit, whose uniqueness under the same bounded-killing regularity is established in Proposition 3

$$(\mu_t^1, \dots, \mu_t^K, L_t^1, \dots, L_t^K),$$

where the limiting state of each type is jointly determined by conditional sub-probability measures and loss feedback. More precisely, $\mu_t^{k,N}$ converges weakly to μ_t^k , and $L_t^{k,N}$ converges uniformly in probability on $[0, T]$.

The proof uses synchronous coupling: the finite particles and the limiting particles share initial states, idiosyncratic noises, common noise, and Poisson thinning random measures. The common-noise adapted control convention preserves conditional independence inside each type once the

common shock history is fixed. The drift error, killing error, and loss-feedback error are controlled by Lipschitz conditions, and the estimate closes by Gronwall's inequality. This result gives the finite-type matrix experiments a rigorous large-system limit foundation. The uniqueness of the limiting controlled system is established independently in Proposition 3, which proves well-posedness under the same bounded-killing and Lipschitz regularity.

Theorem 2 (quantitative convergence rate under explicit empirical-process inputs)

Under the conditions of Theorem 1, assume also that the system has a uniform second-moment bound, that the empirical measures of the conditionally independent limiting particles satisfy the standard bounded-Lipschitz quadratic fluctuation estimate, and that the synchronous Poisson-thinning coupling satisfies the corresponding metric-level quadratic bounded-Lipschitz stability estimate. This last estimate means that the bounded-Lipschitz distance between the finite surviving empirical measure and the paired limiting auxiliary empirical measure is controlled in mean square by the state, measure, and loss coupling errors, up to the usual N_k^{-1} martingale fluctuation term. Then there exists a constant $C_T < \infty$ such that, for each k ,

$$\sup_{t \in [0, T]} \mathbb{E} d_{\text{BL}}^2(\mu_t^{k, N}, \mu_t^k) + \sup_{t \in [0, T]} \mathbb{E} |L_t^{k, N} - L_t^k|^2 \leq \frac{C_T}{N_*}. \quad (16)$$

Therefore, the bounded-Lipschitz empirical-measure mean-square error and the loss-process mean-square error are $O(N_*^{-1})$, and the corresponding first-order errors are $O(N_*^{-1/2})$. The estimate is obtained by first working conditionally on the common-noise path and then integrating over the common noise; the displayed expectation is the resulting unconditional bound.

This rate follows by tracking constants in the synchronous coupling argument for Theorem 1 and by estimating the loss process through its Doob–Meyer martingale decomposition. Common noise changes only the conditional distribution and the constants; it does not change the empirical fluctuation order driven by the minimum type sample size. We state the quadratic bounded-Lipschitz empirical fluctuation input and the synchronous-thinning metric stability input explicitly because the killed-particle setting requires metric-level control of the bounded-Lipschitz supremum. Under the localized finite-entropy/Donsker condition recorded in Appendix SM2, these inputs hold with constants independent of the type sample size, and the displayed estimate is an unconditional consequence after integrating over the common noise. The empirical one-dimensional W_2^2 diagnostics reported below are numerical stability diagnostics consistent with the same first-order $N_*^{-1/2}$ particle scale; the theorem itself is stated in the bounded-Lipschitz metric, where the alive/dead mismatch created by killing is controlled at the mean-square level.

Remark (primitive sufficient condition for Theorem 2) The abstract inputs in Theorem 2 are verified under the localized finite-entropy condition for the killed-path bounded-Lipschitz class and the compact-state localization recorded in Lemma SM2.3 and Remark SM2.4 of Appendix SM2. In that primitive regime, the estimate in (16) follows from the bounded-killing, Lipschitz, and finite-entropy assumptions rather than from an additional rate postulate.

Proposition 1 (conditional same-type propagation of chaos) Under the conditions of Theorem 1, fix a type k , a time $t \in [0, T]$, and any fixed finite index set $\{i_1, \dots, i_m\} \subset \{i : g(i) = k\}$. Let $\hat{X}_i^{k, N}(t)$ denote the lifted state that equals $X_i^{k, N}(t)$ while the particle is alive and equals the cemetery state ∂ after default, and define $\hat{\mu}_t^k = \mu_t^k + L_t^k \delta_\partial$. Then, conditionally on the common-noise history,

$$\mathcal{L}(\hat{X}_{i_1}^{k, N}(t), \dots, \hat{X}_{i_m}^{k, N}(t) \mid \mathcal{F}_t^{W^0}) \Rightarrow (\hat{\mu}_t^k)^{\otimes m}$$

in probability. Equivalently, the finite collection becomes asymptotically independent after conditioning on the common noise. If $\sigma_0 = 0$, the random measure $\hat{\mu}_t^k$ is deterministic, and this conditional statement reduces to the standard unconditional propagation of chaos.

This proposition follows directly from Theorem 1, but it has an interpretive role in our setting: the finite-type matrix model averages individual heterogeneity within the same type, and its validity relies on this conditional exchangeability.

5 Matrix approximation error, tail gap, and spectral radius

This section formulates testable theoretical propositions for the relation between sparse graphs and finite-type matrices. Full proofs are given in Appendix SM3.

Theorem 3 (finite-grid absolute structural bound for the expected-loss gap of matrix approximation) Let $\hat{\Gamma}$ be the type-aggregated matrix of a sparse graph A , and run the graph system and the finite-type matrix system on the same fixed discrete grid $\Delta = \{0 = t_0 < \dots < t_M = T\}$. The two systems are synchronously coupled: they use the same initial states, Brownian increments, common-noise increments, and killing randomization variables. At each grid step, drift and diffusion are applied first, new defaults are sampled from the bounded state-dependent killing probability, and contagion shocks are then applied as one-step state jumps through the graph pressure $\sum_j A_{ij} \Delta N_{j,t_m}^{\text{graph}}$ in the graph system and through the matrix pressure $\sum_\ell \hat{\Gamma}_{g(i)\ell} \Delta L_{\ell,t_m}^{\text{matrix}}$ in the matrix system.

For i with $g(i) = k$, define the block row exposure

$$B_{i\ell} := \sum_{j:g(j)=\ell} A_{ij}, \quad \hat{\Gamma}_{k\ell} := \frac{1}{n_k} \sum_{i:g(i)=k} B_{i\ell}.$$

Define the row-dispersion and square-edge-spread diagnostics by

$$R_k(A) := \left[\frac{1}{n_k} \sum_{i:g(i)=k} \left(\sum_{\ell=1}^K |B_{i\ell} - \hat{\Gamma}_{k\ell}| \right)^2 \right]^{1/2},$$

$$S_k(A) := \frac{1}{n_k} \sum_{i:g(i)=k} \sum_{\ell=1}^K \left(\sum_{j:g(j)=\ell} A_{ij}^2 \right)^{1/2}.$$

Assume that the kill intensity is uniformly bounded and Lipschitz in the state variable, the drift update is Lipschitz in the state and measure variables, the weights are nonnegative, the total row exposures are uniformly bounded on the considered graph sequence, and the type shares satisfy $\pi_k \geq \underline{\pi} > 0$ uniformly over k and over the graph sequence. Assume also the following within-type sampling regularity: for every target node i , source type ℓ , and grid time t_m ,

$$\mathbb{E} \left[\left| \sum_{j:g(j)=\ell} A_{ij} (\Delta N_{j,t_m}^{\text{graph}} - \Delta L_{\ell,t_m}^{\text{graph}}) \right|^2 \right] \leq C_{\text{mix}} \sum_{j:g(j)=\ell} A_{ij}^2,$$

with C_{mix} independent of N . Then there exists a constant $C_{T,\Delta} < \infty$, depending only on T , the fixed grid through M and the grid-stability constants, the Lipschitz constants, the killing-intensity bound, the total exposure bound, C_{mix} , and $\underline{\pi}^{-1}$, such that

$$\left| \mathbb{E} L_T^{\text{graph}} - \mathbb{E} L_T^{\text{matrix}} \right| \leq C_{T,\Delta} \sum_{k=1}^K \pi_k (R_k(A) + S_k(A)). \quad (17)$$

The absolute value in (17) targets the magnitude of the graph–matrix expected-loss discrepancy; its sign is model-dependent. The finite-grid bound matches the jump-pressure update used in the graph and matrix systems. The term $R_k(A)$ measures heterogeneity of total type-block exposure among target nodes of the same type, while $S_k(A)$ measures the sampling volatility of realized default sets under edge concentration. The proof decomposes graph pressure minus matrix pressure into within-type default-set sampling, row-dispersion, and type-loss feedback terms, transfers these terms to state and killing-probability differences through Lipschitz stability, and closes the estimate by a discrete Gronwall argument. The constant is a fixed-grid constant tied to this jump-pressure discretization.

A share-controlled version follows directly. Define the maximum edge-share diagnostic

$$q_k(A) := \max_{i:g(i)=k, 1 \leq \ell \leq K, j:g(j)=\ell} \frac{A_{ij}}{B_{i\ell} \vee \varepsilon_0}, \quad \varepsilon_0 > 0.$$

If the block row exposures $B_{i\ell}$ are uniformly bounded, then $S_k(A) \leq C\sqrt{q_k(A)}$, and hence

$$\left| \mathbb{E}L_T^{\text{graph}} - \mathbb{E}L_T^{\text{matrix}} \right| \leq C_T \sum_{k=1}^K \pi_k(R_k(A) + \sqrt{q_k(A)}).$$

This corollary gives the economic reading of the theorem: matrix approximation is accurate when nodes within the same type have similar row-block exposures and source-type exposure is not concentrated in a few edges.

Scope of the structural bound When the within-type sampling regularity holds, Theorem 3 gives a certified finite-grid upper bound on the expected-loss discrepancy. In heavy-tailed, core–periphery, and other locally concentrated networks, $D_R(A)$ and $D_S(A)$ remain computable diagnostics for the row-dispersion and square-edge channels through which type aggregation loses information. The theorem identifies these structural mechanisms, and the simulations measure their realized size under strong local concentration.

The same discrete proof is stable under Euler-type scalings in which one-step default probabilities are of order Δt and the per-default shock scale $\alpha(\Delta t_m)$ satisfies boundedness of $\sum_m \alpha(\Delta t_m)\Delta t_m$ and $\sum_m \alpha(\Delta t_m)^2\Delta t_m$. A continuous-time refinement leads to a jump or measure-valued impulse formulation and is separated from the finite-grid theorem used here.

Theorem 4 (nonzero lower bound for the tail-loss gap in heterogeneous networks) There exist two sequences of weighted directed networks A_n^{hom} and A_n^{hub} with the same type assignment, the same aggregate matrix, the same type-wise initial-state distributions, the same killing parameters, and the same control rule, but A_n^{hub} contains locally concentrated hub exposure. Then there exist $\delta > 0$ and n_0 such that, for all $n \geq n_0$,

$$Q_{0.95}(L_T^{\text{hub}}) - Q_{0.95}(L_T^{\text{hom}}) \geq \delta. \quad (18)$$

Theorem 4 shows that local exposure concentration can create a strictly positive tail-loss gap even when the aggregate matrices are identical. The aggregate matrix therefore identifies average exposure but leaves the local shock distribution and tail cascade probability underdetermined once concentrated exposures are allowed. This result is the theoretical prototype for the sparse graph experiments in Sections 10–11.

The same construction embeds in any finite type system with at least two active types under the identical inactive-block or no-atom shifted-quantile conditions recorded in Appendix SM2ary

embedding result. Choosing one source type and one target type, setting all other type blocks identical across the two graph sequences, and keeping their terminal losses deterministic or common across the two systems preserves the same positive quantile separation after a harmless change in type shares. Thus the two-type construction is the minimal version of a general finite-type information-loss mechanism.

The separation condition has an explicit feasible parameter region. For example, take the core-type share $\pi_1 = 0.05$ and the periphery-type share $\pi_2 = 0.95$. Let the additional shock-probability difference from the core hub to periphery nodes be $p_+ - p_- = 0.05$, and take the initial perturbation cost of the first type to be $\rho = 0.08$. Then

$$\pi_2(p_+ - p_-) - \pi_1\rho = 0.95 \times 0.05 - 0.05 \times 0.08 = 0.0435 > 0.$$

Under the exponential kill intensity $\lambda_0(x) = \lambda_{\text{base}} \exp\{\gamma(x_b - x)\}$, the stated $p_+ - p_-$ can be generated by pushing the post-hub-shock state near the boundary while the uniform shock with the same aggregate exposure remains outside the boundary, and by increasing γ or the near-critical exposure multiplier. The construction is taken on a discrete-time sampling convention in which first-stage defaults are sampled before the contagion shock of that step is fed back into later kill probabilities; choosing the first window sufficiently short keeps earlier feedback effects of order Δt and preserves the separation. The lower bound in Theorem 4 is therefore realized in an explicit transition region with feasible parameters.

Theorem 5 (local spectral-radius threshold for low-loss cascade perturbations) Let Γ be a nonnegative irreducible contagion matrix and let $Y_m \in \mathbb{R}_+^K$ denote a low-loss perturbation vector on a fixed local time grid. Set $M = \chi\Gamma$, where $\chi > 0$ is the local sensitivity coefficient determined by killing sensitivity and drift damping. Assume the following low-loss linearization conditions. First, while $\|Y_m\| \leq \eta$,

$$\mathbb{E}[Y_{m+1} \mid \mathcal{F}_m] = MY_m + R(Y_m) + \varepsilon_m^N, \quad \|R(y)\| \leq C_q \|y\|^2,$$

and the finite-sample residual has conditional second moment bounded by C/N in the same region. Second, the initial seed has a nondegenerate positive projection on the Perron–Frobenius left eigenvector of M . Third, the seed size, deterministic residual, and martingale fluctuation satisfy the logarithmic-window domination condition from the appendix proof: on the hitting window considered below, the quadratic remainder and finite-sample residual are smaller than the linear Perron–Frobenius gap generated by the initial seed.

If $\rho(M) < 1$, then every sufficiently small positive seed satisfying the low-loss condition decays on this logarithmic window in the Perron–Frobenius direction and remains in the low-loss region with high probability. If $\rho(M) > 1$, then there exists $\varepsilon_* > 0$ such that, for every local cascade threshold $\varepsilon \in (0, \varepsilon_*)$, every positive seed with nondegenerate Perron–Frobenius projection and the stated finite-sample compatibility reaches level ε with probability tending to one before leaving the low-loss linearization region. Equivalently, when the local effective feedback strength is written as

$$R_{\text{eff}} = \rho(\Gamma)\chi,$$

the boundary $R_{\text{eff}} = 1$ is the linear critical boundary for sufficiently small local cascade perturbations.

Remark on the finite-sample compatibility in Theorem 5 The logarithmic-window condition is the finite-sample requirement that makes the probability statement meaningful. In the formulation recorded in the appendix proof, the deterministic residual and martingale fluctuation scale must be

small relative to the Perron–Frobenius growth generated by the initial seed before the trajectory reaches the local threshold. A representative compatible regime has residual order $O(N^{-1})$ and seed size $\|Y_0^N\|_1 = N^{-\alpha}$ with $0 < \alpha < 1$, together with the corresponding martingale domination on the same window.

This is a local theorem, not a global cascade theorem. The assumptions above are the main-text version of the low-loss expansion, finite-sample fluctuation control, nondegenerate Perron–Frobenius seed, and logarithmic-window domination conditions proved and interpreted in the appendix approximation proofs. The result explains why we use $\rho(\Gamma)$ as a diagnostic for contagion amplification in the low-loss linearized region. The spectral radius organizes local amplification, while the graph diagnostics in Theorem 3 describe the within-type information that remains invisible after type aggregation. The numerical sparse-graph experiments combine these two diagnostics.

Proposition 2 (continuous dependence of loss processes on the contagion matrix)

Consider two systems that share the same initial distribution, killing parameters, drift, and noise structure, and differ only in their contagion matrices Γ and Γ' . Suppose $\|\Gamma\|_\infty \vee \|\Gamma'\|_\infty \leq \bar{\Gamma}$. Then there exists a constant $C_T = C(T, \Lambda, L_b, L_\lambda, \bar{\Gamma})$ such that, for each type k ,

$$\mathbb{E} \sup_{t \in [0, T]} |L_t^k(\Gamma) - L_t^k(\Gamma')| \leq C_T \|\Gamma - \Gamma'\|_\infty.$$

Equivalently, the same Lipschitz estimate holds almost surely after conditioning on a fixed common-noise path whenever the conditional version of the coupled systems is used. This proposition ensures that the risk–cost frontiers in the finite-type matrix experiments are not unstable artifacts caused by small numerical perturbations of matrix entries inside a bounded contagion-matrix class.

6 Controlled contagion systems and optimal control characterization

This section gives the well-posedness of the controlled model, the optimal control characterization for two populations, and the relation between the regularized killing mechanism and the hard absorbing boundary. Proposition 4 is a singular-limit statement and is separate from the bounded-killing framework used in the main limit and control theorems. Full proofs are given in Appendix SM4.

Proposition 3 (well-posedness of the controlled multi-population contagion system)

Assume that the drift is Lipschitz in the state and measure variables with linear growth, the kill intensity is bounded and Lipschitz in the state variable, the control sets are compact, the contagion matrix is bounded, the diffusion coefficients are bounded, and the initial states have the required first or second moments. Then, for any common-noise adapted admissible type-level control, the controlled multi-population McKean–Vlasov contagion system has a unique solution. The loss process is a càdlàg nondecreasing process, and the solution is stable with respect to the initial distribution and the control flow.

This proposition also supplies the uniqueness input invoked in Theorem 1. The proof constructs a map from measure flows and loss flows to themselves. The map is closed by a contraction argument on short time intervals, and a concatenation argument then gives a global solution on $[0, T]$. This proposition ensures that the optimization object in (15) is well defined.

Theorem 6 (existence of two-population optimal control and conditional common-noise HJB characterization) Consider the controlled contagion system with $K = 2$. The type-level

control takes values in the compact set $[0, \bar{a}]^2$, the kill intensity is bounded and Lipschitz on the compact state-control domain, the controlled martingale problem is well posed, the idiosyncratic alive-state diffusion coefficients satisfy $\sigma_k > 0$, the cost functions c_k are continuous and convex on compact control sets, and the terminal loss penalty $\Phi : [0, 1]^2 \rightarrow \mathbb{R}$ is bounded uniformly continuous; in the numerical quadratic-loss specifications it is bounded automatically because $(L_T^1, L_T^2) \in [0, 1]^2$. Then, in the presence of common noise, the central control problem admits an optimal relaxed control. The compact relaxed-control formulation gives the HJB value. The killed-HJB comparison principle is proved in the appendix comparison-principle proof on the decomposed state space of alive sub-probability measures and cemetery masses, which is identified with $\mathcal{M}_{\leq 1, 2}(\mathbb{R})^2 \times [0, 1]^2$ subject to $|\nu^k| + L^k = 1$. The alive-measure variables are handled by the Wasserstein smooth-gauge/Ishii template, the cemetery masses are handled by finite-dimensional doubled-variable penalties, and the killing jump becomes a bounded Lipschitz first-order perturbation from alive mass to cemetery mass. Under the standard chattering and measurable-selection conditions recorded in the appendix control proofs, the relaxed value agrees with the strict-control value whenever the Hamiltonian minimizer admits a progressively measurable strict selector. With this convention, the value function is the unique bounded uniformly continuous viscosity solution of the corresponding decomposed measure-valued HJB equation, equivalently of the lifted killed-state HJB.

The preceding viscosity characterization does not require classical differentiability of the value function. The first-order condition below is a smooth-verification diagnostic rather than an additional viscosity-uniqueness claim. When, in addition, the value function admits a classical verification on a smooth region of the lifted measure space, so that the Lions derivative with respect to μ_k and the corresponding cemetery-jump marginal value are well defined, the Hamiltonian minimizer can be characterized by a first-order condition. In the finite-dimensional projected HJB used numerically, this smoothness requirement can be checked locally by standard parabolic PDE regularity on compact grids when the projected diffusion in the alive coordinates is nondegenerate and the coefficients are bounded Lipschitz; the paper uses the resulting formula only as a Hamiltonian diagnostic. For a quadratic running cost $c_k(a) = \frac{1}{2}r_k a^2$ appearing in the population-weighted objective $\sum_k \pi_k c_k(a^k)$, this smooth-verification condition has the integrated form

$$a_k^*(t) = \Pi_{[0, \bar{a}_k]} \left(\frac{\eta_k}{\pi_k r_k} \int_{\mathbb{R}} \lambda_k(x, a_k^*(t)) \mathcal{M}_k(t, x, \boldsymbol{\mu}_t) \mu_t^k(dx) \right),$$

where $\mathcal{M}_k(t, x, \boldsymbol{\mu}_t)$ denotes the complete marginal value of suppressing a type- k killing event at state x , including both the direct cemetery-jump value and the induced loss-feedback value transmitted through the contagion matrix. In the finite-dimensional projection used in the numerical section, this smooth-verification formula is used as a Hamiltonian diagnostic. The alive distribution of each type is represented by the representative coordinate (x_1, x_2, L_1, L_2) , and the integrated condition reduces formally to the pointwise projected fixed-point feedback condition

$$a_k^*(t, x_1, x_2, L_1, L_2) = \Pi_{[0, \bar{a}_k]} \left(\frac{\eta_k \lambda_k(x_k, a_k^*) \mathcal{M}_k(t, x_1, x_2, L_1, L_2)}{\pi_k r_k} \right).$$

Under the stated comparison, relaxed-control, and selector inputs, this theorem places the rule-based controls used in this paper within a stochastic control framework. In the numerical section, we solve the finite-dimensional projection HJB for $K = 2$ and compare the resulting feedback control with rule-based controls. This projection is the computational counterpart of the measure-valued characterization. The comparison principle is proved for the bounded regularized killing specification with $K = 2$; extension to general K requires a K -fold product Wasserstein smooth-gauge construction and a complete doubled-variable maximum argument for all cross-type terms. The numerical HJB benchmark is correspondingly kept at $K = 2$, a natural core-periphery

partition in which systemically important institutions and other banks form the two interacting populations.

Proposition 4 (steep killing and absorbing-boundary default) The regularized killing mechanism used in the paper has a steep-intensity limit that selects an absorbing-boundary model. The appendix control proofs give a primitive verification of the bridge under a localized one-dimensional boundary-regularity package: compact localization, bounded Lipschitz frozen coefficients, nondegenerate idiosyncratic diffusion near the boundary, uniform killed-diffusion boundary estimates, and an absorbing no-atom modulus. The appendix proof verifies localized uniqueness of the absorbing fixed point by a Volterra short-time contraction and concatenation argument, derives eventual entrance into a stability neighborhood, proves local stability of the absorbing fixed point, and obtains uniform convergence of the driven steep-killing maps on compact localized classes. The limiting object depends on the baseline intensity above the boundary, which makes the proposition a precise bridge between regularized killing and the absorbing-boundary convention used in hard-default models.

If

$$\lambda_0^{(n)}(x) = \lambda_{\text{base}} \exp(n(x_b - x)^+),$$

then the selected limit is absorbing-boundary default together with the baseline Cox killing that remains active while $x \geq x_b$. If the intended limiting object is a pure hard absorbing boundary with no baseline Cox killing above the boundary, one can use

$$\tilde{\lambda}_0^{(n)}(x) = \lambda_{\text{base}} \{\exp(n(x_b - x)^+) - 1\},$$

or let the above-boundary baseline intensity vanish with n . The numerical experiments use the bounded regularized state-dependent killing mechanism directly; this singular-limit proposition identifies the absorbing-boundary model selected by a separate steep-killing limit.

7 Information value of local pressure information

Theorem 7 (variational sufficient condition for positive value of local pressure information) Let $\mathbb{F}^{\text{type}}$ be the filtration generated by type-average losses and type-average pressures, and let $\mathbb{F}^{\text{graph}}$ be the filtration that further includes node-level local pressure $P_{i,t}^{\text{graph}}$. Let V^{type} and V^{graph} denote the minimal costs when admissible controls are adapted to these two filtrations under the same running-cost budget.

Assume that there is a type k , a stopping time τ , and an event $\mathcal{E} \in \mathbb{F}_{\tau}^{\text{graph}}$ with $\mathbb{P}(\mathcal{E}) > 0$ on which the node-level graph pressures of surviving type- k nodes have nondegenerate within-type variance. The comparison is made on a strong construction in which the baseline type-level optimizer and the graph-pressure reallocated control share the same Brownian motions and killing randomization variables; the reallocated control agrees with the baseline control up to τ , so $\mathcal{E}$ and the pressure vector used for reallocation are observed under both controls at the intervention time. Writing

$$\bar{P}_{k,\tau} := \frac{1}{N_k} \sum_{i:g(i)=k} P_{i,\tau}^{\text{graph}}, \quad Z_i := P_{i,\tau}^{\text{graph}} - \bar{P}_{k,\tau},$$

assume that $S_k := N_k^{-1} \sum_{i:g(i)=k} Z_i^2 \geq v_k > 0$ on $\mathcal{E}$, and that the type-level optimizer is interior on this event. Assume also that the conditional terminal-risk functional is twice Gateaux differentiable along budget-preserving node-level reallocations in the centered pressure direction Z . Finally, assume

a strict marginal-benefit condition: there exists $m_k > 0$ such that reallocating an infinitesimal unit of control toward nodes with larger Z_i lowers the conditional terminal cost by at least $m_k S_k$ up to the stated type-share and active-window constants, and the second-order remainder is bounded by $C\varepsilon^2 S_k$. Then, for sufficiently small $\varepsilon > 0$, the reallocated control is feasible, preserves the running-cost budget, and strictly lowers the conditional objective on $\mathcal{E}$. Consequently there exists $\delta_{\text{info}} > 0$ such that

$$V^{\text{type}} - V^{\text{graph}} \geq \delta_{\text{info}}. \quad (19)$$

The theorem is a sufficient condition for positive information value. The proof constructs a graph-measurable reallocation that shifts intervention from low-pressure nodes to high-pressure nodes while preserving the first-order control budget. The strict marginal-benefit condition turns this reallocation into a strict reduction in terminal loss risk. The appendix control proofs verify primitive pressure monotonicity for exponential killing in a nonsaturated local window, showing how the marginal-benefit constant is generated by the model. They also record a sufficient smoothness regime for the twice Gateaux differentiability input: smooth bounded drift and killing coefficients, bounded smooth terminal loss penalty, compact controls away from saturation corners, and a local window on which conditional killing compensators have uniformly bounded derivatives. In saturated regions, the forward experiments complement the variational lower bound by reporting realized risk-cost comparisons. The graph-pressure threshold control experiments in Section 11 are the numerical counterpart of this variational mechanism.

Corollary 1 (primitive pressure-monotonicity window for Theorem 7) Suppose the kill intensity has the exponential form $\lambda_k(x, a) = \lambda_{0,k}(x) \exp\{-\eta_k a\}$, the relevant stopping-time event lies in a nonsaturated window in which controls are interior and the post-pressure state remains inside a compact neighborhood of the boundary, and the second-order feedback terms along centered pressure reallocations are bounded. Then the marginal-benefit condition in Theorem 7 holds with a positive constant determined by the killing sensitivity η_k , the local baseline intensity, the drift damping, and the active-window survival mass. Consequently, whenever the within-type local-pressure variance is nondegenerate on such a window, node-level graph-pressure information has strictly positive value relative to type-average information. The appendix proof gives the corresponding pressure-monotonicity calculation and the explicit second-order remainder bound.

8 Numerical design

The finite-type experiments use four classes of institutions: small banks, regional banks, major banks, and systemically important banks. The moderate-stress specification sets the baseline kill-intensity multiplier to 2.5 and shifts the initial state downward by -0.25 . The matrix scalings are 45, 48, 50, the particle number is 10000, the number of time steps is 500, and the number of replications is 200. This specification is used to display near-critical loss growth and the risk-cost frontier in the matrix system. The convergence validation experiment changes the particle number under the same moderate-stress structure and estimates the reference loss and reference empirical distribution using a large-sample system.

The sparse graph experiments examine three network classes. The stochastic block network serves as the regular-mixing benchmark, with edge probabilities and edge weights mainly determined by type pairs. The core-periphery network increases the connection strength and spillover exposure of the core type, and is used to represent locally concentrated risk around systemically important nodes. The heavy-tailed network introduces source-node fitness and hub exposure multipliers, and is used to represent cascade risk driven by highly connected nodes. Each graph is aggregated into a

matrix approximation through (11), and the graph system and matrix system are compared under the same default mechanism.

Sparse graph exposure intensity is amplified by a common multiplier x , with scan range $x = 1, \dots, 30$. The representative control experiments choose three points: the heavy-tailed network at $x = 12$, the core-periphery network at $x = 15$, and the stochastic block network at $x = 26$. At these points, the true graph systems already exhibit significant cascades, while the corresponding aggregate matrix approximations have not entered the cascade region at the same time.

The synthetic numerical experiments use unified inputs and outputs. Inputs include type shares π_k , the initial safety-margin distribution, drift and volatility parameters, kill-intensity parameters, the contagion matrix or sparse adjacency matrix, common-noise intensity, control rules and their budgets or thresholds, the time grid, the particle number, and the number of Monte Carlo replications. The finite-type matrix experiments specify Γ directly. The sparse graph experiments first generate A and then obtain the paired matrix $\hat{\Gamma}$ from (11). Outputs are unified as terminal mean loss, 95% tail loss, cascade probability, control cost, risk reduction per unit cost, and the mean and tail gaps between the graph system and the matrix system. This organization allows different network families, control rules, and parameter scans to be compared under the same set of risk-cost metrics, rather than treating each experiment as an independent simulation case.

To test the consistency between the theoretical results and numerical mechanisms, we include several closed-loop experiments. First, we vary the particle number N and estimate the reference loss using a large-sample system, and we use mean-square error to validate the $N_*^{-1/2}$ particle scale in the conditional quantitative rate of Theorem 2. Since the figure reports MSE, the corresponding reference slope is -1 , and we report the log-log regression slope, standard error, R^2 , and significance level. Second, we scan the common-noise intensity σ_0 to study how common shocks amplify terminal loss, tail gaps, and local control value. Third, we compute the row-dispersion component $R_k(A)$ and the square-edge-spread component $S_k(A)$, and compare the resulting matrix-approximation diagnostic with realized graph-matrix gaps. Fourth, we rescale the three network classes to the same aggregate spectral radius to isolate the contribution of network heterogeneity itself to the matrix approximation gap. Fifth, we compare type-uniform control and graph-pressure threshold control again under a fixed total budget, so that the risk-cost comparison does not depend on cost differences naturally generated by different strategies. Sixth, we densely sample the transition region $x = 8, \dots, 16$ and use high-replication Monte Carlo estimation to report cascade probabilities, tail gaps, and confidence intervals. Seventh, we add time-varying networks, extra degree-distribution networks, and finite-type matrix experiments with $K = 3, 4$ to test the robustness of the conclusions to network dynamics, degree distributions, and the dimension of the population partition.

The two-population optimal control experiment uses a core-periphery structure. The core type has share 5%, and the periphery type has share 95%. The control computation solves a finite-dimensional projection HJB on (x_1, x_2, L_1, L_2) , and compares HJB feedback, type-uniform control, graph-pressure threshold control, and no control in the same forward particle system. The HJB calibration selects a nondegenerate near-critical region in which the uncontrolled system exhibits clear cascades and the optimal feedback has nonzero intervention intensity at comparable cost. The numerical discretization uses a semi-implicit diffusion step for the second-order terms in the x direction, retains the common-noise cross term, and uses a one-way semi-Lagrangian loss jump with monotone piecewise-linear interpolation in the loss coordinates to handle the advance of the killing term in the L coordinate. We then solve the system repeatedly at four resolutions, $N_x = 20, 40, 80, 160$, and perform forward validation for each feedback. We report grid-by-grid changes in the value function, control field, and cascade probability. The killing-parameter sensitivity experiment scans $\gamma \in \{5, 10, 15, 20\}$ and $\lambda_{\text{base}} \in \{0.01, 0.03, 0.05, 0.10\}$ to test how the soft killing

mechanism changes tail loss levels in high-pressure and near-saturation regions.

The external-data calibration experiments use two categories of public regulatory data. The first is the EBA 2024 EU-wide Transparency Exercise data, used to extract bank-level asset-size proxies, risk-exposure-amount proxies, institutional jurisdictions, and country-level credit-exposure proxies. The second is the official reporting package from the EBA Pillar 3 Data Hub / European Data Access Portal, used to record institution-level regulatory disclosure modules and reporting-file structures. These public data constrain bank size distributions, risk-exposure proxies, jurisdictional structure, and disclosure coverage. We use them to construct EBA and Pillar 3 calibrated synthetic core-periphery networks, with bilateral edges generated inside those public constraints. The jurisdictional distribution is visualized using a public country-boundary base map for geographic presentation. In addition to a single representative network, we fix the public-data constraints and independently generate multiple calibrated synthetic networks, using the cross-instance distribution to test the stability of the control conclusions across unobserved bilateral edge realizations.

9 Finite-type matrix experiments

The finite-type matrix system exhibits a moderate but clear near-critical transition under the moderate-stress specification. Without control, as the matrix scaling increases from 45 to 50, the terminal mean loss, tail loss, and cascade probability increase together. Figure 1 reports this transition. The result shows that the contagion matrix is not a mere calibration parameter, but a structural variable that organizes systemic loss paths.




Figure 1: Finite-type matrix scaling and risk-cost frontier in the near-critical region.

The control experiments show that different rules form a risk-cost frontier in the same near-critical region. Risk-threshold control gives the lowest terminal loss and tail loss, but has the highest control cost. Uniform control is a strong benchmark. Type-priority control and contagion-pressure threshold control leave more residual loss but require lower intervention cost. Figure 1 shows that the control conclusion is not global dominance of one strategy over all others. Different information structures and control intensities correspond to different positions on the frontier.

The quantitative convergence result in Theorem 2 is numerically tested in Figure 2. The horizontal axis is the minimum particle number across types, N_* . The vertical axes are the terminal loss MSE and the one-dimensional empirical W_2^2 diagnostic of the lifted empirical measure. The theorem itself is stated in the bounded-Lipschitz metric, while the empirical W_2^2 panel is reported as a numerical stability diagnostic for the same first-order particle scale. The curves for all types have the same order as the C_T/N_* reference line, showing that the finite-particle system is consistent with

the $N_*^{-1/2}$ error scale predicted by the theory. Table 1 reports log-log slopes and fitted constants. Slopes near -1 correspond to the $N_*^{-1/2}$ rate at the RMSE level in Theorem 2.




Figure 2: Particle-number convergence validation for Theorem 2.

Table 1: Log-log convergence regressions for Theorem 2.

Type	Metric	Slope	SE	R^2	C_T
major bank	$E L_T^{k,N} - L_T^k ^2$	-1.730	0.234	0.901	3.399
major bank	$EW_2^2(\hat{\mu}_T^{k,N}, \hat{\mu}_T^k)$	-1.053	0.187	0.840	554.6
regional bank	$E L_T^{k,N} - L_T^k ^2$	-1.455	0.183	0.913	3.766
regional bank	$EW_2^2(\hat{\mu}_T^{k,N}, \hat{\mu}_T^k)$	-0.974	0.164	0.855	470.8
small bank	$E L_T^{k,N} - L_T^k ^2$	-1.329	0.160	0.920	3.648
small bank	$EW_2^2(\hat{\mu}_T^{k,N}, \hat{\mu}_T^k)$	-0.922	0.150	0.864	369.6
systemically important bank	$E L_T^{k,N} - L_T^k ^2$	-1.902	0.266	0.895	3.149
systemically important bank	$EW_2^2(\hat{\mu}_T^{k,N}, \hat{\mu}_T^k)$	-1.123	0.188	0.856	527.4

10 Matrix approximation boundary for sparse graphs

Figure 3 first gives structural diagnostics for the three sparse network classes. The adjacency sparsity pattern of the stochastic block network is close to within-block uniform sampling. The core-periphery network forms denser local connections around core targets and core sources. The heavy-tailed network generates stronger node-level concentration through hub exposure and the upper tail of weighted in-exposure. Figure 4 shows the same differences through representative subgraph layouts. These figures are not additional modeling assumptions. They provide structural background for the matrix approximation gaps studied below: the aggregate matrix preserves average exposure across types, but compresses within-block row-exposure dispersion, edge concentration, and local neighborhood shape.

Figure 5 reports cascade probabilities for the three sparse network classes under the exposure multiplier sweep. The heavy-tailed network enters the graph cascade region first, followed by the core-periphery network, while the stochastic block network enters last. In these near-critical regions, the corresponding aggregate matrix approximations still have low cascade probabilities. Cascades in




Figure 3: Adjacency sparsity patterns and weighted in-exposure distributions for the three sparse network classes. The bottom-row histograms use a common horizontal scale, so one unit has the same exposure meaning across networks.




Figure 4: Representative subgraph layouts for the three sparse network classes. Node positions are used only to display local connection patterns and do not enter the risk computation.

the true local networks are therefore not fully explained by type-average exposure. They are jointly driven by local exposure concentration, hub nodes, and core neighborhoods.




Figure 5: Sparse graph systems versus aggregate matrix approximations under exposure scaling.

The tail-loss gap is clearer in Figure 5. As the exposure multiplier increases, the tail-loss gap of the graph system relative to the matrix approximation expands rapidly. The gap appears earlier in the heavy-tailed and core-periphery networks, showing that locally concentrated exposure can worsen tail paths before aggregate matrix risk indicators raise an alarm.

Figure 6 converts this pattern into critical exposure multipliers and includes Barabasi-Albert scale-free networks and Watts-Strogatz small-world networks in the same comparison. The heavy-tailed network requires the lowest multiplier to reach a 10% graph cascade probability, and the core-periphery network enters the critical region next. The small-world network, stochastic block network, and scale-free network require higher exposure multipliers to reach the same threshold. This five-network ordering shows that local exposure concentration, core neighborhoods, and clustering structure all shift the systemic cascade boundary, while the type-aggregated matrix cannot fully capture these differences from average exposure alone.




Figure 6: Exposure multipliers at which different network structures reach the graph cascade threshold.

Three representative near-critical points make the approximation boundary concrete. In the heavy-tailed network at multiplier 12, graph tail loss is 0.3637 and matrix tail loss is 0.0798, giving a tail gap of 0.2838, while graph cascade probability is 0.5000 and matrix cascade probability remains 0. In the core-periphery network at multiplier 15, the corresponding tail gap is 0.5373 and graph cascade probability is 0.4800; in the stochastic-block network at multiplier 26, the tail gap is 0.2914 and graph cascade probability is 0.4000. These three points are the main evidence for the approximation boundary: the type-aggregated matrix of the same graph can preserve average exposure, but it cannot fully preserve the tail paths generated by locally concentrated exposure.

To test whether the transition ordering depends on Monte Carlo error, we re-estimate the three network classes with 500 replications on the key interval $x = 8, \dots, 16$. Figure 7 gives confidence bands for graph-system cascade probabilities and tail gaps. The refined estimates preserve the same ordering: the heavy-tailed network first enters the cascade region at exposure multiplier 11 with maximum tail gap 0.796 and maximum cascade probability 1.000; the core-periphery network reaches the 10% cascade threshold at multiplier 14 with maximum tail gap 0.544 and maximum cascade probability 0.940; and the stochastic block network still does not reach a 10% cascade probability on this interval, with maximum tail gap only 0.037. The differences across the three network classes are therefore stable transition differences caused by local exposure concentration, rather than sampling fluctuations from a small number of replications.




Figure 7: Confidence bands for cascade probabilities and tail gaps with 500 replications in the exposure multiplier transition region.

The matrix approximation error bound in Theorem 3 is controlled by row-dispersion and square-edge-spread quantities. In the numerical diagnostics, we use

$$D_{T3}(A) = D_R(A) + D_S(A), \quad D_R(A) = \sum_k \pi_k R_k(A), \quad D_S(A) = \sum_k \pi_k S_k(A),$$

where $R_k(A)$ and $S_k(A)$ are the two structural terms in Theorem 3. Figure 8 compares $D_{T3}(A)$ with the realized gap $|\mathbb{E}L_T^{\text{graph}} - \mathbb{E}L_T^{\text{matrix}}|$ across exposure multipliers. The fitted upper-bound line uses a single conservative coefficient multiplying $D_{T3}(A)$. The decomposition shows how row-dispersion and square-edge spread jointly raise the graph-matrix gap in the heterogeneous network classes.

At the largest realized mean-gap points, the heavy-tailed network has multiplier 19, mean gap 0.8246, tail gap 0.8111, $D_R(A) = 33.0271$, $D_S(A) = 25.6111$, $D_{T3}(A) = 58.6381$, and fitted bound 0.9049. The core-periphery network has multiplier 24, mean gap 0.8390, tail gap 0.8154,




Figure 8: Theorem 3 diagnostics for the matrix-approximation boundary.

$D_R(A) = 31.3849$, $D_S(A) = 27.4080$, $D_{T3}(A) = 58.7929$, and fitted bound 0.9073. The stochastic-block network reaches its largest mean gap at multiplier 30, where the mean gap is 0.3973, the tail gap is 0.5228, and $D_{T3}(A) = 41.9001$. These values show that the structural diagnostic is conservative in level but separates the high-gap heterogeneous networks from the more regular stochastic-block benchmark. For the core-periphery and heavy-tailed graphs, the plotted $D_{T3}(A)$ is a conditional structural diagnostic under the within-type sampling regularity in Theorem 3.

We also rescale the three network classes to the same aggregate spectral radius $\rho(\hat{\Gamma}) = 5$ to separate total contagion amplification from local aggregation error. Figure 8 shows that, under this common aggregate-amplification scale, the realized gaps in this calibration become small and comparable. The mean gaps are 0.0055 for the heavy-tailed network, 0.0048 for the core-periphery network, and 0.0054 for the stochastic-block network; the corresponding tail gaps are 0.0063, 0.0061, and 0.0069. The fixed-spectral-radius diagnostics are also close, with $D_{T3}(A) = 9.7896$, 8.5289, and 9.8056, respectively. This comparison first normalizes aggregate feedback strength and then reads the residual differences through the local row-dispersion and square-edge-spread channels in Theorem 3.

The common-noise experiment brings the case $\sigma_0 > 0$ from the theory into the sparse graph numerical analysis. Figure 9 shows that, with 500 replications, terminal tail loss and the matrix approximation gap rise overall as common-noise intensity increases, especially around $\sigma_0 = 0.15$ and 0.20. Across $\sigma_0 \in \{0, 0.05, 0.10, 0.15, 0.20\}$, no-control 95% loss ranges from 0.385 to 0.448, graph-pressure 95% loss ranges from 0.238 to 0.297, and cascade reduction per unit cost remains between 16.41 and 20.77. Common shocks move the safety margins of different nodes in the same direction and therefore amplify synchronization in local network pressure, but they do not eliminate the value of local pressure information.

11 Graph-pressure threshold control and control transfer

The matrix approximation boundary itself concerns risk identification. The next question is whether local pressure information in the sparse graph can support more effective control when the aggregate matrix misses local cascades. We therefore compare no control, type-uniform control, and graph-pressure threshold control at three representative points. Graph-pressure threshold control triggers intervention according to node-level local pressure in (6), and can therefore react to locally high-pressure nodes.

With 500 replications, the no-control graph cascade probabilities are 0.762 for the heavy-tailed network, 0.736 for the core-periphery network, and 0.658 for the stochastic-block network, while




Figure 9: Common-noise effects on sparse graph risk and graph-pressure control.

the matrix approximation cascade probabilities remain zero. Type-uniform control eliminates the observed graph cascades, but its cost is about 0.259. Graph-pressure threshold control leaves a small amount of residual risk, but reduces the cascade probabilities to 0.150, 0.116, and 0.074 at about one tenth of the cost. Type-average control is a strong suppression benchmark, while graph-pressure threshold control is a low-cost local intervention rule.

Figure 10 shows the same comparison visually. Under no control, the representative points for the heavy-tailed, core-periphery, and stochastic-block networks all have high cascade probabilities. Graph-pressure threshold control lowers these cascade probabilities substantially, and type-uniform control lowers them further to zero.




Figure 10: Control strategies and cascade probabilities in representative sparse graph regions.

The cost-efficiency comparison is more compactly summarized in text. For the heavy-tailed, core-periphery, and stochastic-block networks, graph-pressure threshold control achieves cascade reductions per unit cost of 24.21, 23.38, and 24.05, respectively, whereas type-uniform control

achieves only 2.94, 2.84, and 2.54. In regions where matrix approximation fails, local pressure information obtains substantial cascade reduction at much lower intervention cost, while uniform control remains the stronger high-cost suppression benchmark.

Figure 11 gives the risk–cost relation from the perspective of tail loss. Uniform control lies at the low-risk, high-cost end, while graph-pressure threshold control lies at the low-cost end with higher residual risk. This figure mirrors the risk–cost frontier in the finite-type matrix experiments and shows that the same frontier structure persists in sparse graph local contagion systems.




Figure 11: Tail risk–control cost relation in representative sparse graph regions.

Because type-uniform control and graph-pressure threshold control naturally operate at different cost levels, we also repeat the comparison under fixed budgets. Figure 12 gives the cascade probability frontier under budget constraints. Graph-pressure threshold control usually achieves substantial cascade reduction at lower cost in the low-budget region. As the budget increases, uniform control gradually approaches the strong-suppression end. The representative budgets confirm this pattern numerically.

At budget 0.030, graph-pressure threshold control has lower cascade probability than type-uniform control in all three network classes: 0.060 versus 0.140 in the core–periphery network, 0.125 versus 0.170 in the heavy-tailed network, and 0.035 versus 0.065 in the stochastic-block network. The realized-cost and tail-loss comparison has the same shape: graph-pressure threshold control stays in the low-cost range 0.027–0.044 with 95% loss about 0.186–0.220, while type-uniform control moves from low-budget partial suppression to high-budget strong suppression, reaching realized cost 0.151 and 95% loss around 0.085–0.088 at budget 0.200. At budgets 0.100 and 0.200, type-uniform control reaches the strong-suppression regime with zero observed cascade probability in most cases and substantially larger realized costs, while graph-pressure control remains a low-cost frontier point with positive residual risk.

12 Two-population optimal control numerical loop

Theorem 6 gives the HJB characterization, within the lifted comparison framework, and the smooth-verification first-order condition for the controlled McKean–Vlasov system with $K = 2$ under common noise. To connect this theoretical structure to the numerical analysis, we solve




Figure 12: Cascade probability comparison between type-uniform control and graph-pressure threshold control under fixed budgets.

a finite-dimensional projection HJB on the state (x_1, x_2, L_1, L_2) , with controls taking values in $[0, \bar{a}]^2$. The discretization uses a semi-implicit step for the second-order diffusion terms in the two x directions to avoid numerical drift caused by the explicit diffusion CFL condition on fine grids. The common-noise cross term $\sigma_0^2 \partial_{x_1 x_2}^2 V$ is retained explicitly. The killing term corresponds to a one-way upward jump to the default cemetery state and the loss coordinates. Numerically, we approximate $V(L + \Delta L) - V(L)$ using a semi-Lagrangian loss-jump scheme, which preserves monotonicity in the loss direction and reduces the ill-conditioned extrapolation caused by central differences or coarse-grid projection. The resulting feedback control is then used in a forward $K = 2$ core-periphery particle system.

The projected first-order control formula is evaluated on this compact finite-difference state grid. On the interior grid points, the semi-implicit diffusion operator is nondegenerate in the alive-state coordinates and the drift, killing, and interpolation coefficients are bounded Lipschitz; the stored value surface is evaluated through the same smooth finite-difference and monotone interpolation operators used in the Hamiltonian step. Thus the numerical minimization is a discrete smooth-verification analogue of the formula in Theorem 6. Projection onto $[0, \bar{a}]^2$ handles the saturation boundary, and nonsmooth grid points are handled by direct projected Hamiltonian minimization.

The HJB risk-cost comparison places the HJB feedback point, no control, type-uniform control, and graph-pressure threshold control on the same metrics. This finite-dimensional discretization shows that the Hamiltonian structure in Theorem 6 generates computable feedback and can be compared with rule-based controls. In the same forward $K = 2$ system, no control and the representative two-population graph-pressure threshold control both have mean terminal loss about 0.362, 95% loss about 0.375, and cascade probability 1.000. Type-uniform control reduces the 95% loss to 0.211 and cascade probability to 0.455 at cost 0.0143. The finite-dimensional HJB feedback further reduces the 95% loss to 0.202 and cascade probability to 0.095 at cost 0.0162. Thus, in this low-dimensional projection, the HJB feedback achieves lower tail loss and lower cascade probability at a comparable intervention scale.

Figure 13 reports the multi-grid diagnostics under the semi-Lagrangian loss-jump scheme. The four resolutions use the same calibration and terminal penalty, and each feedback is evaluated in the

same $K = 2$ forward system. As the grid is refined from $N_x = 20$ to $N_x = 160$, the initial value V_0 decreases from 0.1713 to 0.0459, the initial control decreases from (1.625, 1.750) to (0.625, 0.625), and the average control cost decreases from 0.0680 to 0.0087. Forward tail loss and cascade probability still change sharply, with cascade probability rising from 0 on the coarse grid to 0.790 and 1.000. These grid diagnostics track the finite-dimensional feedback approximation, while the forward validation metrics below provide the primary policy-performance test.




Figure 13: $K = 2$ HJB multi-grid and Richardson-type diagnostics.

The Richardson-type panel in Figure 13 gives a power-law diagnostic for the four available grids. The estimated exponent is $p \simeq 0.476$, the fitted R^2 is close to 1, and the extrapolated intercept is negative. This value identifies the dominant discretization effect: when $N_L = N_x/2$, the semi-Lagrangian term $V(L + \Delta L) - V(L)$ still smears the jump discontinuity created by killing and loss advancement. We use the panel as a discretization-sensitivity diagnostic and evaluate policy performance through direct forward simulation.

To separate diffusion discretization in the x directions from loss-coordinate projection error, we fix $N_x = 320$ and compare $N_L = 20, 40, 60, 80$. Figure 14 shows that V_0 decreases from 0.1158 to 0.0460, the control cost decreases from 0.0400 to 0.0099, and the forward cascade probability rises from 0 to 0.610 and 1.000. Since the second-order HJB terms act only in the x directions, this pattern locates the dominant sensitivity in the loss-coordinate projection and the killing-jump interpolation.

Fixed-loss slice diagnostic The fixed-loss slice diagnostic fixes $L_1 = L_2 = 0$ and refines only the (x_1, x_2) plane. The HJB initial value is 0.3134 at $N_x = 320$ and 0.3137 at $N_x = 640$, with average slice controls about 1.641 on both grids. Figure 15 reports the corresponding control slices. This confirms that the dominant four-dimensional feedback sensitivity comes from loss-coordinate projection and killing-jump interpolation; diffusion discretization in the alive coordinates is comparatively stable on this slice.

Large-scale forward policy validation We evaluate the stored HJB feedback from the $N_x = 160, N_L = 80$ grid in a larger forward particle system with $N = 100,000$ and 200 replications. It yields mean terminal loss 0.1569, 95% loss 0.1638, cascade probability 0.000, and mean control cost 0.0087. This is the primary policy-performance metric: although V_0 and the control field remain




Figure 14: Asymmetric loss-coordinate refinement diagnostic for the $K = 2$ HJB.




Figure 15: (x_1, x_2) heat maps of $K = 2$ HJB feedback control on the fixed-loss slice.

sensitive to the loss coordinate, the stored feedback achieves substantial tail-risk reduction in direct forward evaluation.

Figure 16 shows control slices at a fixed loss state. Control intensity is more concentrated in high-pressure regions, and core-type control increases earlier as system pressure rises. This structure is consistent with the sign interpretation of the projected fixed-point marginal control condition stated in the appendix HJB characterization: when the local pressure of a type has a larger marginal contribution to future loss, the optimal feedback tilts intervention resources toward that type and the corresponding state region.




Figure 16: (x_1, x_2) slices of $K = 2$ HJB feedback control at a fixed loss state.

Figure 17 further gives time slices of the HJB feedback. Early control is closer to a preventive allocation. As the terminal time approaches, feedback becomes increasingly concentrated near the boundary and in regions where the loss state has already deteriorated. This dynamic form is consistent with the marginal benefit term in the Hamiltonian: control has higher value during periods in which future losses can still be changed, and becomes more state-selective when risk has concentrated near the critical region.




Figure 17: Time slices of $K = 2$ HJB feedback control at $t = 0, 0.25, 0.50, 0.75$.

Graph-pressure threshold control in the two-population projection requires a separate explanation. In the representative-state HJB comparison, the graph-pressure variable is generated by type-average new losses, so local pressure heterogeneity within the same type is projected out. The threshold

rule only observes two representative pressure coordinates and cannot identify differences between high-pressure and low-pressure nodes in the sparse graph. The information value conclusion in Theorem 7 requires node-level local pressure information to be strictly finer than type-average information. The representative $K = 2$ projection therefore illustrates the mechanism cleanly: the value of local pressure information comes from node-level information refinement inside a type, not from the label assigned to a type-average pressure coordinate.

To test this interpretation, we recalibrate graph-pressure threshold control in a large-scale $K = 2$ forward system. This experiment restores local pressure heterogeneity in the core-periphery system, so the threshold parameters can form nontrivial risk-cost points. Together with the representative projection above, it shows that the local pressure rule becomes effective precisely when the information domain contains node-level pressure differences.

Threshold recalibration in the $K = 2$ system After large-scale recalibration, graph-pressure threshold control at threshold 0.0005 and upper scale 1.5 attains 95% loss 0.2025, cascade probability 0.205, and cost 0.0300. The HJB feedback attains 95% loss 0.2034, cascade probability 0.120, and cost 0.0166, indicating that the HJB feedback reaches comparable tail loss with lower cost and lower cascade probability.

The explicit core-periphery experiment further fixes $N_1 = 500$, $N_2 = 9500$, with the core type having higher spillover exposure and the periphery type dominating in population size. Figure 18 gives the two-population loss paths, and the control comparison under the same specification is summarized in text. This experiment connects the two-population structure in Theorem 6, the information value mechanism in Theorem 7, and the sparse graph control frontier in the same low-dimensional example.




Figure 18: Population loss paths in the $K = 2$ core-periphery system.

In this explicit core-periphery case, no control and the graph-pressure threshold rule both have mean loss 0.361, 95% loss 0.377, cascade probability 1.000, and zero cost because the two-type projection does not expose within-type pressure differences. Type-uniform control reduces mean loss to 0.200, 95% loss to 0.210, and cascade probability to 0.465 at cost 0.014, while HJB feedback reaches mean loss 0.194, 95% loss 0.204, and cascade probability 0.130 at cost 0.016.

13 Robustness: control parameters and network heterogeneity

To rule out the possibility that graph-pressure threshold control is only the result of a single threshold choice, we vary the graph-pressure threshold and the control upper bound in three representative near-critical regions. The threshold grid covers seven values from 0.0075 to 0.0300, and the control upper-bound ratios are 0.70, 1.00, 1.30. Each parameter point uses 500 Monte Carlo replications. Figure 19 shows that graph-pressure threshold control forms a stable risk–cost frontier in all three network classes: low-intensity control usually has the highest cascade-reduction efficiency per unit cost, while high-intensity control further lowers residual cascade probability.




Figure 19: Risk–cost frontiers under the parameter grid for graph-pressure threshold control.

Peak graph-pressure efficiency across network families The parameter points with the highest cascade reduction per unit cost show that graph-pressure thresholds reduce cascade risk at low cost across the three benchmark networks and the two additional degree-distribution networks. The efficiency ranking remains largest in locally heterogeneous networks: heavy-tailed, core–periphery, and stochastic-block networks attain high-efficiency points, Barabási–Albert networks are intermediate, and Watts–Strogatz networks are lowest. This ordering is consistent with the mechanism that local pressure information becomes more valuable when exposure neighborhoods are uneven.

The highest-efficiency and lowest-residual-cascade parameter points are separated in a stable way. In the heavy-tailed network, the highest-efficiency point uses threshold 0.0300 and upper-bound ratio 0.7000, with cascade probability 0.278 and cost 0.0117; the lowest-cascade point uses threshold 0.0075 and ratio 1.3000, with cascade probability 0.054 and cost 0.0445. The corresponding pairs are (0.256, 0.0123) versus (0.046, 0.0471) for the core–periphery network, and (0.166, 0.0120) versus (0.016, 0.0426) for the stochastic-block network. High-efficiency points therefore use lower control upper bounds, retain some residual cascades, and have very low costs, while the lowest-cascade points require higher upper bounds and higher costs. Parameter robustness analysis reveals the continuous shape of the risk–cost frontier more clearly than a single representative point.

We also conduct network heterogeneity comparisons under fixed spectral radius and fixed average weighted in-exposure to separate total exposure intensity from local exposure concentration. These comparisons standardize the aggregate matrix spectral radius or the average weighted in-exposure of different networks to the same level and then compare the graph system with the matrix system again. The results support our structural interpretation: after controlling for aggregate intensity, heterogeneous networks still display a stronger tendency toward tail gaps. Matrix approximation gaps therefore arise not only from amplification of total edge weight, but also from the concentration pattern of edge weights at the node level.

The time-window robustness comparison extends the same sparse graph transition-region experiment to $T = 2.0$. The longer time window moves all three network classes into the cascade region earlier: the core-periphery and heavy-tailed networks both reach a 10% cascade probability at exposure multiplier 8, with maximum tail gaps 0.722 and 0.658, respectively, while the stochastic-block network enters the cascade region at multiplier 14 with maximum tail gap 0.151. The time window mainly shifts the cascade onset, while local exposure concentration and matrix approximation tail gaps still determine the relative ordering across networks.




Figure 20: Sparse graph robustness under longer horizons and time-varying networks.

Figure 21 reports sensitivity to killing-mechanism parameters. At exposure multiplier 12, all three network classes are in high-pressure regions, and cascade probabilities equal one throughout the scanned grid. The figure therefore omits the saturated cascade-probability panel and reports terminal mean loss and 95% tail loss, where parameter variation remains visible. Increasing the killing-function slope γ or the baseline kill intensity λ_{base} raises tail loss levels. The stochastic block network retains the widest region of continuous variation, while the core-periphery and heavy-tailed networks enter near-saturation states more quickly. The parameter effect of the soft killing mechanism therefore depends on whether the network has already crossed the cascade threshold.

The scanned values confirm the visual pattern. At $\gamma = 5$, cascade probabilities remain close to zero for $\lambda_{\text{base}} \leq 0.05$ and jump to 1.000 at 0.10, with 95% loss rising from 0.101 to 0.259. At larger slopes, saturation arrives earlier: for $\gamma = 10$, cascade probability is already 1.000 once $\lambda_{\text{base}} \geq 0.03$, and for $\gamma \geq 15$ all reported baseline intensities generate cascade probability 1.000.

Figure 20 compares static and time-varying networks. Under the same network family and the same exposure parameters, the time-varying network resamples the adjacency structure at fixed time intervals. It preserves the overall network-family characteristics but does not fix a single edge set throughout the stress period. This comparison tests whether our conclusions rely on the




Figure 21: Effects of the killing-function slope γ and the baseline kill intensity λ_{base} on terminal mean loss and 95% tail loss. Cascade probability is saturated on this grid and is summarized in the text.

static-network assumption. The results show that time-varying resampling changes the arrival paths of local pressure and the magnitude of tail loss, but the relative ordering across the three network classes and the value of local pressure information remain stable. Matrix approximation gaps therefore mainly come from exposure heterogeneity and local pressure information, with the network-ordering result stable across static and time-varying edge realizations.

Quantitatively, the static and time-varying comparisons remain close. In the heavy-tailed network, no-control cascade probabilities are 0.715 and 0.710 under static and time-varying edge sets, while graph-pressure control gives 0.145 and 0.170. In the core-periphery and stochastic-block networks, graph-pressure control keeps cascade probability at zero in both regimes, with 95% losses changing only modestly. The time-varying experiment therefore perturbs path details without changing the network ordering.

Figure 22 further adds Barabási–Albert scale-free networks and Watts–Strogatz small-world networks. The former emphasizes the upper tail of the source-node degree distribution, while the latter emphasizes local clustering and short paths. These two additional network classes do not change the main mechanism. In the cascade onset region, larger structural concentration is accompanied by a larger graph-matrix tail gap. In extremely high-exposure regions, some matrix systems also enter high-loss states, and the gap contracts because the two systems saturate at the same time. This result shows that row dispersion and edge concentration apply not only to the three benchmark networks used in this paper, but also to the onset conditions for matrix approximation distortion under more general degree distributions.




Figure 22: Robustness under additional network families and larger sparse systems.

At multiplier 30, the additional network families remain heterogeneous after matching the experimental scale. Barabási–Albert networks have tail gap 0.7697 and cascade gap 1.000, while Watts–Strogatz networks have tail gap 0.8053 and cascade gap 0.280. For comparison, the stochastic-block network has tail gap 0.5289 and cascade gap 1.000, while the core-periphery and heavy-tailed networks have tail gaps 0.2549 and 0.0502 at this high multiplier. These values show that degree distribution and clustering both alter matrix-approximation failure modes.

Numerical scale of the information-value constants Several constants in the information-value lower bound of Theorem 7 are informative as scale diagnostics. Here q_k is estimated from type terminal losses in the uncontrolled forward system, the matrix-approximation concentration score is computed from the adjacency matrix, and m_k is estimated from the marginal elasticity of the kill

intensity with respect to control near the initial state. The estimated lower-bound scales are 0.1830 for the heavy-tailed network, 0.0754 for the core-periphery network, 0.0362 for the Barabási-Albert network, 0.0301 for the stochastic-block network, and 0.0293 for the Watts-Strogatz network. This ordering is used as a structural scale check rather than as an exact prediction formula, and it is consistent with the interpretation that local pressure information has greater value when within-block exposure heterogeneity is larger.

The large-scale sparse experiment extends the sparse graph experiments to a larger particle scale. The three network classes preserve the same ordering at the larger scale: heavy-tailed and core-periphery structures show tail gaps earlier, while the gap in the stochastic block network appears later. This result shows that the matrix approximation boundary and the value of graph-pressure threshold control are not small-simulation artifacts and continue to appear in larger sparse systems.

At $N = 50,000$ and multiplier 30, all three benchmark networks reach cascade probability 1.000. The 95% losses are 0.9842 for the core-periphery network, 0.9939 for the heavy-tailed network, and 0.5486 for the stochastic-block network, with corresponding tail gaps 0.4331, 0.0720, and 0.4734. The large-scale check therefore preserves the qualitative boundary: the graph system remains substantially more fragile than its type-aggregated matrix approximation in heterogeneous sparse regimes.

The multi-group extension in Table 2 and Figure 23 extends the finite-type matrix system to $K = 3$ and $K = 4$. The $K = 3$ specification merges small banks and regional banks into the periphery type, while the $K = 4$ specification keeps the original four-type structure. In both dimensions, the matrix spectral radius and scaling parameter organize the increase in terminal mean loss and tail loss. On this grid, the $K = 3$ system remains below the cascade threshold, while the $K = 4$ system starts to produce cascades after scaling 48 and reaches a cascade probability of 0.460 at scaling 54. This shows that our matrix diagnostics are not by-products of the $K = 2$ optimal control example, but stable structures in multi-population finite-type contagion systems.




Figure 23: Tail loss and cascade probability in $K = 3$ and $K = 4$ finite-type matrix systems.

14 External-data calibrated network

We combine the preceding synthetic network mechanism with public regulatory data to construct a semi-realistic calibrated core-periphery network. The calibration inputs come from the EBA 2024 EU-wide Transparency Exercise and the official disclosure packages of the EBA Pillar 3 Data Hub / European Data Access Portal [60, 61]. The transparency data provide bank-level

Table 2: Finite-type matrix experiments for $K = 3$ and $K = 4$.

K	Matrix scale	$\rho(\Gamma)$	95% loss	Cascade
3	36.0	20.385	0.0837	0.000
3	42.0	23.783	0.0915	0.000
3	48.0	27.180	0.1066	0.000
3	54.0	30.578	0.1268	0.000
4	36.0	23.206	0.1002	0.000
4	42.0	27.073	0.1249	0.000
4	48.0	30.941	0.1857	0.035
4	54.0	34.809	0.8388	0.460

size, risk-exposure amount, and jurisdictional information. The Pillar 3/EDAP disclosure packages provide institution-level regulatory disclosure modules and reporting-file structures. Together, they constrain the size concentration, core-node share, country-level credit-exposure proxies, and disclosure coverage structure in the synthetic network. The jurisdictional map uses the Natural Earth country-boundary base map [62], which is used only to visualize country outlines and does not provide financial exposure information.

Public EBA and Pillar 3 data do not identify bank-to-bank bilateral exposures. We therefore construct a calibrated synthetic core-periphery network that is compatible with public regulatory statistics. Specifically, nodes are sampled according to EBA asset-size proxies, core nodes are determined by the upper tail of the size distribution, row exposure budgets are adjusted by institution size and country-level credit-exposure proxies, and a sparse core-periphery adjacency matrix is then generated with the aggregate matrix spectral radius rescaled to $\rho(\hat{\Gamma}) \simeq 5$. We use this network to test whether our mechanisms for the matrix approximation boundary and graph-pressure threshold control continue to appear in a calibrated environment constrained by external data.

The inputs, outputs, and parameter choices in this section are as follows. The input is the bank-level record

$$\mathcal{D}_{\text{EBA}} = \{(S_b, R_b, q_b, m_b) : b = 1, \dots, B\},$$

where S_b is an asset-size proxy, R_b is a risk-exposure-amount proxy, q_b is the jurisdiction, and m_b is the set of available regulatory disclosure modules. Country-level credit-exposure proxies are denoted by $\{E_q\}$. The output is a directed weighted calibrated synthetic network

$$\mathcal{G}_{\text{cal}} = (A^{\text{cal}}, g, \omega, q, \hat{\Gamma}^{\text{cal}}),$$

where A^{cal} is the edge-weight matrix, g is the core-periphery type label, ω is the node size weight, q is the node jurisdiction, and $\hat{\Gamma}^{\text{cal}}$ is the aggregate matrix obtained from (11). The forward simulation outputs include L_T^{graph} , $Q_{0.95}(L_T^{\text{graph}})$, $\mathbb{P}(C^{\text{graph}})$, $\mathcal{K}_T^{\text{graph}}$, and the corresponding matrix approximation metrics.

The parameter choices are designed for comparability across the paper. We fix the number of nodes at $N = 10000$, matching the order of magnitude in the sparse graph and control experiments above so that Monte Carlo metrics are directly comparable. Core nodes are determined by the upper tail of the size distribution, so the core share comes from public size constraints. The spectral radius target is set to $\rho(\hat{\Gamma}^{\text{cal}}) \simeq 5$, placing the synthetic network in a near-critical but not fully saturated contagion region and making the differences arise mainly from node-level heterogeneity. The multi-instance experiment fixes these public-data constraints and resamples the unobserved bilateral edge sets, thereby characterizing the network uncertainty left open by public regulatory statistics.

Figure 24 describes the jurisdictional and size structure of the calibrated sample. The European jurisdiction map is colored by bank-level asset-size proxies aggregated at the country level, and the bar chart on the right reports the allocation of synthetic nodes across major jurisdictions. France, Germany, Spain, Italy, and the Netherlands contribute the main size weights, consistent with size concentration in the sample of large European banks. The asset-concentration calculation below records the corresponding core–periphery assignment: core nodes come from the upper tail of the size distribution, so their node share is much lower than their asset-size share. This structure is the main external-data constraint behind core–periphery contagion paths in the calibrated network below.




Figure 24: Jurisdictional distribution and asset-size proxies in the EBA and Pillar 3 calibrated sample. Map labels mark the major reporting jurisdictions; country boundaries are used only for geographic visualization.

Asset concentration in the calibrated core–periphery partition The calibrated sample contains 112 EBA reporting banks and 27 bank jurisdictions, and the synthetic network contains 10000 nodes and about 711000 directed weighted edges. Core nodes represent 8.04% of institutions but 47.78% of total asset-size proxy mass; periphery nodes represent 91.96% of institutions and 52.22% of the proxy mass. The core–periphery split therefore comes from the upper tail of the size distribution rather than from arbitrary assignment, and the row-dispersion and edge-concentration diagnostics show that node-level exposure concentration remains relevant even after total contagion strength is controlled through the spectral radius.

Figure 25 reports forward risk results on this calibrated network. Under the near-critical stress specification, the uncontrolled graph system has terminal cascade probability 0.486 and mean terminal loss 0.2000. Graph-pressure threshold control lowers the cascade probability to 0.174 at control cost 0.0227. Type-uniform control further eliminates the observed cascade events, but its cost rises to 0.2195. This ordering is consistent with the fixed-budget and sparse graph control results above: local pressure information has substantial marginal value in the low-cost region, while uniform control is a high-cost strong-suppression benchmark.

Figure 26 shows the average cumulative loss paths under the three control rules. The no-control path crosses the neighborhood of the cascade threshold in the middle-to-late part of the horizon. Graph-pressure threshold control reduces the path slope and suppresses tail paths. Uniform control produces stronger overall suppression at higher cost. This semi-realistic calibration experiment




Figure 25: Risk and control results in the EBA and Pillar 3 calibrated synthetic core-periphery network.

shows that the theoretical diagnostics and control mechanisms can be combined with network calibration under public regulatory data constraints.




Figure 26: Average loss paths in the EBA and Pillar 3 calibrated synthetic core-periphery network.

We independently generate 20 calibrated synthetic core-periphery networks under the same EBA and Pillar 3 constraints, and perform 500 forward risk comparisons on each network. Figure 27 reports the cross-instance distribution. This experiment fixes the size distribution, jurisdictional structure, core-node share, and spectral-radius target given by public data, and changes the unobserved bilateral edge sets in the synthetic network generator. The cross-instance variation characterizes the bilateral network uncertainty left open by public regulatory statistics. The cross-instance averages show that the no-control cascade probability is 0.4769, graph-pressure threshold control lowers it to 0.1698, and uniform control gives 0. The corresponding average control costs are 0, 0.0227, and

0.2195, respectively. The low-cost risk reduction from graph-pressure threshold control persists in the cross-instance distribution, while uniform control remains a higher-cost strong-suppression benchmark.




Figure 27: Multi-instance calibrated synthetic networks under EBA and Pillar 3 constraints.

15 Discussion and conclusion

The results identify a clear effective range for finite-type matrix approximation. In low-pressure or regular-mixing regions, the aggregate contagion matrix gives a useful stress-testing object because it preserves the main type-level exposure channels and supports interpretable spectral and row-exposure diagnostics. As local exposure concentration rises, the same matrix can preserve aggregate exposure while missing tail paths through hubs, core nodes, or high-pressure neighborhoods. The diagnostic terms in Theorem 3, the tail-gap construction in Theorem 4, and the spectral threshold in Theorem 5 describe this boundary from complementary angles.

The control experiments translate the approximation boundary into regulatory design. Type-uniform control is a strong benchmark when the intervention budget is large and broad suppression is feasible. Graph-pressure threshold control uses node-level local pressure and therefore performs best when the budget is limited and the cascade mechanism is concentrated in a small set of high-pressure neighborhoods. The finite-dimensional $K = 2$ HJB loop provides a marginal-value benchmark: it converts the Hamiltonian structure into a feedback rule that reduces tail loss and cascade probability at comparable cost. The forward policy evaluation applies the stored feedback in a large particle system and serves as the primary validation metric for the computed control.

The HJB diagnostics also clarify the meaning of the Richardson-style extrapolation. A negative extrapolated value such as $V_\infty = -0.029$ identifies a finite-dimensional projection effect: the loss-coordinate grid smears the jump discontinuity created when killing transfers alive mass into the cemetery state and advances the loss coordinates near the contagion threshold. The representative-coordinate projection also suppresses within-type distributional heterogeneity that the full measure-valued HJB would retain. The semi-Lagrangian loss-jump scheme reduces the instability of central differences, and the remaining loss-grid sensitivity is reported through the grid diagnostics. The value-function grid tests discretization sensitivity, while the forward controlled particle system tests realized risk reduction by the computed feedback.

The external-data layer gives calibrated synthetic evidence under public regulatory constraints. EBA Transparency Exercise data and Pillar 3/EDAP disclosure packages constrain size, jurisdictional distribution, concentration, and disclosure coverage, while the confidential bilateral exposure matrix remains unobserved. The calibrated core–periphery networks use these public constraints to test robustness across admissible synthetic edge realizations. The multi-instance results show that the matrix-approximation gap and the value of local pressure information are stable across network draws satisfying the same external constraints.

Open problems Four directions follow naturally from the present framework: sparse-graph McKean–Vlasov limits with killing jumps, common noise, and local weak convergence; scalable feedback classes based on neighborhood distributions or graph neural networks; comparison theory for degenerate, hard-boundary, or unbounded-intensity versions beyond the bounded killed HJB treated here; and richer calibration pipelines using supervisory bilateral data, when available, to separate model error from disclosure-driven network uncertainty. The central conclusion is that finite-type matrices are informative stress-testing tools in regular regimes, while heterogeneous sparse networks require local diagnostics for tail-risk measurement and intervention.

Code and data availability

Replication code is available at https://github.com/Hopper-221/systemic_mkv. The repository contains the forward simulation, finite-dimensional HJB feedback, calibrated synthetic-network construction, and figure/table generation scripts used for the numerical experiments. Public regulatory inputs are obtained from the EBA Transparency Exercise and Pillar 3 Data Hub and are used to construct calibrated synthetic core–periphery networks under public disclosure constraints.

SM1 Numerical discretization and parameter calibration

This appendix records two key discretization choices used in the numerical experiments in the main text. First, the two-population HJB in Theorem 6 is projected onto the representative state (x_1, x_2, L_1, L_2) and solved by backward induction from T to 0. The first-order terms in the x directions use central differences. The two pure second-order diffusion terms are handled by semi-implicit tridiagonal steps, and the common-noise cross second-order term $\sigma_0^2 \partial_{x_1 x_2}^2 V$ is retained explicitly. The control is searched on a discrete grid over the compact set $[0, \bar{a}]^2$. The killing term enters the loss coordinates through a one-way semi-Lagrangian jump, using monotone piecewise-linear interpolation in the loss coordinates to approximate $V(x_1, x_2, L_1 + \Delta L_1, L_2) - V(x_1, x_2, L_1, L_2)$ and the corresponding term for the second type. This preserves the direction in which losses can only increase and reduces ill-conditioned extrapolation caused by central differences or coarse-grid projection. The explicit treatment of the cross term is used only in the calibrated range of common-noise intensities considered in the experiments; the time step and grid are checked through the multi-grid diagnostics, which serve as an ex post CFL-type stability check for this finite-dimensional projection. The multi-grid diagnostic tables in the main text repeat this discretization at $N_x = 20, 40, 80, 160$ and $N_L = N_x/2$, and validate the feedback control in the same forward system by its cascade probability. This diagnostic identifies the sensitivity of finite-dimensional feedback to grid resolution. The rigorous value-function characterization is still given by the measure-valued HJB in Theorem 6 and Appendix SM4.

Second, the $K = 2$ numerical experiments must lie in a nondegenerate near-critical region: if pressure is too low, the marginal control benefit in the Hamiltonian is zero; if pressure is too high, all rules enter the saturated-control region. We scan the kill-intensity multiplier, the terminal-loss weight multiplier, and the running-cost multiplier on a small grid, and choose a parameter region that satisfies three conditions at once: the uncontrolled system has significant cascade risk, HJB feedback gives nonzero control, and the HJB point and rule-based controls have comparable cost magnitudes. Figure 28 and Table 3 give the numerical basis for this calibration. The final calibration used in the main text is kill-intensity multiplier 25, terminal-loss weight multiplier 6, and running-cost multiplier 0.05.

Computational environment and runtime

The numerical experiments were carried out on a Linux workstation equipped with two AMD EPYC 7542 processors, 64 physical CPU cores, 128 hardware threads, 128 GB of system memory, and one NVIDIA GeForce RTX 4090 GPU with 24 GB of memory. The software environment used Python 3.12.7 together with NumPy 1.26.4, SciPy 1.13.1, pandas 2.2.2, Matplotlib 3.9.2, and a development build of PyTorch 2.8.0 with CUDA 12.8 support. The finite-difference HJB computations and sparse-network Monte Carlo experiments were implemented as CPU-based NumPy/SciPy workloads. Although the workstation also contained an NVIDIA RTX 4090 GPU, the reported results do not rely on GPU execution.

The final numerical campaign required approximately 145 aggregate wall-clock job-hours, covering the sparse-control robustness recomputation, the non-HJB numerical experiments, the monotone-jump and semi-Lagrangian HJB computations, and the additional Watts–Strogatz and Barabási–Albert robustness checks. Random seeds were fixed before each experiment family was run. The principal sparse-graph, matrix-approximation, Theorem 3 diagnostic, and two-type HJB experiments use base seed 2026. The later robustness checks, EBA/Pillar 3 calibration experiments, multi-instance experiments, and auxiliary network-family experiments use fixed seeds recorded with the corresponding experimental outputs. Across all experiment families, seeds were specified ex ante

and were not selected based on numerical outcomes.

Numerical implementation summary for the main experiments

The following pseudocode summarizes the numerical implementation of the definitions and experiments in the main text.

Algorithm A.1 (paired forward simulation of the sparse graph system and matrix approximation)

Input. Node types $g(i)$, adjacency matrix A , aggregate matrix $\hat{\Gamma}$, initial states $X_{i,0}$, kill intensity $\lambda_k(x, a)$, control rule $a_{i,t}$, and time grid $0 = t_0 < \dots < t_M = T$.

Intensity convention. In the theoretical experiments, λ_k denotes the bounded regularized intensity specified in the convention before Theorem 1. On compact HJB grids this coincides with the exponential intensity whenever the cap lies above the grid maximum.

Steps.

1. For each Monte Carlo replication, initialize the graph system and matrix system with the same initial states, idiosyncratic noises, common noise, and killing-sampling random variables.
2. At each time step, first update drift, mean reversion, diffusion, and common noise to obtain the pre-contagion state X_{i,t_m}^{pre} .
3. Sample new defaults $\Delta D_{i,t_m}$ according to the kill probability

$$p_{i,m} = 1 - \exp\{-\lambda_{g(i)}(X_{i,t_m}, a_{i,t_m})\Delta t\}$$

4. In the graph system, shock surviving nodes by $P_{i,t_m}^{\text{graph}} = \sum_j A_{ij} \Delta D_{j,t_m}$. In the matrix system, transmit type-average shocks by $\sum_\ell \hat{\Gamma}_{g(i)\ell} \Delta L_{t_m}^{\ell, \text{matrix}}$.
5. Record L_T , $Q_{0.95}(L_T)$, the cascade event C , the control cost $\mathcal{K}_T$, and the graph-matrix gap.

Output. Mean loss, tail loss, cascade probability, cost, risk reduction per unit cost, and the corresponding confidence intervals.

Algorithm A.2 (Theorem 3 diagnostic and fixed spectral radius comparison)

Input. Adjacency matrix A , node types g , and target spectral radius ρ_0 .

Steps.

1. Compute $\hat{\Gamma}$ by averaging total weighted exposure from each source type to each target type. In the fixed spectral radius experiment, multiply all edge weights by $\rho_0/\rho(\hat{\Gamma})$.
2. For each type k , compute the row-dispersion term $R_k(A)$ and the square-edge-spread term $S_k(A)$ in Theorem SM3.4.
3. Form the diagnostic quantity $D_{T3}(A) = \sum_k \pi_k(R_k(A) + S_k(A))$.
4. Run Algorithm A.1 under the same default mechanism and compare $D_{T3}(A)$ with $|\mathbb{E}L_T^{\text{graph}} - \mathbb{E}L_T^{\text{matrix}}|$ and the tail gap.

Output. Theoretical diagnostic, realized gap, network ordering, and heterogeneity comparison after fixing the spectral radius.

Algorithm A.3 ($K = 2$ finite-dimensional HJB feedback and forward validation)

Input. $K = 2$ parameters, terminal loss penalty Φ , control costs c_k , grid (x_1, x_2, L_1, L_2) , time step, and control grid.

Steps.

1. Set $V(T, x_1, x_2, L_1, L_2) = \Phi(L_1, L_2)$.
2. Apply backward induction from T to 0: update diffusion terms semi-implicitly in the x directions, retain the common-noise cross term explicitly, and compute $V(L + \Delta L) - V(L)$ for the killing term using one-way semi-Lagrangian interpolation.
3. At each grid point, search over discrete controls in $[0, \bar{a}]^2$, minimize the Hamiltonian, and store the feedbacks a_1^*, a_2^* for the required time slices.
4. Interpolate the feedback into the forward $K = 2$ particle system, and repeat Monte Carlo validation for each grid resolution.
5. Compute V_0 , initial control, adjacent-grid control-field changes, terminal tail loss, cascade probability, and control cost.

Output. HJB risk-cost point, multi-grid diagnostic, asymmetric refinement diagnostic, time-slice heat maps, and forward policy evaluation.

Algorithm A.4 (EBA and Pillar 3 calibrated synthetic core-periphery network)

Input. Bank size proxy S_b , risk-exposure-amount proxy R_b , jurisdiction q_b , disclosure-module set m_b , country-level credit-exposure proxy E_q , and target spectral radius ρ_0 .

Steps.

1. Construct the node size distribution from S_b , and determine the core-node set from the upper tail of the size distribution.
2. Allocate node row exposure budgets using S_b , R_b , and E_q , while preserving jurisdictional and disclosure-coverage structures.
3. Generate a sparse directed core-periphery edge set, and rescale edge weights so that $\rho(\hat{\Gamma}) \simeq \rho_0$.
4. Fix public-data constraints and independently generate multiple synthetic networks to measure network uncertainty from unobserved bilateral edge sets.
5. Run Algorithm A.1 on each network, comparing no control, graph-pressure threshold control, and uniform control.

Output. Calibrated network diagnostics, jurisdictional and size distribution figures, and single-instance and multi-instance forward risk results.

Table 3: Top six candidate parameters in the $K = 2$ HJB numerical calibration.

λ multiplier	Terminal-weight multiplier	Cost multiplier	Nonzero-control share	HJB 95% loss	HJB cost
25	6	0.050	0.800	0.102	0.073
25	4	0.050	0.800	0.122	0.050
25	6	0.075	0.800	0.122	0.075
25	4	0.075	0.800	0.135	0.060
25	6	0.100	0.800	0.135	0.079
20	6	0.050	0.800	0.103	0.057




Figure 28: Candidate points in the $K = 2$ HJB numerical calibration grid. Point color represents the kill-intensity multiplier, and point size represents the nonzero-control share.

SM2 Proofs for the mean-field limit, quantitative rate, and propagation of chaos

Lemma SM2.1 (Conditional empirical LLN for killed diffusion paths). *Fix a type k . Given the common-noise path W^0 and a common-noise adapted type-level control, the limiting killed particles*

$$(\bar{X}_i^k, \bar{\xi}_i^k, \bar{\tau}_i^k)_{i \geq 1}$$

are conditionally independent and identically distributed. Assume that the conditional law of $\bar{\tau}_i^k$ has no fixed-time atoms on $[0, T]$, which is satisfied for the bounded Cox killing construction used here. Then

$$\mathbb{E} \sup_{t \leq T} d_{\text{BL}}(\tilde{\mu}_t^{k,N}, \mu_t^k) \longrightarrow 0,$$

and

$$\mathbb{E} \sup_{t \leq T} |\tilde{L}_t^{k,N} - L_t^k| \longrightarrow 0.$$

If the conditional empirical process of the killed paths satisfies the standard quadratic bounded-Lipschitz fluctuation bound, then the same argument gives the corresponding N_k^{-1} mean-square estimate used in Theorem SM2.7.

Proof. Condition on $\mathcal{F}_T^{W^0}$. The common-noise adapted control is then a deterministic path, and the remaining randomness comes only from the independent initial states, idiosyncratic Brownian motions, and Poisson thinning measures. Hence the limiting killed paths are i.i.d. on the path space $D([0, T]; \mathbb{E})$, with $\mathbb{E} = \mathbb{R} \cup \{\partial\}$. The bounded-Lipschitz unit ball on this Polish path space has a countable determining subclass. The ordinary Glivenko–Cantelli theorem for this countable determining class, followed by monotone approximation of the time supremum over a dense grid and the no-fixed-atom property of the killing times, gives the two uniform convergence statements after

removing the conditioning. The quadratic version is the standard empirical-process strengthening of the same conditional LLN and is stated separately in Theorem SM2.7 because it is the finite-sample input needed for the N_k^{-1} rate. $\square$

Lemma SM2.2 (Metric-level quadratic bounded-Lipschitz stability). *Under the synchronous coupling used in Theorem SM2.6, suppose the system has a uniform second-moment bound and the bounded-Lipschitz empirical process of the paired killed particles satisfies the usual separability and quadratic fluctuation estimate on the bounded-Lipschitz unit ball. Then, for the auxiliary empirical surviving measure $\tilde{\mu}_t^{k,N}$,*

$$\begin{aligned} & \mathbb{E} \sup_{r \leq t} d_{\text{BL}}^2(\mu_r^{k,N}, \tilde{\mu}_r^{k,N}) \\ & \leq C \int_0^t \left(E_k(s) + \sum_{\ell=1}^K D_\ell(s) + \sum_{\ell=1}^K V_\ell(s) \right) ds + \frac{C}{N_k}, \end{aligned}$$

where

$$E_k(t) = \mathbb{E}|X_i^{k,N}(t) - \bar{X}_i^k(t)|^2, \quad D_k(t) = \mathbb{E} d_{\text{BL}}^2(\mu_t^{k,N}, \mu_t^k), \quad V_k(t) = \mathbb{E}|L_t^{k,N} - L_t^k|^2.$$

Proof. For each bounded Lipschitz test function, the synchronous Poisson thinning construction bounds the squared difference of the paired empirical integrals by the squared state-coupling error, the squared loss-coupling error, and the martingale fluctuation of order N_k^{-1} . To pass from a single test function to d_{BL} , take a countable dense determining subclass of the bounded-Lipschitz unit ball, use the assumed quadratic empirical-process bound on this class, and then pass to the full supremum by separability. This is precisely the metric-level form of the synchronous stability estimate; it is recorded as a lemma so that Theorem SM2.7 does not rely on an invalid interchange between a pointwise test-function estimate and the bounded-Lipschitz supremum. $\square$

Lemma SM2.3 (Sufficient finite-entropy verification of the quadratic empirical inputs). *Fix a type k and work conditionally on $\mathcal{F}_T^{W^0}$. Suppose, after localization to a compact alive-state interval and adjoining the cemetery point, that the bounded-Lipschitz test class used to metrize the lifted killed paths admits a countable envelope-normalized subclass $\mathcal{F}_k$ with entropy integral*

$$\int_0^1 \sqrt{\log N(\varepsilon, \mathcal{F}_k, L^2(Q))} d\varepsilon < \infty$$

uniformly over all probability laws Q generated by the conditional limiting killed paths in the model class. Assume also the bounded killing intensity, the Lipschitz state dependence of λ_k , and the uniform second-moment bound used in Theorem SM2.7. Then the auxiliary empirical measure satisfies

$$\mathbb{E} \sup_{t \leq T} d_{\text{BL}}^2(\tilde{\mu}_t^{k,N}, \mu_t^k) \leq \frac{C}{N_k}, \quad \mathbb{E} \sup_{t \leq T} |\tilde{L}_t^{k,N} - L_t^k|^2 \leq \frac{C}{N_k},$$

and the metric-level synchronous-thinning estimate in Lemma SM2.2 holds with the same order. The constant depends on the entropy integral, the localization radius, T , L_λ , Λ , and the moment bound, but not on N_k .

Proof. Conditional on $\mathcal{F}_T^{W^0}$, the limiting killed paths are i.i.d. random elements of the localized Polish path space. The assumed uniform entropy integral makes the bounded-Lipschitz determining

class $\mathcal{F}_k$ conditionally Donsker and gives the maximal inequality

$$\mathbb{E} \sup_{f \in \mathcal{F}_k} \left| \frac{1}{N_k} \sum_{i=1}^{N_k} (f(\bar{X}_i^k, \bar{\xi}_i^k) - \mathbb{E}[f(\bar{X}_i^k, \bar{\xi}_i^k) \mid \mathcal{F}_T^{W^0}]) \right|^2 \leq \frac{C}{N_k}.$$

Applying this estimate to the time-indexed alive-measure and loss-coordinate tests gives the two displayed auxiliary empirical bounds. For the paired finite and limiting systems, write the difference of empirical integrals as the sum of an alive-state displacement term, a cemetery-state mismatch term, and a centered compensator martingale. The displacement term is controlled by the squared state-coupling error. The cemetery mismatch is controlled by synchronous Poisson thinning:

$$\mathbb{E} \sup_{r \leq t} |\xi_i^{k,N}(r) - \bar{\xi}_i^k(r)|^2 \leq C \int_0^t \mathbb{E} |X_i^{k,N}(s) - \bar{X}_i^k(s)|^2 ds.$$

The centered compensator martingale has quadratic variation bounded by C/N_k , because the intensity is bounded by Λ . The same entropy maximal inequality transfers these pointwise estimates to the bounded-Lipschitz supremum. This yields Lemma SM2.2 under the stated finite-entropy condition. $\square$

Remark SM2.4 (Concrete localized entropy verification). In the compact alive-state version used for the bounded-killing theory, the finite-entropy condition in Lemma SM2.3 can be checked from standard bracketing entropy bounds. After adjoining the isolated cemetery point, the lifted killed-path space is the union of a compact one-dimensional diffusion path class with one cemetery jump coordinate. The bounded drift, bounded diffusion coefficient, and bounded killing rate give a uniform modulus-in-probability for the alive diffusion paths, and the cemetery coordinate is a monotone indicator with no fixed-time atom. The bounded-Lipschitz unit ball restricted to this localized class has polynomial bracketing entropy under the laws generated by the model; hence the Dudley–bracketing integral is finite by the entropy integral criterion of van der Vaart and Wellner [63, Theorem 2.7.11]. Therefore, on compact state grids or after localization before removing the localization error, the quadratic empirical input in Theorem SM2.7 follows from the stated bounded-Lipschitz and bounded-killing primitives.

Remark SM2.5 (Empirical-process input for the conditional rate). The quadratic empirical inputs in Theorem SM2.7 are verified by Lemma SM2.3 under a localized finite-entropy condition, and are otherwise stated explicitly because the killed-particle state space is path-valued and contains a cemetery state. They are standard once one works conditionally on the common-noise path. Conditional on $\mathcal{F}_T^{W^0}$, the limiting killed paths are i.i.d. random elements of the Polish space $D([0, T]; \mathbb{E})$, with $\mathbb{E} = \mathbb{R} \cup \{\partial\}$. The bounded-Lipschitz unit ball on this space admits a countable determining subclass, and the usual separability and symmetrization arguments for empirical processes give the N_k^{-1} second-moment fluctuation scale for bounded Lipschitz test functions; see, for example, van der Vaart and Wellner [63]. The additional metric-level stability term in Lemma SM2.2 is the synchronous Poisson-thinning analogue of the same argument: the alive-state displacement, cemetery-state mismatch, and compensator martingale terms are controlled before taking the bounded-Lipschitz supremum. Thus Theorem SM2.7 should be read as a conditional quantitative propagation estimate under these standard empirical-process and thinning-stability inputs, not as a pointwise-test-function estimate upgraded without justification.

Theorem SM2.6 (McKean–Vlasov limit for the multi-class bank particle system). *Fix $T > 0$. The system contains K classes of banks. Class k contains N_k particles, and write*

$$N_* = \min_{1 \leq k \leq K} N_k.$$

For $k = 1, \dots, K$ and $i = 1, \dots, N_k$, the finite-particle system satisfies

$$dX_i^{k,N}(t) = b_k(X_i^{k,N}(t), \mu_t^{1,N}, \dots, \mu_t^{K,N}, a_t^k) dt + \sigma_k dW_i^k(t) + \sigma_0 dW^0(t) - \sum_{\ell=1}^K \Gamma_{k\ell} dL_t^{\ell,N}.$$

Here W^0 is the common noise, W_i^k is the idiosyncratic noise, and $\Gamma = (\Gamma_{k\ell})_{1 \leq k, \ell \leq K}$ is the contagion matrix. The surviving empirical measure and loss process of class k are defined by

$$\mu_t^{k,N} = \frac{1}{N_k} \sum_{i=1}^{N_k} \xi_i^{k,N}(t) \delta_{X_i^{k,N}(t)}, \quad L_t^{k,N} = \frac{1}{N_k} \sum_{i=1}^{N_k} (1 - \xi_i^{k,N}(t)),$$

where

$$\xi_i^{k,N}(t) = \mathbf{1}_{\{\tau_i^{k,N} > t\}}.$$

The killing time is given by Poisson thinning. Choose $\Lambda > 0$ such that

$$0 \leq \lambda_k(x, a) \leq \Lambda.$$

Let $\mathcal{N}_i^k(ds, du)$ be an independent Poisson random measure on $[0, \infty) \times [0, \Lambda]$ with intensity $ds du$, and define

$$\tau_i^{k,N} = \inf \left\{ t : \int_0^t \int_0^\Lambda \mathbf{1}_{\{u \leq \lambda_k(X_i^{k,N}(s-), a_s^k)\}} \mathcal{N}_i^k(ds, du) \geq 1 \right\}.$$

Assume that the initial states are independent and identically distributed within each class and have uniformly finite first moments. Assume that the drift function b_k is Lipschitz in the state variable and measure variables: there exists a constant $L_b > 0$ such that

$$\left| b_k(x, \nu^1, \dots, \nu^K, a) - b_k(y, \eta^1, \dots, \eta^K, a) \right| \leq L_b \left(|x - y| + \sum_{\ell=1}^K d_{\text{BL}}(\nu^\ell, \eta^\ell) \right),$$

where d_{BL} denotes the bounded Lipschitz distance on finite measures,

$$d_{\text{BL}}(\nu, \eta) = \sup_{\|f\|_\infty \leq 1, \text{Lip}(f) \leq 1} \left| \int f d\nu - \int f d\eta \right|.$$

Assume also that the kill intensity λ_k is Lipschitz in the state variable, namely

$$|\lambda_k(x, a) - \lambda_k(y, a)| \leq L_\lambda |x - y|,$$

and that the contagion matrix Γ is bounded. The type-level control $a_t = (a_t^1, \dots, a_t^K)$ is progressively measurable with respect to the common-noise filtration $\mathbb{F}^{W^0}$ and takes values in a compact set. This convention is the open-loop mean-field control convention used in the limit theorem; node-level graph-feedback controls are treated later as finite-network rules. Assume that both the finite-particle system and the limiting system below are well posed.

The limiting McKean–Vlasov system is defined as follows. Given the common noise W^0 , the limiting particle $\bar{X}_i^k$ of class k satisfies

$$d\bar{X}_i^k(t) = b_k(\bar{X}_i^k(t), \mu_t^1, \dots, \mu_t^K, a_t^k) dt + \sigma_k dW_i^k(t) + \sigma_0 dW^0(t) - \sum_{\ell=1}^K \Gamma_{k\ell} dL_t^\ell.$$

The limiting surviving measure is the conditional sub-probability measure

$$\mu_t^k = \mathbb{E}[\bar{\xi}_i^k(t) \delta_{\bar{X}_i^k(t)} | \mathcal{F}_t^{W^0}],$$

and the limiting loss process is

$$L_t^k = 1 - \mathbb{E}[\bar{\xi}_i^k(t) | \mathcal{F}_t^{W^0}].$$

Equivalently,

$$L_t^k = \int_0^t \mathbb{E}[\lambda_k(\bar{X}_i^k(s), a_s^k) \bar{\xi}_i^k(s) | \mathcal{F}_s^{W^0}] ds.$$

Then, as $N_* \rightarrow \infty$, for each $k = 1, \dots, K$,

$$\sup_{t \leq T} d_{\text{BL}}(\mu_t^{k,N}, \mu_t^k) \longrightarrow 0 \quad \text{in probability,}$$

and

$$\sup_{t \leq T} |L_t^{k,N} - L_t^k| \longrightarrow 0 \quad \text{in probability.}$$

Thus the empirical surviving measures of the finite-particle system converge weakly to the coupled McKean–Vlasov limit, and the loss processes converge uniformly in probability on $[0, T]$.

Proof. Construct the finite-particle system and the limiting system on the same probability space. For each i and k , let $X_i^{k,N}$ and $\bar{X}_i^k$ use the same initial state, the same idiosyncratic Brownian motion W_i^k , the same common Brownian motion W^0 , and the same Poisson random measure $\mathcal{N}_i^k$. Define the limiting killing time by

$$\bar{\tau}_i^k = \inf \left\{ t : \int_0^t \int_0^\Lambda \mathbf{1}_{\{u \leq \lambda_k(\bar{X}_i^k(s-), a_s^k)\}} \mathcal{N}_i^k(ds, du) \geq 1 \right\},$$

and write

$$\bar{\xi}_i^k(t) = \mathbf{1}_{\{\bar{\tau}_i^k > t\}}.$$

This synchronous Poisson thinning construction ensures that the difference between the killing indicators of the finite particle and the limiting particle comes only from the difference between the two kill-intensity paths

$$\lambda_k(X_i^{k,N}(t), a_t^k), \quad \lambda_k(\bar{X}_i^k(t), a_t^k).$$

Define the coupling error

$$e_i^k(t) = X_i^{k,N}(t) - \bar{X}_i^k(t).$$

Since the finite particle and the limiting particle use the same Brownian motions, the diffusion terms cancel. Hence

$$\begin{aligned} e_i^k(t) = & \int_0^t \left[b_k(X_i^{k,N}(s), \mu_s^{1,N}, \dots, \mu_s^{K,N}, a_s^k) \right. \\ & \left. - b_k(\bar{X}_i^k(s), \mu_s^1, \dots, \mu_s^K, a_s^k) \right] ds - \sum_{\ell=1}^K \Gamma_{k\ell} (L_t^{\ell,N} - L_t^\ell). \end{aligned}$$

By the Lipschitz property of b_k and the boundedness of Γ , there exists a constant $C > 0$, independent of $\mathbf{N} = (N_1, \dots, N_K)$, such that

$$\sup_{r \leq t} |e_i^k(r)| \leq C \int_0^t \left(|e_i^k(s)| + \sum_{\ell=1}^K d_{\text{BL}}(\mu_s^{\ell,N}, \mu_s^\ell) \right) ds + C \sum_{\ell=1}^K \sup_{r \leq t} |L_r^{\ell,N} - L_r^\ell|.$$

Write

$$u^k(t) = \mathbb{E} \sup_{r \leq t} |e_i^k(r)|, \quad m^k(t) = \mathbb{E} \sup_{r \leq t} d_{\text{BL}}(\mu_r^{k,N}, \mu_r^k),$$

and

$$v^k(t) = \mathbb{E} \sup_{r \leq t} |L_r^{k,N} - L_r^k|.$$

Taking expectations gives

$$u^k(t) \leq C \int_0^t \left(u^k(s) + \sum_{\ell=1}^K m^\ell(s) \right) ds + C \sum_{\ell=1}^K v^\ell(t).$$

We next estimate the empirical measure error. Introduce the auxiliary empirical measure generated by the limiting particles,

$$\tilde{\mu}_t^{k,N} = \frac{1}{N_k} \sum_{i=1}^{N_k} \bar{\xi}_i^k(t) \delta_{\bar{X}_i^k(t)}.$$

By the triangle inequality,

$$d_{\text{BL}}(\mu_t^{k,N}, \mu_t^k) \leq d_{\text{BL}}(\mu_t^{k,N}, \tilde{\mu}_t^{k,N}) + d_{\text{BL}}(\tilde{\mu}_t^{k,N}, \mu_t^k).$$

For any test function f with $\|f\|_\infty \leq 1$ and $\text{Lip}(f) \leq 1$,

$$\begin{aligned} \left| \langle f, \mu_t^{k,N} - \tilde{\mu}_t^{k,N} \rangle \right| &\leq \frac{1}{N_k} \sum_{i=1}^{N_k} \left| \xi_i^{k,N}(t) f(X_i^{k,N}(t)) - \bar{\xi}_i^k(t) f(\bar{X}_i^k(t)) \right| \\ &\leq \frac{1}{N_k} \sum_{i=1}^{N_k} |X_i^{k,N}(t) - \bar{X}_i^k(t)| + \frac{1}{N_k} \sum_{i=1}^{N_k} |\xi_i^{k,N}(t) - \bar{\xi}_i^k(t)|. \end{aligned}$$

The second term only uses $\|f\|_\infty \leq 1$. Taking the supremum over test functions, then the supremum over $r \leq t$, and finally expectation, yields

$$\mathbb{E} \sup_{r \leq t} d_{\text{BL}}(\mu_r^{k,N}, \tilde{\mu}_r^{k,N}) \leq u^k(t) + \mathbb{E} \sup_{r \leq t} \frac{1}{N_k} \sum_{i=1}^{N_k} |\xi_i^{k,N}(r) - \bar{\xi}_i^k(r)|.$$

Synchronous Poisson thinning gives

$$\mathbb{E} \sup_{r \leq t} |\xi_i^{k,N}(r) - \bar{\xi}_i^k(r)| \leq \mathbb{E} \int_0^t \left| \lambda_k(X_i^{k,N}(s), a_s^k) - \lambda_k(\bar{X}_i^k(s), a_s^k) \right| ds.$$

By the Lipschitz property of λ_k ,

$$\mathbb{E} \sup_{r \leq t} |\xi_i^{k,N}(r) - \bar{\xi}_i^k(r)| \leq L_\lambda \int_0^t u^k(s) ds.$$

Therefore,

$$\mathbb{E} \sup_{r \leq t} d_{\text{BL}}(\mu_r^{k,N}, \tilde{\mu}_r^{k,N}) \leq u^k(t) + C \int_0^t u^k(s) ds.$$

On the other hand, conditional on the common noise W^0 , the limiting particles

$$\{(\bar{X}_i^k, \bar{\xi}_i^k)\}_{i=1}^{N_k}$$

are conditionally independent and identically distributed. Lemma [SM2.1](#) gives

$$\alpha_{N_k}^k(T) := \mathbb{E} \sup_{t \leq T} d_{\text{BL}}(\tilde{\mu}_t^{k,N}, \mu_t^k) \longrightarrow 0 \quad (N_k \rightarrow \infty).$$

Hence

$$m^k(t) \leq u^k(t) + C \int_0^t u^k(s) ds + \alpha_{N_k}^k(T).$$

We now estimate the loss process. Define the auxiliary empirical loss process of the limiting particles by

$$\tilde{L}_t^{k,N} = \frac{1}{N_k} \sum_{i=1}^{N_k} (1 - \bar{\xi}_i^k(t)).$$

Then

$$|L_t^{k,N} - L_t^k| \leq |L_t^{k,N} - \tilde{L}_t^{k,N}| + |\tilde{L}_t^{k,N} - L_t^k|.$$

The first term is controlled by the synchronous killing coupling:

$$\begin{aligned} \mathbb{E} \sup_{r \leq t} |L_r^{k,N} - \tilde{L}_r^{k,N}| &\leq \mathbb{E} \sup_{r \leq t} \frac{1}{N_k} \sum_{i=1}^{N_k} |\xi_i^{k,N}(r) - \bar{\xi}_i^k(r)| \\ &\leq C \int_0^t u^k(s) ds. \end{aligned}$$

The second term is controlled by Lemma [SM2.1](#). Given the common noise W^0 , the killing times

$$\{\bar{\tau}_i^k\}_{i=1}^{N_k}$$

are conditionally independent and identically distributed with conditional distribution function L_t^k . Hence

$$\beta_{N_k}^k(T) := \mathbb{E} \sup_{t \leq T} |\tilde{L}_t^{k,N} - L_t^k| \longrightarrow 0 \quad (N_k \rightarrow \infty).$$

Thus

$$v^k(t) \leq C \int_0^t u^k(s) ds + \beta_{N_k}^k(T).$$

Let

$$U(t) = \sum_{k=1}^K u^k(t), \quad M(t) = \sum_{k=1}^K m^k(t), \quad V(t) = \sum_{k=1}^K v^k(t).$$

The estimates above give

$$U(t) \leq C \int_0^t (U(s) + M(s)) ds + CV(t),$$

$$M(t) \leq CU(t) + C \int_0^t U(s) ds + A_{\mathbf{N}}(T),$$

and

$$V(t) \leq C \int_0^t U(s) ds + B_{\mathbf{N}}(T),$$

where

$$A_{\mathbf{N}}(T) = \sum_{k=1}^K \alpha_{N_k}^k(T), \quad B_{\mathbf{N}}(T) = \sum_{k=1}^K \beta_{N_k}^k(T).$$

As $N_* \rightarrow \infty$,

$$A_{\mathbf{N}}(T) \rightarrow 0, \quad B_{\mathbf{N}}(T) \rightarrow 0.$$

Substituting the estimates for M and V into the estimate for U yields

$$U(t) \leq C \int_0^t U(s) ds + C A_{\mathbf{N}}(T) + C B_{\mathbf{N}}(T).$$

By Gronwall's inequality,

$$U(T) \leq C_T (A_{\mathbf{N}}(T) + B_{\mathbf{N}}(T)) \rightarrow 0.$$

Substituting this bound back gives

$$M(T) \rightarrow 0, \quad V(T) \rightarrow 0.$$

Therefore, for each $k = 1, \dots, K$,

$$\mathbb{E} \sup_{t \leq T} d_{\text{BL}}(\mu_t^{k,N}, \mu_t^k) \rightarrow 0,$$

and

$$\mathbb{E} \sup_{t \leq T} |L_t^{k,N} - L_t^k| \rightarrow 0.$$

Markov's inequality gives

$$\sup_{t \leq T} d_{\text{BL}}(\mu_t^{k,N}, \mu_t^k) \rightarrow 0 \quad \text{in probability,}$$

and

$$\sup_{t \leq T} |L_t^{k,N} - L_t^k| \rightarrow 0 \quad \text{in probability.}$$

Since d_{BL} metrizes weak convergence of finite measures, the empirical surviving measures of the finite-particle system converge weakly to the McKean–Vlasov limit, and the loss processes converge uniformly in probability on $[0, T]$. The proof is complete. $\square$

Theorem SM2.7 (Conditional quantitative convergence rate under strengthened conditions). *Under the conditions of Theorem SM2.6, assume also that the system has a uniform second-moment bound and that the quadratic versions of the two empirical inputs in Lemmas SM2.1 and SM2.2 hold. Equivalently, the auxiliary empirical surviving measure*

$$\tilde{\mu}_t^{k,N} = \frac{1}{N_k} \sum_{i=1}^{N_k} \bar{\xi}_i^k(t) \delta_{\bar{X}_i^k(t)}$$

satisfies

$$\sup_{t \in [0, T]} \mathbb{E} d_{\text{BL}}^2(\tilde{\mu}_t^{k,N}, \mu_t^k) \leq \frac{C_{\text{emp}}(T)}{N_k}, \quad k = 1, \dots, K,$$

and the paired finite and limiting auxiliary systems satisfy the metric-level stability estimate of Lemma SM2.2. This formulation records the empirical-process step explicitly: the rate is stated in the bounded-Lipschitz metric, and the supremum over bounded-Lipschitz test functions is handled by a separability/quadratic-fluctuation input rather than by a pointwise test-function estimate.

Then there exists a constant

$$C_T = C(T, L_b, L_\lambda, \|\Gamma\|, \Lambda, C_{\text{emp}}(T)),$$

such that, for each $k = 1, \dots, K$,

$$\sup_{t \in [0, T]} \mathbb{E} d_{\text{BL}}^2(\mu_t^{k, N}, \mu_t^k) \leq \frac{C_T}{N_*},$$

and

$$\sup_{t \in [0, T]} \mathbb{E} |L_t^{k, N} - L_t^k|^2 \leq \frac{C_T}{N_*}.$$

Thus the first-order bounded-Lipschitz empirical-measure error and the first-order loss-process error are both $O(N_*^{-1/2})$, uniformly over $t \in [0, T]$. The constant has Gronwall-type growth and can be written as

$$C_T \leq C_0 \exp\{C_1 T(1 + \|\Gamma\|)^2\},$$

where C_0, C_1 do not depend on $\mathbf{N} = (N_1, \dots, N_K)$.

Proof. Use the synchronous coupling from the proof of Theorem SM2.6. For each i, k , the finite particle $X_i^{k, N}$ and the limiting particle $\bar{X}_i^k$ use the same initial state, the same idiosyncratic Brownian motion, the same common Brownian motion, and the same Poisson random measure. Let

$$e_i^k(t) = X_i^{k, N}(t) - \bar{X}_i^k(t).$$

Define

$$E_k(t) := \mathbb{E} |e_i^k(t)|^2, \quad D_k(t) := \mathbb{E} d_{\text{BL}}^2(\mu_t^{k, N}, \mu_t^k), \quad V_k(t) := \mathbb{E} |L_t^{k, N} - L_t^k|^2.$$

Since the Brownian terms are fully synchronized,

$$\begin{aligned} e_i^k(t) &= \int_0^t \left[b_k(X_i^{k, N}(s), \boldsymbol{\mu}_s^N, a_s^k) - b_k(\bar{X}_i^k(s), \boldsymbol{\mu}_s, a_s^k) \right] ds \\ &\quad - \sum_{\ell=1}^K \Gamma_{k\ell} (L_t^{\ell, N} - L_t^\ell), \end{aligned}$$

where $\boldsymbol{\mu}_s^N = (\mu_s^{1, N}, \dots, \mu_s^{K, N})$ and $\boldsymbol{\mu}_s = (\mu_s^1, \dots, \mu_s^K)$. By the Lipschitz property of b_k , the boundedness of Γ , and the definition of d_{BL} ,

$$E_k(t) \leq C \int_0^t \left(E_k(s) + \sum_{\ell=1}^K D_\ell(s) \right) ds + C \sum_{\ell=1}^K V_\ell(t).$$

We next estimate the surviving-measure error. By the triangle inequality,

$$d_{\text{BL}}(\mu_t^{k, N}, \mu_t^k) \leq d_{\text{BL}}(\mu_t^{k, N}, \tilde{\mu}_t^{k, N}) + d_{\text{BL}}(\tilde{\mu}_t^{k, N}, \mu_t^k).$$

The assumed quadratic synchronous-thinning stability gives

$$\mathbb{E} d_{\text{BL}}^2(\mu_t^{k, N}, \tilde{\mu}_t^{k, N}) \leq C \int_0^t \left(E_k(s) + \sum_{\ell=1}^K D_\ell(s) + \sum_{\ell=1}^K V_\ell(s) \right) ds + \frac{C}{N_k}.$$

The auxiliary empirical fluctuation estimate gives

$$\mathbb{E} d_{\text{BL}}^2(\tilde{\mu}_t^{k, N}, \mu_t^k) \leq \frac{C_{\text{emp}}(T)}{N_k}.$$

Therefore,

$$D_k(t) \leq C \int_0^t \left(E_k(s) + \sum_{\ell=1}^K D_\ell(s) + \sum_{\ell=1}^K V_\ell(s) \right) ds + \frac{C}{N_k}.$$

For the loss process, the finite empirical loss has the Doob–Meyer decomposition

$$L_t^{k,N} = M_t^{k,N} + \int_0^t \langle \lambda_k(\cdot, a_s^k), \mu_s^{k,N} \rangle ds,$$

where $M^{k,N}$ is a martingale with

$$\mathbb{E}|M_t^{k,N}|^2 \leq \frac{C\Lambda T}{N_k}.$$

The limiting loss process satisfies

$$L_t^k = \int_0^t \langle \lambda_k(\cdot, a_s^k), \mu_s^k \rangle ds.$$

Since λ_k is bounded and Lipschitz, the pairing with λ_k is controlled by the bounded-Lipschitz distance after rescaling the test function by a deterministic constant. Hence

$$\left| \langle \lambda_k(\cdot, a_s^k), \mu_s^{k,N} \rangle - \langle \lambda_k(\cdot, a_s^k), \mu_s^k \rangle \right| \leq C d_{\text{BL}}(\mu_s^{k,N}, \mu_s^k).$$

By Cauchy's inequality,

$$V_k(t) \leq \frac{C}{N_k} + C \int_0^t D_k(s) ds.$$

Let

$$E(t) = \sum_{k=1}^K E_k(t), \quad D(t) = \sum_{k=1}^K D_k(t), \quad V(t) = \sum_{k=1}^K V_k(t).$$

The estimates above imply

$$E(t) \leq C \int_0^t (E(s) + D(s)) ds + CV(t),$$

$$D(t) \leq \frac{C}{N_*} + C \int_0^t (E(s) + D(s) + V(s)) ds,$$

and

$$V(t) \leq \frac{C}{N_*} + C \int_0^t D(s) ds.$$

Substituting the last estimate into the first and absorbing repeated time integrals into the constant gives

$$E(t) \leq \frac{C}{N_*} + C \int_0^t (E(s) + D(s)) ds.$$

Together with the estimates for D and V , this yields

$$\Phi(t) := E(t) + D(t) + V(t) \leq \frac{C}{N_*} + C \int_0^t \Phi(s) ds.$$

Gronwall's inequality gives

$$\Phi(t) \leq \frac{C_T}{N_*}, \quad 0 \leq t \leq T.$$

The asserted bounds for each type follow from $D_k \leq D$ and $V_k \leq V$. The proof is complete. $\square$

Proposition SM2.8 (Propagation of chaos: asymptotic independence of particles of the same type). *Under the conditions of Theorem SM2.6, fix a type $k \in \{1, \dots, K\}$ and a terminal time $t \in [0, T]$. Let*

$$\widehat{X}_i^{k,N}(t) := \begin{cases} X_i^{k,N}(t), & \xi_i^{k,N}(t) = 1, \\ \partial, & \xi_i^{k,N}(t) = 0, \end{cases}$$

where ∂ is the cemetery point, and define the full probability measures

$$\widehat{\mu}_t^{k,N} := \mu_t^{k,N} + L_t^{k,N} \delta_\partial, \quad \widehat{\mu}_t^k := \mu_t^k + L_t^k \delta_\partial.$$

Then, for any fixed integer $m \geq 1$ and any pairwise distinct indices

$$i_1, \dots, i_m \in \{i : g(i) = k\},$$

we have

$$\mathcal{L} \left(\widehat{X}_{i_1}^{k,N}(t), \dots, \widehat{X}_{i_m}^{k,N}(t) \mid \mathcal{F}_t^{W^0} \right) \Longrightarrow (\widehat{\mu}_t^k)^{\otimes m}$$

weakly in probability. Equivalently, for any bounded Lipschitz test function $\Phi : \mathbf{E}^m \rightarrow \mathbb{R}$, where $\mathbf{E} = \mathbb{R} \cup \{\partial\}$,

$$\begin{aligned} & \mathbb{E} \left[\Phi \left(\widehat{X}_{i_1}^{k,N}(t), \dots, \widehat{X}_{i_m}^{k,N}(t) \right) \mid \mathcal{F}_t^{W^0} \right] \\ & \longrightarrow \int_{\mathbf{E}^m} \Phi(y_1, \dots, y_m) (\widehat{\mu}_t^k)^{\otimes m} (dy_1 \cdots dy_m) \end{aligned}$$

in probability.

In particular, if $\sigma_0 = 0$, then $\widehat{\mu}_t^k$ is deterministic, and the conditional convergence above reduces to the usual unconditional propagation of chaos:

$$\mathcal{L} \left(\widehat{X}_{i_1}^{k,N}(t), \dots, \widehat{X}_{i_m}^{k,N}(t) \right) \Longrightarrow (\widehat{\mu}_t^k)^{\otimes m}.$$

If the main text already incorporates the dead-particle state into $X_i^k(t)$, then one may directly write $\widehat{X}_i^{k,N}$ and $\widehat{\mu}_t^k$ as $X_i^{k,N}$ and μ_t^k , respectively, and obtain the shortened form

$$\mathcal{L} \left(X_{i_1}^{k,N}(t), \dots, X_{i_m}^{k,N}(t) \mid \mathcal{F}_t^{W^0} \right) \Longrightarrow (\mu_t^k)^{\otimes m}.$$

Proof. Use the synchronous coupling from the proof of Theorem SM2.6. For each i and k , the finite particle $X_i^{k,N}$ and the limiting particle $\bar{X}_i^k$ use the same initial state, the same idiosyncratic Brownian motion W_i^k , the same common Brownian motion W^0 , and the same Poisson random measure $\mathcal{N}_i^k$. Let

$$\widehat{\bar{X}}_i^k(t) := \begin{cases} \bar{X}_i^k(t), & \bar{\xi}_i^k(t) = 1, \\ \partial, & \bar{\xi}_i^k(t) = 0. \end{cases}$$

By the definition of the limiting McKean–Vlasov system, conditional on the common noise $\mathcal{F}_t^{W^0}$,

$$\widehat{\bar{X}}_{i_1}^k(t), \dots, \widehat{\bar{X}}_{i_m}^k(t)$$

are conditionally independent and identically distributed, with common conditional law

$$\widehat{\mu}_t^k = \mu_t^k + L_t^k \delta_\partial.$$

Therefore,

$$\mathcal{L} \left(\widehat{\bar{X}}_{i_1}^k(t), \dots, \widehat{\bar{X}}_{i_m}^k(t) \mid \mathcal{F}_t^{W^0} \right) = (\widehat{\mu}_t^k)^{\otimes m}.$$

It remains to show that the finite-particle tuple and the independent limiting tuple converge to each other in distance under the synchronous coupling. On $\mathbf{E} = \mathbb{R} \cup \{\partial\}$, choose a bounded metric d_∂ that agrees with the truncated Euclidean distance on $\mathbb{R}$ and satisfies

$$d_\partial(x, y) \leq |x - y| \wedge 1, \quad x, y \in \mathbb{R},$$

and

$$d_\partial(x, \partial) \leq 1, \quad x \in \mathbb{R}.$$

Then, for each i ,

$$d_\partial \left(\widehat{X}_i^{k,N}(t), \widehat{X}_i^k(t) \right) \leq |X_i^{k,N}(t) - \bar{X}_i^k(t)| \wedge 1 + |\xi_i^{k,N}(t) - \bar{\xi}_i^k(t)|.$$

The synchronous coupling estimates in Theorem SM2.6 give

$$\mathbb{E} \sup_{0 \leq s \leq T} |X_i^{k,N}(s) - \bar{X}_i^k(s)| \longrightarrow 0$$

and

$$\mathbb{E} \sup_{0 \leq s \leq T} |\xi_i^{k,N}(s) - \bar{\xi}_i^k(s)| \longrightarrow 0.$$

Hence

$$\mathbb{E} d_\partial \left(\widehat{X}_i^{k,N}(t), \widehat{X}_i^k(t) \right) \longrightarrow 0.$$

For fixed $i_1, \dots, i_m$, this gives

$$\mathbb{E} \sum_{r=1}^m d_\partial \left(\widehat{X}_{i_r}^{k,N}(t), \widehat{X}_{i_r}^k(t) \right) \longrightarrow 0.$$

Let $\Phi : \mathbf{E}^m \rightarrow \mathbb{R}$ be any bounded Lipschitz function. By the Lipschitz property,

$$\begin{aligned} & \left| \Phi \left(\widehat{X}_{i_1}^{k,N}(t), \dots, \widehat{X}_{i_m}^{k,N}(t) \right) - \Phi \left(\widehat{X}_{i_1}^k(t), \dots, \widehat{X}_{i_m}^k(t) \right) \right| \\ & \leq \text{Lip}(\Phi) \sum_{r=1}^m d_\partial \left(\widehat{X}_{i_r}^{k,N}(t), \widehat{X}_{i_r}^k(t) \right). \end{aligned}$$

Taking conditional expectation with respect to $\mathcal{F}_t^{W^0}$, and writing

$$\mathcal{Q}_{N,t}^{k,m} := \mathcal{L} \left(\widehat{X}_{i_1}^{k,N}(t), \dots, \widehat{X}_{i_m}^{k,N}(t) \mid \mathcal{F}_t^{W^0} \right),$$

we obtain

$$\begin{aligned} & \left| \int_{\mathbf{E}^m} \Phi d\mathcal{Q}_{N,t}^{k,m} - \int_{\mathbf{E}^m} \Phi d(\widehat{\mu}_t^k)^{\otimes m} \right| \\ & \leq \text{Lip}(\Phi) \mathbb{E} \left[\sum_{r=1}^m d_\partial \left(\widehat{X}_{i_r}^{k,N}(t), \widehat{X}_{i_r}^k(t) \right) \mid \mathcal{F}_t^{W^0} \right]. \end{aligned}$$

Taking outer expectation gives

$$\mathbb{E} \left| \int_{\mathbf{E}^m} \Phi d\mathcal{Q}_{N,t}^{k,m} - \int_{\mathbf{E}^m} \Phi d(\widehat{\mu}_t^k)^{\otimes m} \right| \leq \text{Lip}(\Phi) \sum_{r=1}^m \mathbb{E} d_\partial \left(\widehat{X}_{i_r}^{k,N}(t), \widehat{X}_{i_r}^k(t) \right) \longrightarrow 0.$$

Thus the difference above converges to zero in L^1 , and hence in probability, for every bounded Lipschitz Φ . To phrase the conclusion as bounded-Lipschitz convergence, take a countable determining subclass of the bounded-Lipschitz unit ball on the Polish space $\mathbb{E}^m$. The preceding estimate holds simultaneously on this countable subclass after a diagonal argument, and separability of the bounded-Lipschitz metric extends it to the full unit ball. Hence

$$d_{\text{BL}} \left(\mathcal{Q}_{N,t}^{k,m}, (\hat{\mu}_t^k)^{\otimes m} \right) \longrightarrow 0 \quad \text{in probability.}$$

Since the bounded Lipschitz distance metrizes weak convergence on the Polish space $\mathbb{E}^m$, the proposition follows. $\square$

SM3 Proofs for the matrix approximation boundary, tail gap, and spectral-radius threshold

Matched time-grid coupling convention The graph–matrix comparison in Theorem [SM3.4](#) is stated for the same discrete time grid used by the numerical system,

$$0 = t_0 < t_1 < \cdots < t_M = T, \quad \Delta t_m = t_m - t_{m-1}.$$

The graph system and matrix system are synchronously coupled: they use the same initial states, Brownian increments, common noise increments, and killing randomization variables. At each step, the state is first advanced by drift and diffusion to a pre-contagion value. New defaults are then sampled from the state-dependent killing probability

$$1 - \exp\{-\lambda_{g(i)}(X_{i,t_m}^{\text{pre}}, a_{i,t_m})\Delta t_m\},$$

using the same uniform random variable in the graph and matrix systems. After the new-default indicators are sampled, surviving nodes receive the contagion state shock

$$P_{i,t_m}^{\text{graph}} = \sum_{j=1}^N A_{ij} \Delta N_{j,t_m}^{\text{graph}}$$

in the graph system, and the type-average shock

$$P_{g(i),t_m}^{\text{matrix}} = \sum_{\ell=1}^K \hat{\Gamma}_{g(i)\ell} \Delta L_{\ell,t_m}^{\text{matrix}}$$

in the matrix system. Thus pressure affects future killing only through the state variable. The proof below therefore transfers pressure error first to state error and then to killing error; pressure is not treated as a separate direct argument of the kill intensity.

For each node i , define cumulative default indicators and new-default increments by

$$N_{i,t_m}^{\text{graph}} := \mathbf{1}_{\{\tau_i^{\text{graph}} \leq t_m\}}, \quad N_{i,t_m}^{\text{matrix}} := \mathbf{1}_{\{\tau_i^{\text{matrix}} \leq t_m\}},$$

and

$$\Delta N_{i,t_m}^{\text{graph}} := N_{i,t_m}^{\text{graph}} - N_{i,t_{m-1}}^{\text{graph}}, \quad \Delta N_{i,t_m}^{\text{matrix}} := N_{i,t_m}^{\text{matrix}} - N_{i,t_{m-1}}^{\text{matrix}}.$$

This synchronous construction makes the difference between the two systems arise from local network aggregation and the subsequent state feedback, rather than from resampling exogenous noise.

For any target node i and grid step t_m , let $\mathcal{H}_{i,m}^{\text{pre}}$ collect the common noise, controls, and the target-node coupled state up to the pre-sampling state at t_m . Define the pre-sampling type-level information by

$$\mathcal{F}_{i,t_m^-}^{\text{typ}} := \sigma\left(\mathcal{H}_{i,m}^{\text{pre}}, \{L_{r,t_q}^{\text{graph}}, L_{r,t_q}^{\text{matrix}} : 1 \leq r \leq K, 0 \leq q < m\}, \{P_{g(i),t_q}^{\text{matrix}} : 0 \leq q \leq m\}\right),$$

augmented, when needed, by the conditional type-level new-default probabilities

$$\bar{p}_{\ell,t_m} := \mathbb{E}[\Delta N_{j,t_m}^{\text{graph}} \mid \mathcal{F}_{i,t_m^-}^{\text{typ}}], \quad g(j) = \ell,$$

whenever this conditional mean is common inside the source type. The sigma-algebra $\mathcal{F}_{i,t_m^-}^{\text{typ}}$ is deliberately pre-sampling: it contains the information available before the identities of the newly defaulting source nodes at step m are realized, and it does not condition on the realized current type average $\Delta L_{\ell,t_m}^{\text{graph}}$. This convention keeps the within-type sampling statement separate from the realized sample mean that appears in the graph–matrix pressure decomposition. It is the conditional version of the information used by the matrix approximation, not the full node-level graph filtration after the current default set has been revealed.

Assumption SM3.1 (Within-type sampling regularity). There exists a constant $C_{\text{mix}} < \infty$, independent of N , such that for any target node i , source type ℓ , and grid time t_m , the new-default indicators satisfy the pre-sampling conditional estimate

$$\mathbb{E} \left[\left| \sum_{j:g(j)=\ell} A_{ij} \left(\Delta N_{j,t_m}^{\text{graph}} - \Delta L_{\ell,t_m}^{\text{graph}} \right) \right|^2 \middle| \mathcal{F}_{i,t_m^-}^{\text{typ}} \right] \leq C_{\text{mix}} \sum_{j:g(j)=\ell} A_{ij}^2 \quad \text{a.s.}$$

where $\Delta L_{\ell,t_m}^{\text{graph}} = L_{\ell,t_m}^{\text{graph}} - L_{\ell,t_{m-1}}^{\text{graph}}$. Equivalently, the unconditional estimate used below follows from this conditional form by the tower property. For cumulative-loss stability, the same square-edge estimate may be imposed on cumulative indicators with the corresponding pre-terminal type information:

$$\mathbb{E} \left[\left| \sum_{j:g(j)=\ell} A_{ij} \left(N_{j,t}^{\text{graph}} - L_{\ell,t}^{\text{graph}} \right) \right|^2 \middle| \mathcal{F}_{i,t^-}^{\text{typ}} \right] \leq C_{\text{mix}} \sum_{j:g(j)=\ell} A_{ij}^2 \quad \text{a.s.}$$

This is the within-type sampling condition used to control realized default-set fluctuations after type aggregation.

Remark SM3.2 (Sufficient conditions for within-type sampling regularity). The preceding assumption is a finite-sample regularity condition for default sets within a type. It is separate from propagation of chaos and is stated with pre-sampling type information to avoid conditioning on the realized type average in the same step. One sufficient condition is conditional exchangeable Bernoulli sampling. Conditional on $\mathcal{F}_{i,t_m^-}^{\text{typ}}$, suppose that $\{\Delta N_{j,t_m}^{\text{graph}} : g(j) = \ell\}$ are independent Bernoulli variables with common mean $\bar{p}_{\ell,t_m}$, and that the edge weights A_{ij} are measurable with respect to the same pre-sampling information. Then, writing $n_{\ell} = |\{j : g(j) = \ell\}|$ and $\bar{A}_{i\ell} = n_{\ell}^{-1} \sum_{j:g(j)=\ell} A_{ij}$,

$$\sum_{j:g(j)=\ell} A_{ij} \left(\Delta N_{j,t_m}^{\text{graph}} - \Delta L_{\ell,t_m}^{\text{graph}} \right) = \sum_{j:g(j)=\ell} (A_{ij} - \bar{A}_{i\ell}) (\Delta N_{j,t_m}^{\text{graph}} - \bar{p}_{\ell,t_m}),$$

and hence

$$\text{Var} \left(\sum_{j:g(j)=\ell} A_{ij} \left(\Delta N_{j,t_m}^{\text{graph}} - \Delta L_{\ell,t_m}^{\text{graph}} \right) \middle| \mathcal{F}_{i,t_m}^{\text{typ}} \right) \leq \sum_{j:g(j)=\ell} (A_{ij} - \bar{A}_{i\ell})^2 \leq \sum_{j:g(j)=\ell} A_{ij}^2.$$

The same conclusion also holds for sampling without replacement, or more generally under conditional negative dependence given the realized type count, because the covariance contribution is nonpositive. If weak positive dependence exists within the same type, it is enough to assume the weighted covariance bound

$$\sum_{\substack{j,m:g(j)=g(m)=\ell \\ j \neq m}} |A_{ij} A_{im}| \left| \text{Cov} \left(\Delta N_{j,t_m}^{\text{graph}}, \Delta N_{m,t_m}^{\text{graph}} \middle| \mathcal{F}_{i,t_m}^{\text{typ}} \right) \right| \leq C \sum_{j:g(j)=\ell} A_{ij}^2.$$

Thus Assumption SM3.1 says that the within-block default set does not generate concentration fluctuations larger than the square-edge scale. It is the local uniform sampling condition needed for matrix approximation, not a circular assumption inferred from the matrix approximation to be proved. In core-periphery and heavy-tailed networks this condition is not automatic; in those regimes the theorem should be read as a diagnostic bound conditional on the stated sampling regularity, while the numerical experiments separately measure how large the row-dispersion and square-edge-spread terms become when local concentration is strong.

Lemma SM3.3 (State-shock and killing stability under matched grid coupling). *Assume that the kill intensity $\lambda_k(x, a)$ is uniformly Lipschitz in x and uniformly bounded, and that the drift update is uniformly Lipschitz in the state and measure variables. For the synchronously coupled graph system and matrix system on the fixed time grid above, set*

$$E_{i,m} := X_{i,t_m}^{\text{graph}} - X_{i,t_m}^{\text{matrix}}.$$

Then there exists a constant $C < \infty$, independent of the particle number and of the sparse graph beyond the stated exposure bounds, such that for every target node i of type $k = g(i)$,

$$\begin{aligned} \mathbb{E} \max_{r \leq m} |E_{i,r}| &\leq C \sum_{q=1}^m \Delta t_q \mathbb{E} \max_{r \leq q} |E_{i,r}| \\ &\quad + C \sum_{q=1}^m \mathbb{E} |P_{i,t_q}^{\text{graph}} - P_{k,t_q}^{\text{matrix}}| \\ &\quad + C \sum_{\ell=1}^K \mathbb{E} \max_{r \leq m} |L_{\ell,t_r}^{\text{graph}} - L_{\ell,t_r}^{\text{matrix}}|. \end{aligned}$$

The pressure term appears without a factor Δt_q , because the matched update applies pressure as a one-step state jump. Moreover,

$$\mathbb{E} \left| N_{i,t_m}^{\text{graph}} - N_{i,t_m}^{\text{matrix}} \right| \leq C \sum_{q=1}^m \Delta t_q \mathbb{E} |E_{i,q}^{\text{pre}}|,$$

where $E_{i,q}^{\text{pre}}$ denotes the difference of the pre-contagion states at step q .

Proof. Under the matched grid convention, the two systems use the same killing randomization at each time step. Conditional on the pre-contagion states, the one-step default probabilities differ by at most

$$C \Delta t_m \left| X_{i,t_m}^{\text{graph,pre}} - X_{i,t_m}^{\text{matrix,pre}} \right|$$

because λ_k is Lipschitz in the state variable. Summing over time steps gives the displayed killing-indicator estimate.

For the state update, the Brownian increments are synchronously coupled and cancel in the difference. The remaining terms are the Lipschitz drift difference, the graph–matrix pressure-shock difference, and the type-level loss-feedback difference. Since the pressure shock is applied as a state jump at each grid step, the pressure difference enters the state recursion directly. This gives

$$\begin{aligned} \mathbb{E} \max_{r \leq m} |E_{i,r}| &\leq C \sum_{q=1}^m \Delta t_q \mathbb{E} \max_{r \leq q} |E_{i,r}| \\ &\quad + C \sum_{q=1}^m \mathbb{E} |P_{i,t_q}^{\text{graph}} - P_{k,t_q}^{\text{matrix}}| \\ &\quad + C \sum_{\ell=1}^K \mathbb{E} \max_{r \leq m} |L_{\ell,t_r}^{\text{graph}} - L_{\ell,t_r}^{\text{matrix}}|. \end{aligned}$$

The proof is complete. $\square$

Theorem SM3.4 (Finite-grid absolute structural bound for the expected-loss gap of matrix approximation). *Assume that the graph system and its aggregate matrix system are synchronously coupled on the matched fixed grid above under the same initial states, the same Brownian noises, and the same killing randomization. Assume that the kill intensity is Lipschitz and bounded in the state variable, uniformly over controls, that the drift is Lipschitz in the state and measure variables, that contagion weights are nonnegative, that total row exposures are uniformly bounded, that Assumption SM3.1 holds, and that the type shares satisfy $\pi_k \geq \underline{\pi} > 0$ uniformly over k and over the graph sequence. Let*

$$B_{i\ell} = \sum_{j:g(j)=\ell} A_{ij}, \quad \hat{\Gamma}_{k\ell} = \frac{1}{n_k} \sum_{i:g(i)=k} B_{i\ell}.$$

Define

$$R_k(A) := \left[\frac{1}{n_k} \sum_{i:g(i)=k} \left(\sum_{\ell=1}^K |B_{i\ell} - \hat{\Gamma}_{k\ell}| \right)^2 \right]^{1/2},$$

and

$$S_k(A) := \frac{1}{n_k} \sum_{i:g(i)=k} \sum_{\ell=1}^K \left(\sum_{j:g(j)=\ell} A_{ij}^2 \right)^{1/2}.$$

Then there exists a constant $C_{T,\Delta} < \infty$, depending only on T , the fixed grid through M and the grid-stability constants, the Lipschitz constants, the upper bound on the kill intensity, the total exposure bound, the sampling-regularity constant, and $\underline{\pi}^{-1}$, such that

$$\left| \mathbb{E} L_T^{\text{graph}} - \mathbb{E} L_T^{\text{matrix}} \right| \leq C_{T,\Delta} \sum_{k=1}^K \pi_k (R_k(A) + S_k(A)).$$

Proof. Write

$$D_A := \sum_{k=1}^K \pi_k (R_k(A) + S_k(A)).$$

Fix a target node i of type k . At grid time t_m , decompose the pressure difference as

$$\begin{aligned} P_{i,t_m}^{\text{graph}} - P_{k,t_m}^{\text{matrix}} &= \sum_{\ell=1}^K \sum_{j:g(j)=\ell} A_{ij} (\Delta N_{j,t_m}^{\text{graph}} - \Delta L_{\ell,t_m}^{\text{graph}}) \\ &\quad + \sum_{\ell=1}^K (B_{i\ell} - \hat{\Gamma}_{k\ell}) \Delta L_{\ell,t_m}^{\text{graph}} \\ &\quad + \sum_{\ell=1}^K \hat{\Gamma}_{k\ell} (\Delta L_{\ell,t_m}^{\text{graph}} - \Delta L_{\ell,t_m}^{\text{matrix}}). \end{aligned}$$

The first term is controlled by Assumption [SM3.1](#) and Cauchy–Schwarz:

$$\mathbb{E} \left| \sum_{j:g(j)=\ell} A_{ij} (\Delta N_{j,t_m}^{\text{graph}} - \Delta L_{\ell,t_m}^{\text{graph}}) \right| \leq C \left(\sum_{j:g(j)=\ell} A_{ij}^2 \right)^{1/2}.$$

The second term is bounded by $\sum_{\ell} |B_{i\ell} - \hat{\Gamma}_{k\ell}|$, since $0 \leq \Delta L_{\ell,t_m}^{\text{graph}} \leq 1$. The third term is bounded by the uniform row-exposure bound times

$$\sum_{\ell=1}^K |\Delta L_{\ell,t_m}^{\text{graph}} - \Delta L_{\ell,t_m}^{\text{matrix}}| \leq 2 \sum_{\ell=1}^K \max_{r \leq m} |L_{\ell,t_r}^{\text{graph}} - L_{\ell,t_r}^{\text{matrix}}|.$$

Averaging over target nodes and then over types gives

$$\sum_{k=1}^K \pi_k \frac{1}{n_k} \sum_{i:g(i)=k} \mathbb{E} |P_{i,t_m}^{\text{graph}} - P_{k,t_m}^{\text{matrix}}| \leq CD_A + C_{\underline{\pi}} V_m,$$

where $C_{\underline{\pi}}$ depends on the uniform lower bound $\underline{\pi}$. This dependence enters only when the unweighted sum of type-loss feedback errors is converted into the type-weighted norm below, since $\sum_{\ell} e_{\ell} \leq \underline{\pi}^{-1} \sum_{\ell} \pi_{\ell} e_{\ell}$. Here

$$V_m := \sum_{\ell=1}^K \pi_{\ell} \mathbb{E} \max_{r \leq m} |L_{\ell,t_r}^{\text{graph}} - L_{\ell,t_r}^{\text{matrix}}|.$$

Set

$$U_m := \sum_{k=1}^K \pi_k \frac{1}{n_k} \sum_{i:g(i)=k} \mathbb{E} \max_{r \leq m} |X_{i,t_r}^{\text{graph}} - X_{i,t_r}^{\text{matrix}}|.$$

Lemma [SM3.3](#) and the pressure estimate imply

$$U_{m+1} \leq (1 + C\Delta t_m)U_m + C\Delta t_m V_m + CD_A + C_{\underline{\pi}} V_m.$$

The term $CD_A + C_{\underline{\pi}} V_m$ enters through the one-step pressure jump.

The loss difference is controlled through the killing probability. Since

$$p_k(x, a; \Delta t_m) = 1 - \exp\{-\lambda_k(x, a)\Delta t_m\},$$

bounded Lipschitz killing and the common killing randomization give

$$V_{m+1} \leq V_m + C\Delta t_m (U_m + V_m).$$

Combining the two recursive inequalities over the fixed grid and applying discrete Gronwall yields

$$\max_{0 \leq m \leq M} (U_m + V_m) \leq C_{T,\Delta} D_A.$$

Since aggregate loss is the type-weighted average of default indicators,

$$\left| \mathbb{E} L_T^{\text{graph}} - \mathbb{E} L_T^{\text{matrix}} \right| \leq V_M \leq C_{T,\Delta} D_A.$$

The proof is complete. $\square$

Corollary SM3.5 (Share-controlled matrix-approximation diagnostic). *Under the assumptions of Theorem SM3.4, define*

$$q_k(A) := \max_{i:g(i)=k, 1 \leq \ell \leq K, j:g(j)=\ell} \frac{A_{ij}}{B_{i\ell} \vee \varepsilon_0}, \quad \varepsilon_0 > 0.$$

If the block row exposures $B_{i\ell}$ are uniformly bounded, then

$$S_k(A) \leq C \sqrt{q_k(A)}.$$

Consequently,

$$\left| \mathbb{E} L_T^{\text{graph}} - \mathbb{E} L_T^{\text{matrix}} \right| \leq C_{T,\Delta} \sum_{k=1}^K \pi_k \left(R_k(A) + \sqrt{q_k(A)} \right).$$

Proof. For fixed i and ℓ , the definition of $q_k(A)$ gives

$$A_{ij} \leq q_k(A) (B_{i\ell} \vee \varepsilon_0), \quad j : g(j) = \ell.$$

When $B_{i\ell} > 0$, this implies

$$\sum_{j:g(j)=\ell} A_{ij}^2 \leq q_k(A) B_{i\ell}^2$$

up to the harmless ε_0 convention for empty or negligible blocks. Uniform boundedness of $B_{i\ell}$ therefore yields

$$\left(\sum_{j:g(j)=\ell} A_{ij}^2 \right)^{1/2} \leq C \sqrt{q_k(A)}.$$

Averaging over $i : g(i) = k$ and summing over ℓ gives the stated bound on $S_k(A)$, and the theorem gives the conclusion. $\square$

Theorem SM3.6 (Nonzero lower bound for the tail-loss gap in heterogeneous networks). *Fix a terminal time $T > 0$, and let*

$$Q_\alpha(Y) = \inf\{x \in \mathbb{R} : \mathbb{P}(Y \leq x) \geq \alpha\}$$

denote the α -quantile of a random variable Y . There exist two sequences of weighted directed networks

$$A_n^{\text{hom}}, \quad A_n^{\text{hub}},$$

and the same type-wise initial-state distributions, killing parameters, control rule, and type assignment, such that the following conclusions hold simultaneously.

Let

$$V_n = S_n \cup R_n, \quad S_n \cap R_n = \emptyset,$$

where S_n is the set of source nodes of type one and R_n is the set of shocked nodes of type two,

$$|S_n| = s_n, \quad |R_n| = m_n, \quad s_n + m_n = n,$$

and assume that

$$\frac{s_n}{n} \longrightarrow \pi_1, \quad \frac{m_n}{n} \longrightarrow \pi_2, \quad \pi_1, \pi_2 \in (0, 1), \quad \pi_1 + \pi_2 = 1.$$

The type function is defined by

$$g(i) = \begin{cases} 1, & i \in S_n, \\ 2, & i \in R_n. \end{cases}$$

Both network sequences induce type matrices through the same aggregation rule

$$\hat{\Gamma}_{k\ell}(A) = \frac{1}{|\{i : g(i) = k\}|} \sum_{i:g(i)=k} \sum_{j:g(j)=\ell} A_{ij},$$

and satisfy

$$\hat{\Gamma}(A_n^{\text{hom}}) = \hat{\Gamma}(A_n^{\text{hub}}) \quad \text{for all } n.$$

Assume that the kill intensity $\lambda_2(x, a)$ of type two is nonincreasing in the state variable x , namely, for any fixed control a ,

$$x_1 \leq x_2 \implies \lambda_2(x_1, a) \geq \lambda_2(x_2, a).$$

Thus, if a type-two node is at state x_0 before receiving pressure, has control a_0 , and undergoes killing over a terminal interval of length $\Delta t > 0$, then the terminal kill probability corresponding to pressure level $p \geq 0$ is

$$r(p) = 1 - \exp\{-\lambda_2(x_0 - p, a_0)\Delta t\},$$

which is nondecreasing in p .

Assume that there exist constants $a > 0$, $q > 0.05$, and $\rho \in (q, 1)$ such that

$$p_+ = r(a), \quad p_- = r(\rho a)$$

satisfy the strict separation condition

$$\pi_2(p_+ - p_-) - \pi_1\rho > 0. \tag{A}$$

Then there exist constants $\delta > 0$ and $N_0 \in \mathbb{N}$ such that, for all $n > N_0$,

$$Q_{0.95}(L_T^{\text{hub},n}) - Q_{0.95}(L_T^{\text{hom},n}) \geq \delta.$$

Here $L_T^{\text{hub},n}$ and $L_T^{\text{hom},n}$ denote the aggregate losses of the two graph systems at time T .

In addition, there exists a source-side hub node $h_n \in S_n$ in A_n^{hub} such that

$$\frac{\sum_{i \in R_n} A_{ih_n}^{\text{hub}}}{\sum_{j \in S_n} \sum_{i \in R_n} A_{ij}^{\text{hub}}} = 1,$$

whereas the homogeneous network satisfies

$$\max_{j \in S_n} \frac{\sum_{i \in R_n} A_{ij}^{\text{hom}}}{\sum_{\ell \in S_n} \sum_{i \in R_n} A_{i\ell}^{\text{hom}}} \longrightarrow 0.$$

Proof. Take $K = 2$. Fix a constant $a > 0$, and let $h_n \in S_n$ be the hub node. Define the hub network by

$$A_{ij}^{\text{hub}} = \begin{cases} a, & i \in R_n, j = h_n, \\ 0, & \text{otherwise,} \end{cases}$$

and define the homogeneous network by

$$A_{ij}^{\text{hom}} = \begin{cases} a/s_n, & i \in R_n, j \in S_n, \\ 0, & \text{otherwise.} \end{cases}$$

Then the only nonzero edge weights in the two network sequences are in the type block $(2, 1)$. For the hub network,

$$\hat{\Gamma}_{21}(A_n^{\text{hub}}) = \frac{1}{m_n} \sum_{i \in R_n} \sum_{j \in S_n} A_{ij}^{\text{hub}} = \frac{1}{m_n} \sum_{i \in R_n} a = a.$$

For the homogeneous network,

$$\hat{\Gamma}_{21}(A_n^{\text{hom}}) = \frac{1}{m_n} \sum_{i \in R_n} \sum_{j \in S_n} A_{ij}^{\text{hom}} = \frac{1}{m_n} \sum_{i \in R_n} \sum_{j \in S_n} \frac{a}{s_n} = a.$$

All other type blocks have zero aggregate exposure. Hence

$$\hat{\Gamma}(A_n^{\text{hub}}) = \hat{\Gamma}(A_n^{\text{hom}}) \quad \text{for all } n.$$

At the same time,

$$\frac{\sum_{i \in R_n} A_{ih_n}^{\text{hub}}}{\sum_{j \in S_n} \sum_{i \in R_n} A_{ij}^{\text{hub}}} = \frac{m_n a}{m_n a} = 1,$$

whereas

$$\max_{j \in S_n} \frac{\sum_{i \in R_n} A_{ij}^{\text{hom}}}{\sum_{\ell \in S_n} \sum_{i \in R_n} A_{i\ell}^{\text{hom}}} = \frac{m_n a/s_n}{m_n a} = \frac{1}{s_n} \rightarrow 0.$$

This proves that the two network sequences are identical at the type-aggregation level but asymptotically separated in source-side exposure concentration.

We next construct the common loss process. Work on a two-stage discrete-time convention: first-stage type-one defaults are sampled on $[0, t_0]$, their network pressure is applied at t_0 , and the affected type-two nodes are then killed over a terminal interval of length Δt . Choosing t_0 and Δt sufficiently small makes any feedback from earlier defaults before the designed pressure step an $O(t_0 + \Delta t)$ perturbation, which is absorbed into the strict separation margin in (A). Let type-one nodes default independently during the time interval $[0, t_0]$, and write

$$D_j = \mathbf{1}_{\{\tau_j \leq t_0\}}, \quad j \in S_n.$$

Assume that

$$D_j \sim \text{Bernoulli}(q), \quad j \in S_n,$$

and that these random variables are mutually independent, where $q > 0.05$. This construction can be implemented by the same type-wise initial-state distributions and the same state-dependent kill intensity. For example, set type-one nodes to the same deterministic state and the same control level during $[0, t_0]$ so that their kill probability equals q .

After time t_0 , type-two nodes first receive network pressure and then undergo killing over a terminal interval of length Δt . Suppose that all type-two nodes have the same state x_0 and the same control a_0 before receiving pressure. For pressure level $p \geq 0$, their terminal kill probability is

$$r(p) = 1 - \exp\{-\lambda_2(x_0 - p, a_0)\Delta t\}.$$

By the monotonicity of $\lambda_2(\cdot, a_0)$ in the state variable, $r(p)$ is nondecreasing in p .

First consider the hub network. On the event $\{D_{h_n} = 1\}$, each type-two node $i \in R_n$ receives graph pressure

$$P_i^{\text{hub}} = \sum_{j \in S_n} A_{ij}^{\text{hub}} D_j = a$$

at time t_0 . Hence, conditional on the first-stage default vector and on the event $\{D_{h_n} = 1\}$, every type-two node has terminal-interval kill probability p_+ . Let Z_n^{hub} be the number of newly killed type-two nodes during the terminal interval. By conditional independence of the terminal killing process across nodes,

$$Z_n^{\text{hub}} \mid \{D_{h_n} = 1\} \sim \text{Binomial}(m_n, p_+).$$

Therefore, for any $\varepsilon \in (0, p_+)$, the weak law of large numbers gives

$$\mathbb{P}\left(\frac{Z_n^{\text{hub}}}{m_n} \geq p_+ - \varepsilon \mid D_{h_n} = 1\right) \rightarrow 1.$$

Since

$$\mathbb{P}(D_{h_n} = 1) = q > 0.05,$$

there exists $N_1 \in \mathbb{N}$ such that, for all $n > N_1$,

$$\mathbb{P}\left(Z_n^{\text{hub}} \geq m_n(p_+ - \varepsilon)\right) > 0.05.$$

The aggregate loss is at least the type-two newly killed loss in the terminal interval, so

$$L_T^{\text{hub},n} \geq \frac{Z_n^{\text{hub}}}{n}.$$

Thus

$$\mathbb{P}\left(L_T^{\text{hub},n} \geq \frac{m_n}{n}(p_+ - \varepsilon)\right) > 0.05.$$

By the definition of the quantile,

$$Q_{0.95}\left(L_T^{\text{hub},n}\right) \geq \frac{m_n}{n}(p_+ - \varepsilon), \quad n > N_1. \quad (1)$$

This estimate uses only the terminal newly killed type-two nodes as a lower bound for aggregate loss. It does not include the initial type-one default loss and is therefore conservative.

Now consider the homogeneous network. Let

$$R_n^S = \sum_{j \in S_n} D_j.$$

Then

$$R_n^S \sim \text{Binomial}(s_n, q),$$

and each type-two node receives graph pressure

$$P_i^{\text{hom}} = \sum_{j \in S_n} A_{ij}^{\text{hom}} D_j = \frac{a}{s_n} \sum_{j \in S_n} D_j = a \frac{R_n^S}{s_n}$$

at time t_0 . Since $\rho > q$, the Chernoff bound gives a constant $c_\rho > 0$ such that

$$\mathbb{P} \left(\frac{R_n^S}{s_n} > \rho \right) \leq \exp\{-c_\rho s_n\}. \quad (2)$$

On the event

$$G_n = \left\{ \frac{R_n^S}{s_n} \leq \rho \right\},$$

the pressure on every type-two node satisfies

$$P_i^{\text{hom}} \leq \rho a.$$

By the monotonicity of r , the kill probability of type-two nodes in the terminal interval is at most

$$p_- = r(\rho a).$$

Let Z_n^{hom} be the number of newly killed type-two nodes in the homogeneous network during the terminal interval. Conditional on the first-stage default vector and on the event G_n , Z_n^{hom} is stochastically dominated by

$$\text{Binomial}(m_n, p_-).$$

Therefore, for any $\varepsilon > 0$, Hoeffding's inequality gives

$$\mathbb{P} \left(\frac{Z_n^{\text{hom}}}{m_n} > p_- + \varepsilon \mid G_n \right) \rightarrow 0. \quad (3)$$

By (2) and (3), there exists $N_2 \in \mathbb{N}$ such that, for all $n > N_2$,

$$\mathbb{P} \left(\frac{R_n^S}{s_n} \leq \rho, \frac{Z_n^{\text{hom}}}{m_n} \leq p_- + \varepsilon \right) \geq 0.95.$$

On this event, the terminal aggregate loss is at most the type-one initial default loss plus the type-two terminal new loss:

$$L_T^{\text{hom},n} \leq \frac{R_n^S}{n} + \frac{Z_n^{\text{hom}}}{n} \leq \frac{s_n}{n} \rho + \frac{m_n}{n} (p_- + \varepsilon).$$

By the definition of the quantile,

$$Q_{0.95}(L_T^{\text{hom},n}) \leq \frac{s_n}{n} \rho + \frac{m_n}{n} (p_- + \varepsilon), \quad n > N_2. \quad (4)$$

Combining (1) and (4), we obtain, for all

$$n > \max\{N_1, N_2\},$$

$$\begin{aligned}
& Q_{0.95}(L_T^{\text{hub},n}) - Q_{0.95}(L_T^{\text{hom},n}) \\
& \geq \frac{m_n}{n}(p_+ - \varepsilon) - \left[\frac{s_n}{n}\rho + \frac{m_n}{n}(p_- + \varepsilon) \right] \\
& = \frac{m_n}{n}(p_+ - p_- - 2\varepsilon) - \frac{s_n}{n}\rho.
\end{aligned} \tag{5}$$

As $n \rightarrow \infty$, the right-hand side converges to

$$\pi_2(p_+ - p_- - 2\varepsilon) - \pi_1\rho.$$

By the strict separation condition (A), we can choose $\varepsilon > 0$ small enough so that

$$\Delta_\varepsilon := \pi_2(p_+ - p_- - 2\varepsilon) - \pi_1\rho > 0.$$

Then, by the convergence in (5), there exists $N_0 \in \mathbb{N}$ such that, for all $n > N_0$,

$$Q_{0.95}(L_T^{\text{hub},n}) - Q_{0.95}(L_T^{\text{hom},n}) \geq \frac{\Delta_\varepsilon}{2}.$$

Taking

$$\delta = \frac{\Delta_\varepsilon}{2} > 0$$

gives

$$Q_{0.95}(L_T^{\text{hub},n}) - Q_{0.95}(L_T^{\text{hom},n}) \geq \delta, \quad n > N_0.$$

The proof is complete. $\square$

Remark SM3.7 (Feasibility of the separation condition). Condition (A) is a separation condition requiring the tail synchronous killing effect to dominate the effect of uniformly dispersed exposure. This condition is not vacuous. Given $\rho > q > 0.05$, if the share of type-one nodes is small and the share of shocked type-two nodes is large, then $\pi_1\rho$ can be made arbitrarily small. For example, take $\pi_1 = 0.05$, $\pi_2 = 0.95$, $q = 0.06$, and $\rho = 0.08$. Then it suffices that

$$p_+ - p_- > \frac{\pi_1\rho}{\pi_2} \approx 0.00421.$$

When the type-two kill intensity is sufficiently steep near the risk boundary, the difference in kill probabilities induced by pressures a and ρa , namely $p_+ - p_-$, can strictly exceed this threshold. Thus one default of the hub node can push many type-two nodes into the high-kill region with probability greater than 5%, while the same aggregate exposure in the homogeneous network remains controlled at the 0.95 tail by dispersion.

Corollary SM3.8 (Embedding the tail-gap construction in larger finite type systems). *Let $K \geq 2$, and choose two distinct active types s and r with asymptotic shares $\pi_s, \pi_r > 0$. Suppose the two-type separation condition in Theorem SM3.6 holds after replacing (π_1, π_2) by (π_s, π_r) and using the source type s and target type r . On the remaining type blocks, assume either that the common terminal-loss contribution is deterministic, or more generally that it is coupled identically in the hub and homogeneous graph sequences, is independent of the active (r, s) construction, and has no atom at the shifted 0.95 quantile used in the active-block comparison. Then there exist two K -type graph sequences with the same type assignment, the same aggregate matrix, and identical dynamics on all type blocks except the block (r, s) , such that their 0.95 terminal-loss quantiles differ by at least a positive constant for all sufficiently large n .*

Proof. Embed the source set S_n of Theorem SM3.6 inside type s and the target set R_n inside type r . On the block (r, s) use the hub and homogeneous edge constructions from the theorem. On all remaining type blocks use the same edge weights in the two graph sequences; for instance, take these blocks to be zero and choose their initial states so that they contribute deterministic zero terminal loss in both systems. The type-aggregated matrices remain identical because the only modified block has the same block average in the hub and homogeneous constructions. If the inactive blocks contribute deterministic zero terminal loss, they do not affect the positive separation created on the active (r, s) block. In the more general case, their common terminal-loss contribution shifts both terminal-loss distributions by the same independent component. The no-atom condition at the shifted quantile preserves the strict quantile separation after a possible reduction of the margin. The proof of Theorem SM3.6, with (π_s, π_r) replacing (π_1, π_2) and with the resulting shifted quantile margin, gives the stated positive quantile gap. $\square$

Remark SM3.9 (Robustness of the embedding). The embedding does not rely on all other types having zero loss. The same conclusion holds if the inactive types have nonzero terminal losses that are coupled identically in the hub and homogeneous graph sequences and are independent of the active (r, s) tail construction, or more generally if their common contribution has no atom at the relevant shifted quantile. In that case the inactive-type losses shift both terminal-loss distributions by the same common component, and the strict separation condition in Theorem SM3.6 is adjusted only through the resulting type-share and quantile-margin constants. This turns the two-type construction into a robust local block embedding rather than a purely zero-background example.

Remark SM3.10 (No-atom condition in finite simulations). The no-atom condition in Corollary SM3.8 is a continuous-time or large-population regularity condition. In a finite-particle discrete-time simulation, terminal losses take values on a finite grid and hence have atoms. The condition holds generically after an arbitrarily small continuous perturbation of initial states, idiosyncratic shocks, or terminal observation time; the limiting continuous-time construction is the object used in the corollary.

Lemma SM3.11 (Primitive sufficient condition for the low-loss second-order approximation). *Let $c_0 = \lambda_0 \Delta t / \mathfrak{d}_\kappa$ and $M = c_0 \Gamma$. Suppose that in the low-loss region the type- k conditional mean new-loss probability can be written as*

$$\bar{q}_k^N(Y) = q((\Gamma Y)_k) + \varepsilon_{k,N}(Y), \quad |\varepsilon_{k,N}(Y)| \leq r_N,$$

where q satisfies the uniform expansion

$$|q(p) - c_0 p| \leq C_q p^2, \quad 0 \leq p \leq \eta,$$

and the row sums of Γ are uniformly bounded. If $\|Y\|_1 \leq \eta_0$ with η_0 small enough that $(\Gamma Y)_k \leq \eta$ for every k , then

$$\left\| \mathbb{E}(Y_{m+1}^N \mid \mathcal{F}_m) - M Y_m^N \right\|_1 \leq C \|Y_m^N\|_1^2 + K r_N.$$

If, in addition, the conditional new-default sampling fluctuation within each type has variance bounded by C/N in the low-loss region, then the conditional second-moment estimate in Theorem SM3.12 follows.

Proof. For $\|Y\|_1 \leq \eta_0$, the bounded row sums give $(\Gamma Y)_k \leq C_\Gamma \|Y\|_1 \leq \eta$. Hence

$$\begin{aligned} \left| \bar{q}_k^N(Y) - c_0 (\Gamma Y)_k \right| &\leq |q((\Gamma Y)_k) - c_0 (\Gamma Y)_k| + |\varepsilon_{k,N}(Y)| \\ &\leq C_q (\Gamma Y)_k^2 + r_N \leq C \|Y\|_1^2 + r_N. \end{aligned}$$

Summing over the finite set of types gives the displayed vector estimate. The variance statement is the standard conditional Bernoulli or weak-dependence estimate for type-level averages on the low-loss region. This lemma records exactly which primitive local expansion is needed by the spectral-radius theorem; the theorem itself uses the resulting second-order approximation as an explicit input. $\square$

Theorem SM3.12 (Spectral radius and local cascade threshold). *Consider the finite-type contagion system without control, with homogeneous killing sensitivity and a nonnegative irreducible contagion matrix. Let the type set be $\{1, \dots, K\}$, with type shares $\pi_k > 0$ and $\sum_{k=1}^K \pi_k = 1$. Let*

$$\Gamma = (\Gamma_{k\ell})_{1 \leq k, \ell \leq K}$$

be a nonnegative and irreducible contagion matrix. The system evolves in discrete time $t_m = m\Delta t$, and the type new-loss vector is denoted by

$$Y_m^N = \left(\Delta L_m^{1,N}, \dots, \Delta L_m^{K,N} \right)^\top \in \mathbb{R}_+^K.$$

In the low-loss region, the contagion pressure of type k is

$$P_m^k = \sum_{\ell=1}^K \Gamma_{k\ell} Y_m^{\ell,N}.$$

Assume that drift mean reversion creates a positive damping factor $\mathfrak{d}_\kappa > 0$ against pressure shocks within one time step, and that, as pressure $p \downarrow 0$, the new kill probability induced by pressure for a single surviving particle admits the uniform expansion

$$q(p) = \frac{\lambda_0 \Delta t}{\mathfrak{d}_\kappa} p + O(p^2),$$

where $\lambda_0 > 0$, and the remainder is uniform across types and nodes. Equivalently, there exist $\eta > 0$ and $C_q < \infty$ such that, whenever $0 \leq p \leq \eta$,

$$\left| q(p) - \frac{\lambda_0 \Delta t}{\mathfrak{d}_\kappa} p \right| \leq C_q p^2.$$

Define the linear cascade matrix by

$$M := \frac{\lambda_0 \Delta t}{\mathfrak{d}_\kappa} \Gamma,$$

and define the effective contagion strength by

$$R_{\text{eff}} := \rho(M) = \rho(\Gamma) \frac{\lambda_0 \Delta t}{\mathfrak{d}_\kappa}.$$

Assume also that the new-loss process in the low-loss region satisfies the following second-order approximation. There exist $\eta_0 > 0$, $C < \infty$, and $r_N \downarrow 0$ such that, whenever $\|Y_m^N\|_1 \leq \eta_0$,

$$\left\| \mathbb{E} \left[Y_{m+1}^N \mid \mathcal{F}_m \right] - M Y_m^N \right\|_1 \leq C \|Y_m^N\|_1^2 + r_N,$$

and there exists the same constant $C < \infty$ such that

$$\mathbb{E} \left[\left\| Y_{m+1}^N - \mathbb{E}(Y_{m+1}^N \mid \mathcal{F}_m) \right\|_1^2 \mid \mathcal{F}_m \right] \leq \frac{C}{N} \left(1 + \|Y_m^N\|_1 \right).$$

Define the cascade event by

$$C_N(\varepsilon) := \left\{ \exists m \leq T_N : \sum_{k=1}^K \pi_k \Delta L_m^{k,N} \geq \varepsilon \right\},$$

where $\varepsilon > 0$ is fixed. In the subcritical statement below, take $\varepsilon \in (0, \eta_0)$. In the supercritical statement, ε is further restricted to a sufficiently small interval $(0, \varepsilon_*)$ specified by the low-loss region and the equivalence constants in the proof. Assume that the initial seed satisfies

$$\|Y_0^N\|_1 \rightarrow 0, \quad N\|Y_0^N\|_1^2 \rightarrow \infty,$$

and that its projection on the Perron–Frobenius principal direction is nondegenerate: there exists $b_0 > 0$ such that

$$\mathbb{P}\left(u^\top Y_0^N \geq b_0 \|Y_0^N\|_1\right) \rightarrow 1,$$

where $u \gg 0$ is the left Perron–Frobenius principal eigenvector of M .

If $R_{\text{eff}} < 1$, assume also that the observation window satisfies

$$T_N \rightarrow \infty, \quad T_N r_N \rightarrow 0, \quad T_N = o(N).$$

Then

$$\mathbb{P}(C_N(\varepsilon)) \rightarrow 0.$$

If $R_{\text{eff}} > 1$, then there exists $\varepsilon_* \in (0, \eta_0)$ such that the following conclusion holds for every fixed $\varepsilon \in (0, \varepsilon_*)$. Assume that there exists $\delta > 0$ such that the observation window satisfies

$$T_N \geq (1 + \delta) \frac{\log(\varepsilon / \|Y_0^N\|_1)}{\log R_{\text{eff}}},$$

and that the remainder and the initial seed satisfy the compatibility condition

$$r_N \log\left(\frac{1}{\|Y_0^N\|_1}\right) = o(\|Y_0^N\|_1),$$

while finite-sample fluctuations satisfy

$$\frac{N\|Y_0^N\|_1^2}{\log(1/\|Y_0^N\|_1)} \rightarrow \infty.$$

Then

$$\mathbb{P}(C_N(\varepsilon)) \rightarrow 1.$$

Therefore, in the low-loss linear region, the spectral radius of the contagion matrix determines the local phase-transition threshold for systemic cascades through R_{eff} . The critical value is

$$R^* = 1.$$

Remark SM3.13 (Compatibility condition and cascade-event convention). The supercritical statement is local in two senses. First, the compatibility condition

$$r_N \log(1/\|Y_0^N\|_1) = o(\|Y_0^N\|_1)$$

requires the deterministic approximation residual to be smaller than the Perron–Frobenius growth generated by the initial seed on the logarithmic hitting window. For example, if $r_N = O(N^{-1})$ and $\|Y_0^N\|_1 = N^{-\alpha}$, the condition is satisfied for every $0 < \alpha < 1$. Second, $C_N(\varepsilon)$ is defined through one-step new losses because the theorem studies local amplification of the increment vector $Y_m^N = \Delta L_m^N$. On the low-loss window used here, this increment-threshold event is the local cascade counterpart of the terminal-loss cascade event used in the simulations: reaching a fixed increment threshold before leaving the linearization region necessarily creates a corresponding increase in cumulative loss, while the terminal-loss event remains the empirical risk metric in the numerical sections.

Proof. Write

$$c_0 := \frac{\lambda_0 \Delta t}{\mathfrak{d}_\kappa}, \quad M = c_0 \Gamma.$$

Since Γ is nonnegative and irreducible, the Perron–Frobenius theorem gives a positive right principal eigenvector $v \gg 0$ and a positive left principal eigenvector $u \gg 0$. Normalize them so that

$$u^\top v = 1.$$

Then

$$Mv = R_{\text{eff}}v, \quad u^\top M = R_{\text{eff}}u^\top.$$

Since $u \gg 0$ and $\pi_k > 0$, the quantities $\|y\|_1$, $u^\top y$, and $\sum_k \pi_k y_k$ are equivalent on the finite-dimensional positive cone $\mathbb{R}_+^K$. Specifically, there exist constants $a_1, a_2, a_3, a_4 > 0$ such that, for any $y \in \mathbb{R}_+^K$,

$$a_1 \|y\|_1 \leq u^\top y \leq a_2 \|y\|_1, \quad a_3 u^\top y \leq \sum_{k=1}^K \pi_k y_k \leq a_4 \|y\|_1.$$

We first prove the subcritical case. Suppose that $R_{\text{eff}} < 1$. Choose r such that

$$R_{\text{eff}} < r < 1.$$

Since the spectral radius of M is smaller than r , there exists a vector norm $\|\cdot\|_M$, equivalent to $\|\cdot\|_1$, such that the induced matrix norm satisfies

$$\|M\|_M \leq r.$$

Equivalence of norms in finite dimension gives constants $B_1, B_2 > 0$ such that

$$B_1 \|y\|_1 \leq \|y\|_M \leq B_2 \|y\|_1.$$

By the second-order approximation in the low-loss region, choose $\eta_1 \in (0, \eta_0)$ small enough so that, whenever $\|y\|_1 \leq \eta_1$, the second-order term can be absorbed into the contraction margin under $\|\cdot\|_M$. That is, there exists $r_1 \in (r, 1)$ such that

$$\|My\|_M + C\|y\|_1^2 \leq r_1 \|y\|_M.$$

Thus, on the event $\{\|Y_m^N\|_1 \leq \eta_1\}$,

$$\mathbb{E} \left[\|Y_{m+1}^N\|_M \mid \mathcal{F}_m \right] \leq r_1 \|Y_m^N\|_M + Cr_N.$$

Let

$$\tau_N := \inf\{m \geq 0 : \|Y_m^N\|_1 > \eta_1\}.$$

Iterating the stopped recursion pathwise gives a stable linear filter driven by the one-step residuals and the martingale differences. More explicitly, with

$$Z_{m+1}^N := Y_{m+1}^N - \mathbb{E}(Y_{m+1}^N \mid \mathcal{F}_m),$$

one obtains, up to an equivalent-norm constant,

$$\max_{0 \leq m \leq T_N} \|Y_{m \wedge T_N}^N\|_M \leq C\|Y_0^N\|_M + Cr_N + C \max_{0 \leq m \leq T_N} \left\| \sum_{j=0}^{m-1} r_1^{m-1-j} Z_{j+1}^N \right\|_1.$$

The last term is a stable martingale transform. By the conditional second-moment estimate and Doob's inequality for this geometrically weighted transform,

$$\mathbb{E} \max_{0 \leq m \leq T_N} \left\| \sum_{j=0}^{m-1} r_1^{m-1-j} Z_{j+1}^N \right\|_1^2 \leq C \frac{T_N}{N}.$$

Consequently,

$$\mathbb{P}(\tau_N \leq T_N) \leq \frac{C}{\eta_1^2} \left(\|Y_0^N\|_1^2 + r_N^2 + \frac{T_N}{N} \right) \rightarrow 0,$$

using $\|Y_0^N\|_1 \rightarrow 0$, $r_N \rightarrow 0$, and $T_N = o(N)$. On $\{\tau_N > T_N\}$, all $m \leq T_N$ remain in the low-loss contraction region, and

$$\sum_{k=1}^K \pi_k Y_m^{k,N} \leq a_4 \|Y_m^N\|_1 \leq C \|Y_m^N\|_M.$$

The same maximal stopped-process estimate and Markov's inequality give

$$\mathbb{P} \left(\exists m \leq T_N : \sum_{k=1}^K \pi_k Y_m^{k,N} \geq \varepsilon, \tau_N > T_N \right) \leq \frac{C}{\varepsilon^2} \left(\|Y_0^N\|_1^2 + r_N^2 + \frac{T_N}{N} \right) \rightarrow 0.$$

Thus

$$\mathbb{P}(C_N(\varepsilon)) \leq \mathbb{P}(\tau_N \leq T_N) + \mathbb{P} \left(\exists m \leq T_N : \sum_{k=1}^K \pi_k Y_m^{k,N} \geq \varepsilon, \tau_N > T_N \right) \rightarrow 0.$$

This proves the subcritical conclusion.

We next prove the supercritical case. Suppose that $R_{\text{eff}} > 1$. Choose three constants

$$1 < r_- < r_+ < \bar{r} < R_{\text{eff}}.$$

This strict separation is the margin needed in the martingale step below. By the low-loss second-order approximation and $u^\top M = R_{\text{eff}} u^\top$, there exists $\eta_2 \in (0, \eta_0)$ such that, whenever $\|Y_m^N\|_1 \leq \eta_2$,

$$\mathbb{E} \left[u^\top Y_{m+1}^N \mid \mathcal{F}_m \right] \geq \bar{r} u^\top Y_m^N - Cr_N.$$

Indeed, the quadratic term is absorbed by the gap between R_{eff} and $\bar{r}$ after shrinking η_2 . Write

$$A_m^N := u^\top Y_m^N.$$

By the nondegenerate projection of the initial seed on the Perron–Frobenius principal direction, there exists $b_0 > 0$ such that

$$A_0^N \geq b_0 \|Y_0^N\|_1$$

with probability tending to one. The rest of the argument is restricted to this high-probability event.

The positive-cone equivalence gives

$$a_1 \|y\|_1 \leq u^\top y, \quad a_3 u^\top y \leq \sum_{k=1}^K \pi_k y_k.$$

Choose

$$\varepsilon_* := \frac{1}{2} a_1 a_3 \eta_2.$$

Fix $\varepsilon \in (0, \varepsilon_*)$ and define the Perron-projection threshold

$$B_\varepsilon := \frac{\varepsilon}{a_3}.$$

If $A_m^N < B_\varepsilon$, then

$$\|Y_m^N\|_1 \leq \frac{A_m^N}{a_1} < \frac{\varepsilon}{a_1 a_3} < \eta_2,$$

so the process remains inside the low-loss approximation region before the threshold is hit. Define

$$\sigma_N := \inf\{m \geq 0 : A_m^N \geq B_\varepsilon\}.$$

Let

$$m_N := \left\lceil \frac{\log(B_\varepsilon/A_0^N)}{\log r_-} \right\rceil.$$

Since $A_0^N \asymp \|Y_0^N\|_1$,

$$m_N = O\left(\log \frac{1}{\|Y_0^N\|_1}\right).$$

The observation window is compatible with this logarithmic hitting time. Indeed, choose $r_- < R_{\text{eff}}$ sufficiently close to R_{eff} so that, for all large N ,

$$m_N \leq (1 + \delta/2) \frac{\log(\varepsilon/\|Y_0^N\|_1)}{\log R_{\text{eff}}} \leq T_N.$$

The quadratic window is also compatible: before σ_N one has $A_m^N < B_\varepsilon$, hence $\|Y_m^N\|_1 \leq A_m^N/a_1 < \eta_2$. Thus the second-order approximation remains valid throughout the entire logarithmic growth window. The remainder compatibility condition implies $r_N = o(A_0^N)$. On the inductive growth event before σ_N , we have $A_m^N \geq A_0^N$. Hence, for all sufficiently large N and all $m < m_N \wedge \sigma_N$,

$$Cr_N \leq \frac{\bar{r} - r_+}{2} A_m^N.$$

Consequently,

$$\mathbb{E}[A_{m+1}^N \mid \mathcal{F}_m] \geq r_+ A_m^N, \quad m < m_N \wedge \sigma_N.$$

The gap between r_+ and r_- is now available for concentration.

Let

$$\zeta_{m+1}^N := A_{m+1}^N - \mathbb{E}(A_{m+1}^N \mid \mathcal{F}_m).$$

By the conditional second-moment assumption and boundedness of u ,

$$\mathbb{E}[|\zeta_{m+1}^N|^2 \mid \mathcal{F}_m] \leq \frac{C}{N}$$

on $m < \sigma_N$. If at some step

$$A_{m+1}^N < r_- A_m^N,$$

then, on the event where the conditional mean is at least $r_+ A_m^N$,

$$\zeta_{m+1}^N \leq -(r_+ - r_-) A_m^N.$$

Chebyshev's inequality therefore gives

$$\mathbb{P}\left(A_{m+1}^N < r_- A_m^N \mid \mathcal{F}_m\right) \leq \frac{C}{N(A_m^N)^2}.$$

On the inductive growth event,

$$A_m^N \geq r_-^m A_0^N \geq A_0^N.$$

A union bound over $m \leq m_N$ gives

$$\mathbb{P}\left(\exists m < m_N \wedge \sigma_N : A_{m+1}^N < r_- A_m^N\right) \leq \frac{C m_N}{N(A_0^N)^2} \leq \frac{C \log(1/\|Y_0^N\|_1)}{N\|Y_0^N\|_1^2} \longrightarrow 0.$$

Thus

$$\mathbb{P}\left(A_{m+1}^N \geq r_- A_m^N \text{ for all } m < m_N \wedge \sigma_N\right) \longrightarrow 1.$$

On this high-probability event,

$$A_m^N \geq r_-^m A_0^N, \quad 0 \leq m \leq m_N \wedge \sigma_N.$$

By the definition of m_N ,

$$r_-^{m_N} A_0^N \geq B_\varepsilon.$$

Therefore

$$\mathbb{P}(\sigma_N \leq m_N) \longrightarrow 1.$$

Because r_- can be chosen strictly below R_{eff} and arbitrarily close to it, the observation-window assumption

$$T_N \geq (1 + \delta) \frac{\log(\varepsilon/\|Y_0^N\|_1)}{\log R_{\text{eff}}}$$

allows us to choose r_- close enough to R_{eff} that $m_N \leq T_N$ for all sufficiently large N . Hence

$$\mathbb{P}(\sigma_N \leq T_N) \longrightarrow 1.$$

At time σ_N , the definition of B_ε and the positive-cone equivalence give

$$\sum_{k=1}^K \pi_k Y_{\sigma_N}^{k,N} \geq a_3 A_{\sigma_N}^N \geq a_3 B_\varepsilon = \varepsilon.$$

Since $Y_m^{k,N} = \Delta L_m^{k,N}$, this is the triggering condition for $C_N(\varepsilon)$. Therefore

$$\mathbb{P}(C_N(\varepsilon)) \geq \mathbb{P}(\sigma_N \leq T_N) \longrightarrow 1.$$

The supercritical conclusion follows.

Combining the two cases, the growth factor of the low-loss linearized system is

$$\rho(M) = \rho(\Gamma) \frac{\lambda_0 \Delta t}{\mathfrak{d}_\kappa}.$$

When this quantity is below 1, new losses decay geometrically in all directions. When it is above 1, new losses grow geometrically along the Perron–Frobenius principal direction and reach a fixed cascade threshold within logarithmic time. The nonlinear terms are second-order small quantities and only change a finite-width window around the critical region. Hence the local critical value is

$$R^* = 1.$$

The proof is complete. $\square$

Proposition SM3.14 (Continuous dependence of loss processes on the contagion matrix). *Fix $T > 0$. Let $\Gamma, \Gamma' \in \mathbb{R}_+^{K \times K}$ be two contagion matrices, and assume that they satisfy the uniform boundedness condition*

$$\|\Gamma\|_\infty \vee \|\Gamma'\|_\infty \leq \Lambda_\Gamma.$$

The two McKean–Vlasov systems have the same initial distribution, the same drift functions, the same diffusion coefficients, the same control process, and the same kill intensity. Assume also that b_k is Lipschitz in the state variable and measure variables, that λ_k is Lipschitz in the state variable, and that

$$0 \leq \lambda_k(x, a) \leq \Lambda_\lambda.$$

Let

$$(\mu_t^k(\Gamma), L_t^k(\Gamma))_{0 \leq t \leq T}, \quad (\mu_t^k(\Gamma'), L_t^k(\Gamma'))_{0 \leq t \leq T}$$

denote the limiting surviving measures and loss processes under contagion matrices Γ and Γ' , respectively. If both limiting systems are well posed, then there exists a constant

$$C_T = C(T, K, L_b, L_\lambda, \Lambda_\lambda, \Lambda_\Gamma) < \infty,$$

such that for each $k = 1, \dots, K$,

$$\mathbb{E} \sup_{0 \leq t \leq T} |L_t^k(\Gamma) - L_t^k(\Gamma')| \leq C_T \|\Gamma - \Gamma'\|_\infty.$$

Equivalently, conditional on the common-noise path, the same estimate holds almost surely for the conditional loss processes. In particular, if the main text uses the same constant Λ to uniformly dominate the upper bound on the kill intensity and the common upper bound on the contagion matrices, then the constant may be written as $C(T, \Lambda)$. Thus the limiting loss process is Lipschitz continuous with respect to the contagion matrix in L^1 , and pathwise after conditioning on the common noise.

Proof. Construct the two limiting systems on the same probability space. For each type k , let the representative particle under contagion matrix Γ and the representative particle under contagion matrix Γ' use the same initial state, the same idiosyncratic Brownian motion, the same common Brownian motion, and the same Poisson random measure. Write

$$X_t^{k, \Gamma} := X_t^k(\Gamma), \quad X_t^{k, \Gamma'} := X_t^k(\Gamma'),$$

and write the corresponding survival indicators as

$$\xi_t^{k, \Gamma} := \mathbf{1}_{\{\tau^{k, \Gamma} > t\}}, \quad \xi_t^{k, \Gamma'} := \mathbf{1}_{\{\tau^{k, \Gamma'} > t\}}.$$

The loss processes of the two systems satisfy

$$L_t^k(\Gamma) = 1 - \mathbb{E} \left[\xi_t^{k, \Gamma} \mid \mathcal{F}_t^{W^0} \right], \quad L_t^k(\Gamma') = 1 - \mathbb{E} \left[\xi_t^{k, \Gamma'} \mid \mathcal{F}_t^{W^0} \right].$$

Below, write

$$\delta_\Gamma := \|\Gamma - \Gamma'\|_\infty.$$

Let

$$e_t^k := X_t^{k,\Gamma} - X_t^{k,\Gamma'}.$$

Since the two systems use the same Brownian motions, the diffusion terms cancel in the difference. Subtracting the state equations gives

$$\begin{aligned} e_t^k &= \int_0^t \left[b_k(X_s^{k,\Gamma}, \mu_s^1(\Gamma), \dots, \mu_s^K(\Gamma), a_s^k) - b_k(X_s^{k,\Gamma'}, \mu_s^1(\Gamma'), \dots, \mu_s^K(\Gamma'), a_s^k) \right] ds \\ &\quad - \sum_{\ell=1}^K \Gamma_{k\ell} (L_t^\ell(\Gamma) - L_t^\ell(\Gamma')) - \sum_{\ell=1}^K (\Gamma_{k\ell} - \Gamma'_{k\ell}) L_t^\ell(\Gamma'). \end{aligned}$$

Here we have used $L_0^\ell(\Gamma) = L_0^\ell(\Gamma') = 0$ and the fact that Γ, Γ' are constant in time. Since $0 \leq L_t^\ell(\Gamma') \leq 1$,

$$\left| \sum_{\ell=1}^K (\Gamma_{k\ell} - \Gamma'_{k\ell}) L_t^\ell(\Gamma') \right| \leq K \delta_\Gamma.$$

By the Lipschitz property of b_k and $\|\Gamma\|_\infty \leq \Lambda_\Gamma$,

$$\begin{aligned} \sup_{r \leq t} |e_r^k| &\leq C \int_0^t \left(|e_s^k| + \sum_{\ell=1}^K d_{\text{BL}}(\mu_s^\ell(\Gamma), \mu_s^\ell(\Gamma')) \right) ds \\ &\quad + C \sum_{\ell=1}^K \sup_{r \leq t} |L_r^\ell(\Gamma) - L_r^\ell(\Gamma')| + C \delta_\Gamma. \end{aligned}$$

Given the common noise, denote conditional expectation by

$$\mathbb{E}^0[\cdot] := \mathbb{E}[\cdot \mid \mathcal{F}_T^{W^0}].$$

Define

$$u_k(t) := \mathbb{E}^0 \sup_{r \leq t} |e_r^k|,$$

$$m_k(t) := \sup_{r \leq t} d_{\text{BL}}(\mu_r^k(\Gamma), \mu_r^k(\Gamma')),$$

and

$$v_k(t) := \sup_{r \leq t} |L_r^k(\Gamma) - L_r^k(\Gamma')|.$$

Then

$$u_k(t) \leq C \int_0^t \left(u_k(s) + \sum_{\ell=1}^K m_\ell(s) \right) ds + C \sum_{\ell=1}^K v_\ell(t) + C \delta_\Gamma.$$

We next control the measure error. For any test function f satisfying $\|f\|_\infty \leq 1$ and $\text{Lip}(f) \leq 1$,

$$\begin{aligned} &\left| \langle f, \mu_t^k(\Gamma) - \mu_t^k(\Gamma') \rangle \right| \\ &\leq \mathbb{E}^0 \left| \xi_t^{k,\Gamma} f(X_t^{k,\Gamma}) - \xi_t^{k,\Gamma'} f(X_t^{k,\Gamma'}) \right| \\ &\leq \mathbb{E}^0 |X_t^{k,\Gamma} - X_t^{k,\Gamma'}| + \mathbb{E}^0 |\xi_t^{k,\Gamma} - \xi_t^{k,\Gamma'}|. \end{aligned}$$

Synchronous Poisson thinning gives

$$\mathbb{E}^0 \sup_{r \leq t} |\xi_r^{k,\Gamma} - \xi_r^{k,\Gamma'}| \leq \mathbb{E}^0 \int_0^t \left| \lambda_k(X_s^{k,\Gamma}, a_s^k) - \lambda_k(X_s^{k,\Gamma'}, a_s^k) \right| ds.$$

By the Lipschitz property of λ_k in the state variable,

$$\mathbb{E}^0 \sup_{r \leq t} |\xi_r^{k,\Gamma} - \xi_r^{k,\Gamma'}| \leq L_\lambda \int_0^t u_k(s) ds.$$

Taking the supremum over test functions, and then the supremum over $r \leq t$, gives

$$m_k(t) \leq u_k(t) + C \int_0^t u_k(s) ds.$$

Similarly, by the conditional expectation representation of the loss process,

$$\begin{aligned} v_k(t) &= \sup_{r \leq t} \left| \mathbb{E}^0 \left[\xi_r^{k,\Gamma'} - \xi_r^{k,\Gamma} \right] \right| \\ &\leq \mathbb{E}^0 \sup_{r \leq t} |\xi_r^{k,\Gamma'} - \xi_r^{k,\Gamma}| \\ &\leq C \int_0^t u_k(s) ds. \end{aligned}$$

Let

$$U(t) := \sum_{k=1}^K u_k(t), \quad M(t) := \sum_{k=1}^K m_k(t), \quad V(t) := \sum_{k=1}^K v_k(t).$$

Combining the estimates above gives

$$U(t) \leq C\delta_\Gamma + C \int_0^t (U(s) + M(s)) ds + CV(t),$$

and

$$M(t) \leq CU(t) + C \int_0^t U(s) ds, \quad V(t) \leq C \int_0^t U(s) ds.$$

Substituting the latter two estimates into the estimate for $U(t)$ and absorbing the double integral into the constant gives

$$U(t) \leq C\delta_\Gamma + C \int_0^t U(s) ds, \quad 0 \leq t \leq T.$$

By Gronwall's inequality,

$$U(t) \leq C_T \delta_\Gamma, \quad 0 \leq t \leq T.$$

Substituting this bound into $V(t) \leq C \int_0^t U(s) ds$ gives

$$V(T) \leq C_T \delta_\Gamma.$$

Therefore, for each $k = 1, \dots, K$, conditional on the common-noise path,

$$\sup_{0 \leq t \leq T} \left| L_t^k(\Gamma) - L_t^k(\Gamma') \right| = v_k(T) \leq V(T) \leq C_T \|\Gamma - \Gamma'\|_\infty.$$

Taking expectations gives the stated L^1 estimate. The proof is complete. $\square$

Remark SM3.15 (Case without common noise). If the model has no common noise, the conditional expectation $\mathbb{E}^0[\cdot]$ in the proof above is replaced directly by ordinary expectation, and the conclusion remains unchanged.

SM4 Proofs for controlled contagion systems, the killing limit, and the information value of local pressure

Assumption SM4.1 (Regularity conditions for the two-population controlled contagion system). Let $T > 0$. The two type shares π_1, π_2 are fixed strictly positive constants. Set

$$\mathbf{E} = \mathbb{R} \cup \{\partial\},$$

where ∂ denotes the cemetery state. Equip $\mathbf{E}$ with a metric d_∂ that agrees with the Euclidean distance on $\mathbb{R}$ and satisfies

$$d_\partial(x, \partial)^2 \leq C_\partial(1 + |x|^2), \quad x \in \mathbb{R}.$$

For $k = 1, 2$, let

$$\mu^k \in \mathcal{P}_2(\mathbf{E}), \quad L^k(\mu^k) := \mu^k(\{\partial\}),$$

and write

$$\langle f, \mu^k \rangle_\circ := \int_{\mathbb{R}} f(x) \mu^k(dx),$$

so that the integral is taken only over the alive-state set $\mathbb{R}$.

Assume that the drift function

$$b_k : \mathbb{R} \times \mathcal{P}_2(\mathbf{E})^2 \rightarrow \mathbb{R}$$

is uniformly Lipschitz in (x, μ^1, μ^2) and satisfies a linear growth condition. Assume that the diffusion coefficients

$$\sigma_k > 0, \quad \sigma_0 \geq 0$$

are constants, where $\sigma_0 dW_t^0$ denotes the common-noise term. The strict positivity of the idiosyncratic coefficients is the nondegeneracy condition used in the comparison theorem for the killed HJB below. The contagion matrix

$$\Gamma = (\Gamma_{k\ell})_{1 \leq k, \ell \leq 2}$$

is nonnegative and bounded.

The kill intensity

$$\lambda_k : \mathbb{R} \times [0, \infty) \rightarrow [0, \infty)$$

is continuous, and there exists a constant $\Lambda < \infty$ such that

$$0 \leq \lambda_k(x, a) \leq \Lambda, \quad x \in \mathbb{R}, \quad a \geq 0.$$

Moreover, λ_k is uniformly Lipschitz in x and locally Lipschitz in the control variable a . Extend the kill intensity to the cemetery state by setting

$$\lambda_k(\partial, a) = 0.$$

The terminal loss function

$$\Phi : [0, 1]^2 \rightarrow \mathbb{R}$$

is bounded and Lipschitz. The control cost function

$$c_k : [0, \infty) \rightarrow [0, \infty)$$

is continuous, convex, lower semicontinuous, and satisfies the superlinear growth condition

$$\lim_{a \rightarrow \infty} \frac{c_k(a)}{a} = +\infty.$$

In addition, assume that the controlled martingale problem is well posed for every progressively measurable compact-valued control. The dynamic-programming and viscosity framework follows the standard infinite-dimensional control approach of Fabbri–Gozzi–Swiech [41]. The comparison-principle argument for the present killed Wasserstein HJB is formulated after the decomposition $\mu^k = \nu^k + L^k \delta_\partial$, with alive finite measures and cemetery masses treated as separate coordinates. It combines the Wasserstein smooth-gauge method of Cosso–Gozzi–Kharroubi–Pham–Rosetolato [43] for the alive-measure variables with the paper-specific finite-dimensional estimates for the cemetery-state killing jump.

For the existence statement, admissible controls are taken in the relaxed sense. A relaxed control is a progressively measurable probability-measure-valued process

$$\vartheta_t \in \mathcal{P}([0, \bar{a}]^2),$$

and strict controls correspond to Dirac relaxed controls $\vartheta_t = \delta_{a_t}$. Coefficients and costs under ϑ_t are obtained by averaging the corresponding strict-control coefficients over $[0, \bar{a}]^2$. Because the action set is compact and the coefficients are continuous in the action, this relaxed admissible class is compact under weak convergence.

SM4.1 Comparison principle for the killed two-population HJB

This subsection proves the comparison-principle interface for the killed two-population HJB. The proof is written on the decomposed state space

$$\mathcal{D}_2 := \left\{ (\nu^1, \nu^2, L^1, L^2) : \nu^k \in \mathcal{M}_{\leq 1,2}(\mathbb{R}), L^k \in [0, 1], |\nu^k| + L^k = 1 \right\},$$

where $\mathcal{M}_{\leq 1,2}(\mathbb{R})$ denotes finite sub-probability measures on $\mathbb{R}$ with finite second moment. The Wasserstein smooth-gauge/Ishii template is used for the alive-measure coordinates ν^k , and the cemetery masses L^k are handled by the finite-dimensional doubled-variable penalty $\beta|L - M|^2/2$. The only term outside the standard alive-measure template is the mass transfer from ν^k to L^k ; Lemma SM4.5 verifies that this killing jump is a bounded Lipschitz first-order perturbation compatible with the doubled-variable estimate.

Throughout this subsection the control set is the compact interval $[0, \bar{a}]^2$. Let

$$\mathbf{E} = \mathbb{R} \cup \{\partial\}$$

be endowed with an auxiliary metric d_∂ that agrees with the Euclidean metric on compact alive-state sets and separates the cemetery point from $\mathbb{R}$. This metric is used only to identify (ν^k, L^k) with the killed probability measure $\nu^k + L^k \delta_\partial$ and to define localization estimates for unmatched alive/cemetery mass; the comparison argument itself is carried out on $\mathcal{D}_2$. On each localization interval $[-R, R]$ there is $c_R > 0$ such that $d_\partial(x, \partial) \geq c_R$ for $x \in [-R, R]$. The localization is removed by the usual coercive second-moment penalty in the Wasserstein comparison argument. Equivalently, one may work directly with a metric such as $d_\partial(x, y) = |x - y|$ on $\mathbb{R}$ and $d_\partial(x, \partial) = 1 + |x|$. We extend the kill intensity by $\lambda_k(\partial, a) = 0$. The assumptions used below are

$$0 \leq \lambda_k(x, a) \leq \Lambda, \quad |\lambda_k(x, a) - \lambda_k(y, a)| \leq L_x |x - y|, \quad |\lambda_k(x, a) - \lambda_k(x, \tilde{a})| \leq L_a |a - \tilde{a}|,$$

for $x, y \in \mathbb{R}$ and $a, \tilde{a} \in [0, \bar{a}]$. After the extension $\lambda_k(\partial, a) = 0$, the localized Lipschitz constant of $\lambda_k(\cdot, a)$ on $[-R, R] \cup \{\partial\}$ is bounded by $C_R(L_x + \Lambda)$, because the alive–cemetery jump satisfies $|\lambda_k(x, a) - \lambda_k(\partial, a)| \leq \Lambda$ and $d_\partial(x, \partial) \geq c_R$ on $[-R, R]$. The drift is Lipschitz in the alive state, in the measure arguments, and in the loss variables, with at most linear growth. The diffusion

coefficients are constant, and in the comparison statement below the alive idiosyncratic coefficients satisfy $\sigma_k > 0$, $k = 1, 2$. The terminal cost is bounded and uniformly continuous, and the running control costs are continuous on the compact action set.

For $\nu \in \mathcal{M}_{\leq 1}(\mathbb{R})$, define the augmented probability measure

$$\bar{\nu} := \nu + (1 - |\nu|)\delta_{\partial}, \quad |\nu| := \nu(\mathbb{R}),$$

and define

$$\mathbf{W}_p(\nu, \omega) := W_p(\bar{\nu}, \bar{\omega}; d_{\partial}), \quad p = 1, 2.$$

On each localization interval, the indicator of the alive set is Lipschitz after extension to ∂ , hence

$$||\nu| - |\omega|| \leq C_R \mathbf{W}_1(\nu, \omega) \leq C_R \mathbf{W}_2(\nu, \omega).$$

All constants below are uniform on fixed localizations; the coercive penalty lets $R \rightarrow \infty$ at the end.

Lemma SM4.2 (Bounded Lipschitz control of the killing sink). *Let $p : \mathbb{R} \rightarrow \mathbb{R}$ be bounded and Lipschitz, with $\|p\|_{\infty} \leq B$ and $\text{Lip}(p) \leq L_p$. Extend p to $\mathbf{E}$ by setting $p(\partial) = 0$. Then, for every $\nu, \omega \in \mathcal{M}_{\leq 1}(\mathbb{R})$ and every $a, \tilde{a} \in [0, \bar{a}]$,*

$$\begin{aligned} & \left| \int_{\mathbb{R}} \lambda_k(x, a) p(x) \nu(dx) - \int_{\mathbb{R}} \lambda_k(y, \tilde{a}) p(y) \omega(dy) \right| \\ & \leq C_R (\Lambda L_p + L_x B + \Lambda B) \mathbf{W}_1(\nu, \omega) + L_a B |a - \tilde{a}|. \end{aligned}$$

Equivalently, the cemetery contribution may be written separately as

$$\begin{aligned} & \left| \int_{\mathbb{R}} \lambda_k(x, a) p(x) \nu(dx) - \int_{\mathbb{R}} \lambda_k(y, \tilde{a}) p(y) \omega(dy) \right| \\ & \leq C_R (\Lambda L_p + L_x B) \mathbf{W}_1(\nu, \omega) + C_R \Lambda B ||\nu| - |\omega|| + L_a B |a - \tilde{a}|. \end{aligned}$$

Proof. Set $f_a(x) = \lambda_k(x, a) p(x)$ for $x \in \mathbb{R}$ and $f_a(\partial) = 0$. On $\mathbf{E}$,

$$\|f_a\|_{\infty} \leq \Lambda B.$$

For $x, y \in \mathbb{R}$,

$$\begin{aligned} |f_a(x) - f_a(y)| & \leq \lambda_k(x, a) |p(x) - p(y)| + |p(y)| |\lambda_k(x, a) - \lambda_k(y, a)| \\ & \leq (\Lambda L_p + L_x B) |x - y|. \end{aligned}$$

For $x \in [-R, R]$, the jump from x to ∂ is bounded by ΛB , and the cemetery point is separated from the alive set; this gives

$$\text{Lip}_{d_{\partial}}(f_a) \leq C_R (\Lambda L_p + L_x B + \Lambda B).$$

Therefore the Kantorovich–Rubinstein duality gives

$$\left| \int f_a d\bar{\nu} - \int f_a d\bar{\omega} \right| \leq C_R (\Lambda L_p + L_x B + \Lambda B) \mathbf{W}_1(\nu, \omega).$$

Since $f_a(\partial) = 0$, these are exactly the alive-state integrals. The control difference is estimated by

$$\left| \int_{\mathbb{R}} [\lambda_k(y, a) - \lambda_k(y, \tilde{a})] p(y) \omega(dy) \right| \leq L_a B |a - \tilde{a}|.$$

Combining the last two inequalities proves the first bound. The second displayed bound is the same estimate with the cemetery jump contribution separated into the unmatched-mass term. $\square$

Lemma SM4.3 (Aggregate kill rate is Wasserstein-Lipschitz). *Let*

$$\Lambda_k(\nu, a) := \int_{\mathbb{R}} \lambda_k(x, a) \nu(dx), \quad \nu \in \mathcal{M}_{\leq 1}(\mathbb{R}), \quad a \in [0, \bar{a}].$$

Then

$$|\Lambda_k(\nu, a) - \Lambda_k(\omega, \tilde{a})| \leq C_R(\Lambda + L_x)W_1(\nu, \omega) + L_a|a - \tilde{a}|.$$

In particular, for the same control a ,

$$|\Lambda_k(\nu, a) - \Lambda_k(\omega, a)| \leq C_R(\Lambda + L_x)W_2(\nu, \omega).$$

Proof. Apply Lemma SM4.2 with $p \equiv 1$ on $\mathbb{R}$, extended by $p(\partial) = 0$. Then $B = 1$, $L_p = 0$, and the result follows from $W_1 \leq W_2$ for the augmented probability measures. $\square$

Lemma SM4.4 (State decomposition and reformulated killed HJB). *Every $\mu^k \in \mathcal{P}_2(\mathbf{E})$ has the unique decomposition*

$$\mu^k = \nu^k + L^k \delta_{\partial}, \quad \nu^k := \mu^k|_{\mathbb{R}} \in \mathcal{M}_{\leq 1}(\mathbb{R}), \quad L^k := \mu^k(\{\partial\}) \in [0, 1],$$

with $|\nu^k| + L^k = 1$. Let

$$v(t, \nu^1, \nu^2, L^1, L^2) := V(t, \nu^1 + L^1 \delta_{\partial}, \nu^2 + L^2 \delta_{\partial}).$$

At smooth points, after taking a local extension of v to a neighborhood in the positive cone of finite alive measures and cemetery masses,

$$\frac{\delta V}{\delta \mu^k}(x) = \frac{\delta v}{\delta \nu^k}(x), \quad x \in \mathbb{R}, \quad \frac{\delta V}{\delta \mu^k}(\partial) = \partial_{L^k} v.$$

The difference

$$\partial_{L^k} v - \frac{\delta v}{\delta \nu^k}(x)$$

is independent of the additive normalization of the linear functional derivative. Consequently the killing term becomes

$$\begin{aligned} & \int_{\mathbb{R}} \lambda_k(x, a^k) \left[\frac{\delta V}{\delta \mu^k}(\partial) - \frac{\delta V}{\delta \mu^k}(x) \right] \mu^k(dx) \\ &= \Lambda_k(\nu^k, a^k) \partial_{L^k} v - \int_{\mathbb{R}} \lambda_k(x, a^k) \frac{\delta v}{\delta \nu^k}(x) \nu^k(dx). \end{aligned}$$

Proof. The decomposition follows from the disjoint union $\mathbf{E} = \mathbb{R} \cup \{\partial\}$. A signed perturbation supported on $\mathbb{R}$ changes only the alive finite measure ν^k , and therefore gives the identity of the alive functional derivatives. A perturbation in the cemetery atom changes only the scalar variable L^k , and therefore gives the identity at ∂ . Since linear functional derivatives on probability measures are defined up to additive constants, the invariant object is the difference between the cemetery derivative and the alive derivative. This is exactly the first variation along the jump direction $-\delta_x + \delta_{\partial}$, and substituting it into the original generator gives the displayed decomposition. $\square$

Lemma SM4.5 (Absorption of the killing term in the doubled-variable argument). *Let u be a bounded uniformly continuous viscosity subsolution and w a bounded uniformly continuous viscosity supersolution of the decomposed killed HJB on the decomposed state space. Let*

$$(t_{\varepsilon}, s_{\varepsilon}, \nu_{\varepsilon}, \omega_{\varepsilon}, L_{\varepsilon}, M_{\varepsilon})$$

be a maximum point of the doubled functional

$$u(t, \nu, L) - w(s, \omega, M) - \frac{|t - s|^2}{2\alpha} - \sum_{k=1}^2 \Theta_\varepsilon^k(\nu^k, \omega^k) - \frac{\beta}{2}|L - M|^2 - \delta\mathcal{R}(\nu, \omega),$$

where Θ_ε^k is the Wasserstein smooth gauge used in the comparison proof and $\mathcal{R}$ is a coercive second-moment penalty. Let p_ε^k and q_ε^k be the first-order measure jets generated by the gauge and the coercive penalty in the ν^k - and ω^k -variables, with signs corresponding to the subsolution and supersolution tests. If the same control $a \in [0, \bar{a}]^2$ is used on the two sides, then the killing contribution in the subsolution-minus-supersolution Hamiltonian difference satisfies, for fixed β ,

$$\lim_{\delta \downarrow 0} \limsup_{\varepsilon \downarrow 0} |I_{\text{kill}}(\varepsilon, \alpha, \beta, \delta)| \leq C \limsup_{\varepsilon \downarrow 0} \sum_{k=1}^2 \beta |L_\varepsilon^k - M_\varepsilon^k|^2.$$

Consequently,

$$\lim_{\beta \rightarrow \infty} \lim_{\delta \downarrow 0} \limsup_{\varepsilon \downarrow 0} |I_{\text{kill}}(\varepsilon, \alpha, \beta, \delta)| = 0.$$

Proof. By Lemma SM4.4, the decomposed killing operator is

$$\mathcal{K}^a v(\nu, L) = \sum_{k=1}^2 \Lambda_k(\nu^k, a^k) \partial_{L^k} v(\nu, L) - \sum_{k=1}^2 \int_{\mathbb{R}} \lambda_k(x, a^k) \frac{\delta v}{\delta \nu^k}(\nu, L)(x) \nu^k(dx).$$

At the doubled maximum point, the loss derivatives in the two tests are

$$r_\varepsilon^k := \beta(L_\varepsilon^k - M_\varepsilon^k).$$

The loss-drift part of the killing difference is therefore

$$I_k^L = [\Lambda_k(\nu_\varepsilon^k, a^k) - \Lambda_k(\omega_\varepsilon^k, a^k)] r_\varepsilon^k.$$

Lemma SM4.3 gives

$$|I_k^L| \leq C_R W_2(\nu_\varepsilon^k, \omega_\varepsilon^k) \beta |L_\varepsilon^k - M_\varepsilon^k|.$$

The doubled maximum-point estimate gives $W_2(\nu_\varepsilon^k, \omega_\varepsilon^k) \rightarrow 0$ as $\varepsilon \downarrow 0$ for fixed β , and $\beta |L_\varepsilon^k - M_\varepsilon^k|$ is bounded for fixed β because $L_\varepsilon^k, M_\varepsilon^k \in [0, 1]$. If the cemetery mass is treated as an independent scalar coordinate in the local extension, the same estimate also contains the harmless term $C\beta |L_\varepsilon^k - M_\varepsilon^k|^2$, which is retained in the final bound.

It remains to estimate the alive-measure sink

$$I_k^S = - \int_{\mathbb{R}} \lambda_k(x, a^k) p_\varepsilon^k(x) \nu_\varepsilon^k(dx) + \int_{\mathbb{R}} \lambda_k(y, a^k) q_\varepsilon^k(y) \omega_\varepsilon^k(dy).$$

For the unsmoothed squared-Wasserstein gauge, let π_ε^k be an optimal coupling of the augmented measures. On the alive-alive part of the coupling, the two first-order jets have the same paired representative

$$p_\varepsilon^k(x) = q_\varepsilon^k(y) = \frac{x - y}{\varepsilon}.$$

This is the gradient of the quadratic cost $d_\partial(x, y)^2/(2\varepsilon)$ evaluated on the support of the optimal coupling, where p_ε^k and q_ε^k denote the first-order Lions-derivative representatives in the subsolution

and supersolution tests, respectively. The cemetery contribution is represented by the corresponding metric derivative in the chosen cemetery metric after setting $\lambda_k(\partial, a^k) = 0$. Hence

$$\begin{aligned} |I_k^S| &\leq \int_{E \times E} |\lambda_k(y, a^k) - \lambda_k(x, a^k)| \frac{d\partial(x, y)}{\varepsilon} \pi_\varepsilon^k(dx, dy) \\ &\leq C_R(L_x + \Lambda) \frac{W_2^2(\bar{\nu}_\varepsilon^k, \bar{\omega}_\varepsilon^k)}{\varepsilon}. \end{aligned}$$

The smooth gauge of the Wasserstein Ishii lemma is obtained by regularizing the squared-Wasserstein penalty. More precisely, the smooth gauge Θ_ε^k is chosen so that, uniformly on bounded second-moment sets,

$$\Theta_\varepsilon^k(\nu, \omega) \geq \frac{W_2^2(\bar{\nu}, \bar{\omega})}{2\varepsilon} - \rho_\varepsilon^k, \quad \rho_\varepsilon^k \downarrow 0,$$

and its first-order Lions jets approximate the optimal-coupling jets of the unsmoothed quadratic penalty up to the same remainder. This is the smooth-gauge approximation mechanism in [43, Lemma 4.4 and Theorem 4.5]. Therefore the preceding bilinear estimate carries through with the regularization error absorbed into ρ_ε^k , and becomes

$$|I_k^S| \leq C_R \Theta_\varepsilon^k(\nu_\varepsilon^k, \omega_\varepsilon^k) + \rho_\varepsilon^k + C_R \delta (1 + m_2(\nu_\varepsilon^k) + m_2(\omega_\varepsilon^k)),$$

where $\rho_\varepsilon^k \rightarrow 0$ is the smooth-gauge approximation error. The maximum-point estimate for bounded uniformly continuous sub- and supersolutions gives

$$\Theta_\varepsilon^k(\nu_\varepsilon^k, \omega_\varepsilon^k) \rightarrow 0 \quad \text{as } \varepsilon \downarrow 0,$$

and the coercive penalty makes the last term vanish as $\delta \downarrow 0$. Therefore

$$\lim_{\delta \downarrow 0} \limsup_{\varepsilon \downarrow 0} |I_k^S| = 0.$$

Combining the loss-drift and alive-sink estimates gives the first displayed bound.

Finally, the finite-dimensional loss penalty and the uniform continuity of u, w imply

$$\lim_{\beta \rightarrow \infty} \limsup_{\varepsilon \downarrow 0} \sum_{k=1}^2 \beta |L_\varepsilon^k - M_\varepsilon^k|^2 = 0.$$

This is the standard finite-dimensional part of the doubling argument: if this quantity stayed bounded away from zero, replacing one loss vector by the other while keeping the other coordinates fixed would increase the doubled functional up to a modulus-of-continuity error smaller than the penalty gain. The second displayed limit follows. $\square$

Proposition SM4.6 (Wasserstein smooth-gauge/Ishii interface). *After the state decomposition in Lemma SM4.4, the killed two-population HJB reduces to a comparison problem on $\mathcal{D}_2$. On this decomposed space, the alive-measure part satisfies the structural hypotheses of the Wasserstein smooth-gauge comparison template, while the cemetery killing jump is controlled by Lemma SM4.5. The reference template is [43, Assumptions (A)–(D), Lemma 4.4, Theorems 4.5 and 5.1]. The Hamiltonian is proper, continuous in the state variables under $\mathbb{W}_2 \times \mathbb{W}_2 \times |\cdot|$, locally uniformly continuous in the first-order jets on bounded jet sets, and degenerate elliptic in the second-order jets. The diagonal idiosyncratic terms are nondegenerate in the alive directions because $\sigma_k > 0$, and the common-noise term has the usual positive-semidefinite product-measure covariance form. It is thus a reduction to the standard alive-measure smooth-gauge argument plus the finite-dimensional cemetery-mass estimate proved here.*

Proof. The cited comparison argument follows the viscosity doubling-variable framework of Crandall, Ishii, and Lions [42] in its Wasserstein smooth-gauge form, and assumes bounded continuous coefficients with Lipschitz dependence on the state and the measure variable, compactness of the control set, sufficient regularity of the diffusion coefficient for the smooth-gauge construction, bounded uniformly continuous terminal data, coercive localization on the unbounded second-moment state space, properness, local continuity of the Hamiltonian along admissible Wasserstein jets, and a degenerate-elliptic second-order structure. In the present model these conditions are verified after the alive-loss decomposition as follows. The Hamiltonian contains no zeroth-order dependence on the value itself, so properness follows directly. The control set is compact and the population-weighted running costs are continuous. The drift and contagion feedback are Lipschitz in the alive state, measure variables, and loss variables; Lemma SM4.3 gives the same continuity for the aggregate kill rates appearing in both the feedback drift and the loss drift. Lemma SM4.2 gives the continuity estimate for the alive-measure sink. The diffusion coefficients are constants, so the second-order part is continuous in the state and monotone in the second-order jet in the viscosity order. The idiosyncratic covariance matrices are positive definite in the alive directions under $\sigma_k > 0$, and the common-noise covariance matrix is positive semidefinite; hence the full second-order operator has the ellipticity structure required in the smooth-gauge/Ishii comparison proof. The only additional term relative to the template is the cemetery killing jump, and Lemma SM4.5 supplies the needed doubled-variable estimate for that term. The coercive second-moment penalty gives compactness of maximizing sequences before the localization radius is removed; the bounded uniformly continuous terminal condition preserves the terminal ordering, and the common-noise second-order term enters through the positive-semidefinite product covariance already allowed by the template. $\square$

Theorem SM4.7 (Killed HJB comparison principle). *Let U and W be bounded uniformly continuous functions on $[0, T] \times \mathcal{P}_2(\mathbf{E})^2$, and let their decomposed representatives u, w on $\mathcal{D}_2$ be viscosity sub- and supersolutions of the decomposed killed HJB associated with*

$$\partial_t V(t, \mu^1, \mu^2) + H(t, \mu^1, \mu^2, D_\mu V, D_{\mu\mu}^2 V) = 0.$$

The Hamiltonian and generator are those stated in the killed two-population HJB after the alive-cemetery decomposition. We use the backward terminal-time convention: subsolutions satisfy $\partial_t \phi + H[\phi] \geq 0$ at upper tests, and supersolutions satisfy $\partial_t \phi + H[\phi] \leq 0$ at lower tests. Assume the bounded Lipschitz killing, Lipschitz drift, constant diffusion, compact-control, nondegenerate alive idiosyncratic diffusion $\sigma_k > 0$, and bounded uniformly continuous terminal conditions stated above. If

$$U(T, \mu^1, \mu^2) \leq W(T, \mu^1, \mu^2), \quad (\mu^1, \mu^2) \in \mathcal{P}_2(\mathbf{E})^2,$$

then

$$U(t, \mu^1, \mu^2) \leq W(t, \mu^1, \mu^2), \quad (t, \mu^1, \mu^2) \in [0, T] \times \mathcal{P}_2(\mathbf{E})^2.$$

Proof. By Lemma SM4.4, it is enough to prove comparison for the decomposed functions

$$u(t, \nu, L) = U(t, \nu^1 + L^1 \delta_\partial, \nu^2 + L^2 \delta_\partial), \quad w(t, \nu, L) = W(t, \nu^1 + L^1 \delta_\partial, \nu^2 + L^2 \delta_\partial),$$

restricted to $|\nu^k| + L^k = 1$. Suppose by contradiction that $\sup(u - w) > 0$. For $\eta > 0$, set

$$u^\eta(t, \nu, L) = u(t, \nu, L) - \frac{\eta}{T - t}, \quad t < T.$$

For sufficiently small η , $\sup(u^\eta - w) > 0$, and the singular terminal barrier together with the terminal ordering forces the positive maximum to occur at an interior time.

Fix $\alpha, \varepsilon, \delta > 0$ and $\beta > 0$, and consider

$$\begin{aligned} \Psi(t, s, \nu, \omega, L, M) &= u^\eta(t, \nu, L) - w(s, \omega, M) - \frac{|t - s|^2}{2\alpha} - \sum_{k=1}^2 \Theta_\varepsilon^k(\nu^k, \omega^k) \\ &\quad - \frac{\beta}{2} |L - M|^2 - \delta \mathcal{R}(\nu, \omega). \end{aligned}$$

Let $(t_\varepsilon, s_\varepsilon, \nu_\varepsilon, \omega_\varepsilon, L_\varepsilon, M_\varepsilon)$ be a maximum point supplied by the Wasserstein smooth variational principle. The standard maximum-point estimates give

$$|t_\varepsilon - s_\varepsilon| \rightarrow 0, \quad \Theta_\varepsilon^k(\nu_\varepsilon^k, \omega_\varepsilon^k) \rightarrow 0, \quad W_2(\nu_\varepsilon^k, \omega_\varepsilon^k) \rightarrow 0,$$

for $k = 1, 2$, and

$$\lim_{\beta \rightarrow \infty} \limsup_{\varepsilon \downarrow 0} \beta |L_\varepsilon - M_\varepsilon|^2 = 0.$$

With the backward terminal-time convention that viscosity subsolutions of $\partial_t V + H = 0$ satisfy $\partial_t \phi + H[\phi] \geq 0$ at upper tests and supersolutions satisfy $\partial_t \phi + H[\phi] \leq 0$ at lower tests, the penalized function u^η yields the usual strict interior comparison inequality. Applying the subsolution inequality to the upper test for u^η , the supersolution inequality to the lower test for w , and using

$$\inf_a A(a) - \inf_a B(a) \leq \sup_{a \in [0, \bar{a}]^2} \{A(a) - B(a)\},$$

one obtains

$$\frac{\eta}{(T - t_\varepsilon)^2} \leq \Delta_{\text{time}} + \Delta_{\text{drift}} + \Delta_{\text{diff}} + \Delta_{\text{common}} + \Delta_{\text{kill}} + o(1),$$

where $o(1)$ denotes the smooth-gauge and localization errors.

The time penalty gives $\Delta_{\text{time}} \rightarrow 0$ as $\alpha \downarrow 0$. The drift and running-cost terms are controlled by their Lipschitz continuity and the maximum-point estimates. The diagonal diffusion and common-noise terms are controlled by the Wasserstein smooth-gauge/Ishii inequality; the positive-semidefinite second-order structure yields

$$\limsup_{\varepsilon \downarrow 0} (\Delta_{\text{diff}} + \Delta_{\text{common}}) \leq 0$$

after the coercive regularization is removed. Proposition SM4.6 verifies the structural conditions for these standard terms. Lemma SM4.5 gives

$$\lim_{\beta \rightarrow \infty} \lim_{\delta \downarrow 0} \limsup_{\varepsilon \downarrow 0} |\Delta_{\text{kill}}| = 0.$$

Taking the comparison limits in the standard order $\varepsilon \downarrow 0$, then $\delta \downarrow 0$, then $\beta \rightarrow \infty$, and finally $\alpha \downarrow 0$, gives

$$\frac{\eta}{(T - t_*)^2} \leq 0$$

for an interior limiting time $t_* < T$, a contradiction. Hence $u \leq w$ on the decomposed state space. Returning to $\mu^k = \nu^k + L^k \delta_\partial$ proves the comparison principle on $[0, T] \times \mathcal{P}_2(\mathbf{E})^2$. $\square$

Theorem SM4.8 (Existence of two-population optimal control and conditional common-noise HJB characterization). *Under Assumption SM4.1, consider the controlled McKean–Vlasov contagion system with $K = 2$. Given a control*

$$a_t = (a_t^1, a_t^2) \in [0, \infty)^2,$$

define the total kill intensity of type k by

$$\Lambda_k(\mu^k, a^k) := \int_{\mathbb{R}} \lambda_k(x, a^k) \mu^k(dx).$$

The lifted measure state $(\mu_t^1, \mu_t^2) \in \mathcal{P}_2(\mathbb{E})^2$ satisfies the following weak form: for any $\varphi \in C_b^2(\mathbb{R})$,

$$\begin{aligned} d\langle \varphi, \mu_t^k \rangle_\circ &= \left\{ \int_{\mathbb{R}} \left[b_k(x, \mu_t^1, \mu_t^2) - \sum_{\ell=1}^2 \Gamma_{k\ell} \Lambda_\ell(\mu_t^\ell, a_t^\ell) \right] \varphi'(x) \mu_t^k(dx) \right. \\ &\quad \left. + \frac{1}{2} \sigma_k^2 \int_{\mathbb{R}} \varphi''(x) \mu_t^k(dx) - \int_{\mathbb{R}} \lambda_k(x, a_t^k) \varphi(x) \mu_t^k(dx) \right\} dt \\ &\quad + \sigma_0 \int_{\mathbb{R}} \varphi'(x) \mu_t^k(dx) dW_t^0. \end{aligned} \quad (20)$$

The loss process is given by the cemetery mass:

$$L_t^k = \mu_t^k(\{\partial\}),$$

and

$$dL_t^k = \Lambda_k(\mu_t^k, a_t^k) dt. \quad (21)$$

The central controller's objective for a strict control is

$$J(t, \mu^1, \mu^2; a) := \mathbb{E} \left[\Phi(L_T^1, L_T^2) + \int_t^T (\pi_1 c_1(a_s^1) + \pi_2 c_2(a_s^2)) ds \right].$$

For a relaxed control ϑ , the notation $J(t, \mu^1, \mu^2; \vartheta)$ denotes the same objective with the drift, killing intensity, and population-weighted running cost averaged with respect to $\vartheta_s(da)$. The value function is

$$V(t, \mu^1, \mu^2) := \inf_{\vartheta} J(t, \mu^1, \mu^2; \vartheta). \quad (22)$$

Then there exists a constant $\bar{a} < \infty$ such that the original problem has the same value as the compact relaxed-control problem on $[0, \bar{a}]^2$, and the compact relaxed-control problem admits an optimal relaxed control

$$\vartheta_t^* \in \mathcal{P}([0, \bar{a}]^2), \quad t \in [0, T]. \quad (23)$$

If the Hamiltonian minimizer admits a progressively measurable selector, then the relaxed optimum is attained by a strict feedback control $a_t^* \in [0, \bar{a}]^2$. Moreover, after the alive-cemetery decomposition, the value function is a bounded uniformly continuous viscosity solution of the decomposed HJB equation, equivalently of the killed-state HJB written in lifted notation,

$$\partial_t V(t, \mu^1, \mu^2) + H(t, \mu^1, \mu^2, D_\mu V, D_{\mu\mu}^2 V) = 0, \quad (24)$$

with terminal condition

$$V(T, \mu^1, \mu^2) = \Phi(\mu^1(\{\partial\}), \mu^2(\{\partial\})). \quad (25)$$

Here the Hamiltonian is

$$H(t, \mu^1, \mu^2, D_\mu U, D_{\mu\mu}^2 U) = \inf_{a \in [0, \bar{a}]^2} \left\{ \pi_1 c_1(a^1) + \pi_2 c_2(a^2) + \mathcal{L}^a U(\mu^1, \mu^2) \right\}, \quad (26)$$

and for a sufficiently smooth functional $U : \mathcal{P}_2(\mathbb{E})^2 \rightarrow \mathbb{R}$, the common-noise generator $\mathcal{L}^a$ is

$$\begin{aligned} \mathcal{L}^a U(\mu^1, \mu^2) &= \sum_{k=1}^2 \int_{\mathbb{R}} \left[b_k(x, \mu^1, \mu^2) - \sum_{\ell=1}^2 \Gamma_{k\ell} \Lambda_{\ell}(\mu^{\ell}, a^{\ell}) \right] D_{\mu^k} U(\mu^1, \mu^2)(x) \mu^k(dx) \\ &\quad + \frac{1}{2} \sum_{k=1}^2 (\sigma_k^2 + \sigma_0^2) \int_{\mathbb{R}} \partial_x D_{\mu^k} U(\mu^1, \mu^2)(x) \mu^k(dx) \\ &\quad + \frac{1}{2} \sigma_0^2 \sum_{k,\ell=1}^2 \int_{\mathbb{R}} \int_{\mathbb{R}} \partial_{xy}^2 \frac{\delta^2 U}{\delta \mu^k \delta \mu^{\ell}}(\mu^1, \mu^2)(x, y) \mu^k(dx) \mu^{\ell}(dy) \\ &\quad + \sum_{k=1}^2 \int_{\mathbb{R}} \lambda_k(x, a^k) \left[\frac{\delta U}{\delta \mu^k}(\mu^1, \mu^2)(\partial) - \frac{\delta U}{\delta \mu^k}(\mu^1, \mu^2)(x) \right] \mu^k(dx). \end{aligned} \tag{27}$$

Here $D_{\mu^k} U$ denotes the Lions derivative, $\partial_x D_{\mu^k} U$ denotes its spatial derivative,

$$\frac{\delta U}{\delta \mu^k}$$

denotes the first-order linear functional derivative, and

$$\frac{\delta^2 U}{\delta \mu^k \delta \mu^{\ell}}$$

denotes the second-order linear functional derivative. In (27), the first line corresponds to drift and contagion feedback, the second line to the diagonal diffusion terms from idiosyncratic and common noise, the third line to the second-order measure derivative induced by common noise, and the fourth line to the killing jump $x \mapsto \partial$.

In representative two-particle coordinates, if one formally writes

$$v = v(t, x_1, x_2, L_1, L_2),$$

then the second-order local operator corresponding to common noise contains

$$\frac{1}{2}(\sigma_1^2 + \sigma_0^2) \partial_{x_1 x_1}^2 v + \frac{1}{2}(\sigma_2^2 + \sigma_0^2) \partial_{x_2 x_2}^2 v + \sigma_0^2 \partial_{x_1 x_2}^2 v.$$

This finite-dimensional representative form is used only to explain the cross second-order term generated by common noise in the HJB equation; the rigorous equation is still given by (24)–(27).

If, in addition, $c_k \in C^1$ is strictly convex and

$$\lambda_k(x, a) = \lambda_{0,k}(x) e^{-\eta_k a}, \quad \eta_k > 0, \tag{28}$$

then, at differentiability points of the Hamiltonian, every interior optimal control satisfies the first-order condition

$$\pi_k c'_k(a^{k,*}) = \eta_k \int_{\mathbb{R}} \lambda_k(x, a^{k,*}) Q_k^U(x) \mu^k(dx), \tag{29}$$

where

$$Q_k^U(x) := \left[\frac{\delta U}{\delta \mu^k}(\mu^1, \mu^2)(\partial) - \frac{\delta U}{\delta \mu^k}(\mu^1, \mu^2)(x) \right] - \sum_{r=1}^2 \Gamma_{rk} \int_{\mathbb{R}} D_{\mu^r} U(\mu^1, \mu^2)(y) \mu^r(dy). \tag{30}$$

Under the constraint $a^k \in [0, \bar{a}]$, (29) should be interpreted as the projection condition

$$a^{k,*} = \Pi_{[0,\bar{a}]} \left[(c'_k)^{-1} \left(\frac{\eta_k}{\pi_k} \int_{\mathbb{R}} \lambda_k(x, a^{k,*}) Q_k^U(x) \mu^k(dx) \right) \right]. \quad (31)$$

In particular, when

$$c_k(a) = \frac{1}{2} r_k a^2,$$

we have

$$a^{k,*} = \Pi_{[0,\bar{a}]} \left[\frac{\eta_k}{\pi_k r_k} \int_{\mathbb{R}} \lambda_k(x, a^{k,*}) Q_k^U(x) \mu^k(dx) \right]. \quad (32)$$

Furthermore, if for each $k = 1, 2$, the map

$$a \mapsto \pi_k c_k(a) + \int_{\mathbb{R}} \lambda_k(x, a) Q_k^U(x) \mu^k(dx)$$

is strictly convex on $[0, \bar{a}]$, then the projection condition (31) determines a unique optimal control component. A common sufficient condition is

$$Q_k^U(x) \geq 0 \quad \mu^k\text{-a.e.},$$

together with strict convexity of c_k . In that case, the convexity of $a \mapsto e^{-\eta_k a}$ and $Q_k^U \geq 0$ imply convexity of the killing term in a , and adding the strictly convex control cost gives a strictly convex objective. More generally, if $c_k \in C^2$ and there exists $m_k > 0$ such that

$$\pi_k c_k''(a) + \eta_k^2 \int_{\mathbb{R}} \lambda_k(x, a) Q_k^U(x) \mu^k(dx) \geq m_k, \quad a \in [0, \bar{a}],$$

then this one-dimensional minimization problem is strictly convex, and (31) has a unique solution on $[0, \bar{a}]$.

Proof. We first prove stability of the controlled measure flow with respect to the initial measure and the control. Fix two initial states

$$\mu = (\mu^1, \mu^2), \quad \nu = (\nu^1, \nu^2),$$

and let the two systems be constructed on the same probability space with the same common noise W^0 , the same idiosyncratic noises, the same Poisson thinning variables, and the same control a . Denote the corresponding measure flows by

$$\mu_t = (\mu_t^1, \mu_t^2), \quad \nu_t = (\nu_t^1, \nu_t^2).$$

By the Lipschitz property of the drift b_k , the boundedness of the contagion matrix Γ , the Lipschitz property of the kill intensity λ_k , and Wasserstein control after lifting the cemetery state, there exists a constant $C_T < \infty$ such that

$$\mathbb{E} \left[\sup_{s \in [t, T]} \sum_{k=1}^2 W_2(\mu_s^k, \nu_s^k) \right] \leq C_T \sum_{k=1}^2 W_2(\mu^k, \nu^k). \quad (33)$$

More precisely, a quadratic coupling estimate first gives

$$\mathbb{E} \left[\sup_{s \in [t, T]} \sum_{k=1}^2 W_2^2(\mu_s^k, \nu_s^k) \right] \leq C_T \sum_{k=1}^2 W_2^2(\mu^k, \nu^k),$$

and (33) follows by Jensen's inequality. Under synchronous coupling, the diffusion noise terms cancel, and the difference comes only from drift, contagion loss feedback, and the difference in kill intensities. Taking square expectations of the state difference on the alive state and using

$$d_{\text{BL}}(\alpha, \beta) \leq W_1(\hat{\alpha}, \hat{\beta}) \leq W_2(\hat{\alpha}, \hat{\beta})$$

controls the bounded Lipschitz distance between surviving sub-probability measures by the lifted Wasserstein distance. The dead mass has already been placed at ∂ , so

$$|L^k(\mu_s^k) - L^k(\nu_s^k)| \leq W_1(\mu_s^k, \nu_s^k) \leq W_2(\mu_s^k, \nu_s^k).$$

Gronwall's inequality then gives the stated estimates.

Similarly, if two systems start from the same initial state but use controls a and $\tilde{a}$, respectively, then

$$\mathbb{E} \left[\sup_{s \in [t, T]} \sum_{k=1}^2 W_2(\mu_s^{k,a}, \mu_s^{k,\tilde{a}}) \right] \leq C_T \sum_{k=1}^2 \mathbb{E} \int_t^T \left\| \lambda_k(\cdot, a_r^k) - \lambda_k(\cdot, \tilde{a}_r^k) \right\|_{\infty} dr. \quad (34)$$

Since $0 \leq \lambda_k \leq \Lambda$, if the two controls differ only on a set $B \subset [t, T]$, then

$$\mathbb{E} \left[\sup_{s \in [t, T]} \sum_{k=1}^2 W_2(\mu_s^{k,a}, \mu_s^{k,\tilde{a}}) \right] \leq C_T |B|. \quad (35)$$

By the Lipschitz property of Φ and (33), the value function is Lipschitz in the initial measure:

$$\left| V(t, \mu^1, \mu^2) - V(t, \nu^1, \nu^2) \right| \leq C_T \sum_{k=1}^2 W_2(\mu^k, \nu^k). \quad (36)$$

Using the bounded kill intensities, the linear-growth drift, and diffusion estimates in (20), we also obtain uniform continuity in time:

$$|V(t, \mu^1, \mu^2) - V(s, \mu^1, \mu^2)| \leq C_T |t - s|^{1/2} + C_T |t - s|. \quad (37)$$

We next prove compactification of the control set. Since c_k is convex with superlinear growth, there exist $\bar{a} < \infty$ and a constant C_0 such that, whenever $a > \bar{a}$,

$$c_k(a) - c_k(\bar{a}) > C_0, \quad (38)$$

where C_0 is chosen larger than the maximal terminal-loss variation constant generated by (35) and the Lipschitz constant of Φ . For any admissible control a , define the truncated control

$$a_s^{k,\bar{a}} := a_s^k \wedge \bar{a}, \quad k = 1, 2.$$

Let

$$B_k := \{s \in [t, T] : a_s^k > \bar{a}\}.$$

By (35) and the Lipschitz property of Φ , the increase in terminal cost due to truncation is at most

$$C \sum_{k=1}^2 |B_k|. \quad (39)$$

On the other hand, by (38), the decrease in running cost is at least

$$C_0 \sum_{k=1}^2 |B_k|. \quad (40)$$

Taking $C_0 > C$, we get

$$J(t, \mu^1, \mu^2; a^{\bar{a}}) \leq J(t, \mu^1, \mu^2; a). \quad (41)$$

Thus the original noncompact control problem and the compact control problem

$$a_t \in [0, \bar{a}]^2$$

have the same value function, and it is sufficient to search for an optimal control in $[0, \bar{a}]^2$.

We now prove existence of an optimal relaxed control on the compact control set. Take a minimizing sequence of relaxed controls $(\vartheta^n)_{n \geq 1}$, and let $(\mu^{1,n}, \mu^{2,n})$ be the corresponding measure flows. Since $[0, \bar{a}]^2$ is compact, the space $\mathcal{P}([0, \bar{a}]^2)$ is compact under weak convergence. The bounded kill intensity and the linear-growth drift give tightness of the corresponding controlled measure flows in

$$C([t, T]; \mathcal{P}_2(\mathbf{E})^2).$$

By Prokhorov's theorem and the compactness of relaxed controls, there is a subsequence along which the relaxed control-state laws converge weakly. By the stability of the controlled martingale problem under weak convergence in Assumption [SM4.1](#), the limiting state process satisfies the relaxed version of [\(20\)](#); denote the limiting relaxed control by ϑ^* . By lower semicontinuity of the running cost and continuity of the terminal cost,

$$J(t, \mu^1, \mu^2; \vartheta^*) \leq \liminf_{n \rightarrow \infty} J(t, \mu^1, \mu^2; \vartheta^n) = V(t, \mu^1, \mu^2).$$

Thus ϑ^* is an optimal relaxed control.

The compact relaxed-control problem admits an optimal relaxed control. Under the standard chattering condition, every relaxed control can be approximated in the weak occupation-measure topology by progressively measurable strict controls, and the stability estimate [\(34\)](#) transfers this approximation to the measure flow and terminal cost. The relaxed value then agrees with the closure of strict-control values. If the pointwise Hamiltonian minimizer admits a progressively measurable selector, selecting this minimizer gives a strict feedback control attaining the same value. This proves [\(23\)](#).

We now establish the dynamic programming principle. By [\(33\)](#), the controlled measure flow depends continuously on the current state. By [\(36\)](#) and [\(37\)](#), the value function is uniformly continuous on the Wasserstein state space. The controlled martingale problem is stable under concatenation of controls. Hence, for every stopping time $\theta \in [t, T]$,

$$V(t, \mu^1, \mu^2) = \inf_{\vartheta} \mathbb{E} \left[\int_t^\theta \int_{[0, \bar{a}]^2} (\pi_1 c_1(a^1) + \pi_2 c_2(a^2)) \vartheta_s(da) ds + V(\theta, \mu_\theta^1, \mu_\theta^2) \right]. \quad (42)$$

Here (μ_s^1, μ_s^2) is the controlled measure flow starting from (t, μ^1, μ^2) under the relaxed control ϑ . One inequality follows by using an arbitrary relaxed control until θ and then concatenating an ε -optimal relaxed control after θ ; the other follows because the conditional remaining cost of any global relaxed control after θ is bounded below by $V(\theta, \mu_\theta^1, \mu_\theta^2)$.

We next turn [\(42\)](#) into the HJB equation. Take a sufficiently rich nonatomic probability space $(\Omega, \mathcal{F}, \mathbb{P})$. For each $\mu^k \in \mathcal{P}_2(\mathbf{E})$, choose a random variable

$$\xi^k \in L^2(\Omega; \mathbf{E}), \quad \mathcal{L}(\xi^k) = \mu^k.$$

Define the Lions lift

$$\mathcal{V}(t, \xi^1, \xi^2) := V(t, \mathcal{L}(\xi^1), \mathcal{L}(\xi^2)). \quad (43)$$

If U is a smooth functional on $\mathcal{P}_2(\mathbf{E})^2$, set

$$\mathcal{U}(t, \xi^1, \xi^2) := U(t, \mathcal{L}(\xi^1), \mathcal{L}(\xi^2)).$$

The Lions derivative and the Fréchet derivative of the lifted functional satisfy

$$D_{\xi^k} \mathcal{U}(t, \xi^1, \xi^2) = D_{\mu^k} U(t, \mu^1, \mu^2)(\xi^k). \quad (44)$$

The second derivative correspondence gives the common-noise term

$$\frac{1}{2} \sigma_0^2 \sum_{k, \ell=1}^2 \int_{\mathbb{R}} \int_{\mathbb{R}} \partial_{xy}^2 \frac{\delta^2 U}{\delta \mu^k \delta \mu^\ell}(\mu^1, \mu^2)(x, y) \mu^k(dx) \mu^\ell(dy). \quad (45)$$

Apply the Hilbert-space Itô formula with jumps to the lifted test function. The continuous martingale part yields the diffusion terms and the common-noise second-order term; the compensator of the killing jump $x \mapsto \partial$ is

$$\lambda_k(x, a^k) \left[\frac{\delta U}{\delta \mu^k}(\mu^1, \mu^2)(\partial) - \frac{\delta U}{\delta \mu^k}(\mu^1, \mu^2)(x) \right]. \quad (46)$$

Thus the lifted infinite-dimensional HJB equation is

$$\partial_t \mathcal{V} + \inf_{a \in [0, \bar{a}]^2} \left\{ \pi_1 c_1(a^1) + \pi_2 c_2(a^2) + \mathcal{L}^a \mathcal{V} \right\} = 0, \quad (47)$$

where $\mathcal{L}^a$, when projected back to measure space, is exactly (27).

We now verify the viscosity solution property. We use the following sign convention for the backward HJB equation: if a smooth test function touches the value function from above, then

$$\partial_t \phi + H(\phi) \geq 0;$$

if it touches from below, then

$$\partial_t \phi + H(\phi) \leq 0.$$

Let $\mathcal{U}$ be a smooth test function on $L^2(\Omega; \mathbf{E})^2$, and suppose that

$$\mathcal{V} - \mathcal{U}$$

has a local maximum at (t, ξ^1, ξ^2) . Fix any constant control $a \in [0, \bar{a}]^2$. By the dynamic programming principle, for small $h > 0$,

$$\mathcal{V}(t, \xi^1, \xi^2) \leq \mathbb{E} \left[\int_t^{t+h} (\pi_1 c_1(a^1) + \pi_2 c_2(a^2)) ds + \mathcal{V}(t+h, \xi_{t+h}^1, \xi_{t+h}^2) \right].$$

Since $\mathcal{V} - \mathcal{U}$ has a local maximum at the initial point,

$$0 \leq \mathbb{E} \left[\int_t^{t+h} (\pi_1 c_1(a^1) + \pi_2 c_2(a^2)) ds + \mathcal{U}(t+h, \xi_{t+h}^1, \xi_{t+h}^2) - \mathcal{U}(t, \xi^1, \xi^2) \right].$$

Applying the Hilbert-space Itô formula with jumps to $\mathcal{U}$, dividing by h , and sending $h \downarrow 0$, we obtain

$$0 \leq \partial_t \mathcal{U} + \mathcal{L}^a \mathcal{U} + \pi_1 c_1(a^1) + \pi_2 c_2(a^2).$$

Since $a \in [0, \bar{a}]^2$ is arbitrary,

$$\partial_t \mathcal{U} + \inf_{a \in [0, \bar{a}]^2} \left\{ \mathcal{L}^a \mathcal{U} + \pi_1 c_1(a^1) + \pi_2 c_2(a^2) \right\} \geq 0. \quad (48)$$

Conversely, let $\mathcal{U}$ be a smooth test function such that

$$\mathcal{V} - \mathcal{U}$$

has a local minimum at (t, ξ^1, ξ^2) . By the chattering equivalence of relaxed and strict values, an ε -optimal relaxed control can be approximated by a strict representative without changing the first-order Hamiltonian inequality. We therefore write the local argument for an ε -optimal strict control a^ε . By the dynamic programming principle,

$$\mathcal{V}(t, \xi^1, \xi^2) + \varepsilon h \geq \mathbb{E} \left[\int_t^{t+h} (\pi_1 c_1(a_s^{1,\varepsilon}) + \pi_2 c_2(a_s^{2,\varepsilon})) ds + \mathcal{V}(t+h, \xi_{t+h}^1, \xi_{t+h}^2) \right].$$

The local minimum property allows us to replace $\mathcal{V}$ on the right-hand side by $\mathcal{U}$, so

$$\varepsilon h \geq \mathbb{E} \left[\int_t^{t+h} (\pi_1 c_1(a_s^{1,\varepsilon}) + \pi_2 c_2(a_s^{2,\varepsilon})) ds + \mathcal{U}(t+h, \xi_{t+h}^1, \xi_{t+h}^2) - \mathcal{U}(t, \xi^1, \xi^2) \right].$$

Applying the Hilbert-space Itô formula with jumps, dividing by h , sending $h \downarrow 0$, and then letting $\varepsilon \downarrow 0$, gives

$$\partial_t \mathcal{U} + \inf_{a \in [0, \bar{a}]^2} \left\{ \mathcal{L}^a \mathcal{U} + \pi_1 c_1(a^1) + \pi_2 c_2(a^2) \right\} \leq 0. \quad (49)$$

By (48) and (49), $\mathcal{V}$ is a viscosity solution of the lifted Hilbert-space equation. By the correspondence between Lions lifts and derivatives on Wasserstein space, V is a Wasserstein–Lions viscosity solution of (24).

The terminal condition follows directly from the definition of the objective. At $t = T$, the running cost integral vanishes, and no subsequent control can change the cemetery masses already recorded in the state. Therefore

$$\begin{aligned} V(T, \mu^1, \mu^2) &= \inf_{\vartheta} \mathbb{E} \left[\Phi(L_T^1, L_T^2) \right] \\ &= \Phi(\mu^1(\{\partial\}), \mu^2(\{\partial\})), \end{aligned}$$

which proves (25).

The comparison theorem proved in Theorem SM4.7 applies to the killed two-population HJB under the bounded Lipschitz killing, compact-control, Lipschitz-drift, constant-diffusion, and alive-diffusion nondegeneracy assumptions above. Hence any bounded uniformly continuous viscosity subsolution and supersolution ordered at terminal time remain ordered on $[0, T]$. Applying this result to two viscosity solutions of (24) gives uniqueness in the class of bounded uniformly continuous Wasserstein–Lions viscosity solutions. The value function V constructed by dynamic programming is therefore the unique viscosity solution of this HJB equation in that class.

The viscosity characterization above does not require classical differentiability of the value function. The following first-order condition is a smooth-verification statement: assume that, on the region under consideration, a strict Hamiltonian minimizer exists and the value admits Lions derivatives that identify the marginal cemetery-jump and loss-feedback terms. The relaxed formulation gives the same Hamiltonian because the generator and the population-weighted running cost are averaged linearly over actions, so the infimum over relaxed actions is attained at an extreme point whenever a strict measurable minimizer exists. For a smooth functional U , the part of the Hamiltonian depending on a^k is

$$\pi_k c_k(a^k) + \int_{\mathbb{R}} \lambda_k(x, a^k) \left[\frac{\delta U}{\delta \mu^k}(\mu^1, \mu^2)(\partial) - \frac{\delta U}{\delta \mu^k}(\mu^1, \mu^2)(x) \right] \mu^k(dx)$$

together with the a^k -dependent part in the contagion-feedback drift,

$$-\sum_{r=1}^2 \int_{\mathbb{R}} \Gamma_{rk} \Lambda_k(\mu^k, a^k) D_{\mu^r} U(\mu^1, \mu^2)(y) \mu^r(dy).$$

Combining these two parts yields

$$\pi_k c_k(a^k) + \int_{\mathbb{R}} \lambda_k(x, a^k) Q_k^U(x) \mu^k(dx),$$

where Q_k^U is exactly (30). If

$$\lambda_k(x, a) = \lambda_{0,k}(x) e^{-\eta_k a},$$

then

$$\partial_a \lambda_k(x, a) = -\eta_k \lambda_k(x, a).$$

Differentiating the a^k -component of the Hamiltonian, an interior minimizer satisfies

$$0 = \pi_k c'_k(a^{k,*}) - \eta_k \int_{\mathbb{R}} \lambda_k(x, a^{k,*}) Q_k^U(x) \mu^k(dx),$$

which is (29). On the constrained interval $[0, \bar{a}]$, first-order optimality together with endpoint complementarity is equivalent to the projected form (31). If $c_k(a) = \frac{1}{2} r_k a^2$, then the population-weighted derivative is $\pi_k r_k a$, giving (32). The proof is complete. $\square$

Assumption SM4.9 (Regularity conditions for the information value of local pressure information). Fix a terminal time $T > 0$, and suppose that the system contains N nodes and K types. The type of node i is denoted by

$$g(i) \in \{1, \dots, K\}, \quad N_k := |\{i : g(i) = k\}|, \quad \pi_k := \frac{N_k}{N}.$$

The local graph pressure is

$$P_{i,t}^{\text{graph}} := \sum_{j=1}^N A_{ij} \Delta N_{j,t},$$

where $A_{ij} \geq 0$ is the edge-weight shock from source node j to target node i . The type-average pressure is

$$P_{k,t}^{\text{type}} := \frac{1}{N_k} \sum_{i:g(i)=k} P_{i,t}^{\text{graph}}.$$

Let

$$\mathbb{F}^{\text{type}} = \left(\mathcal{F}_t^{\text{type}} \right)_{0 \leq t \leq T}, \quad \mathbb{F}^{\text{graph}} = \left(\mathcal{F}_t^{\text{graph}} \right)_{0 \leq t \leq T}$$

denote the type-average filtration and the node graph-pressure filtration, respectively, and assume that

$$\mathcal{F}_t^{\text{type}} \subset \mathcal{F}_t^{\text{graph}}, \quad 0 \leq t \leq T.$$

Define the admissible type-average control set by

$$\mathfrak{A}^{\text{type}} := \left\{ a : a_{i,t} = \alpha_{g(i),t}, \alpha_{k,t} \text{ is } \mathbb{F}^{\text{type}}\text{-progressively measurable} \right\},$$

and the node graph-pressure control set by

$$\mathfrak{A}^{\text{graph}} := \left\{ a : a_{i,t} \text{ is } \mathbb{F}^{\text{graph}}\text{-progressively measurable} \right\}.$$

Then

$$\mathfrak{A}^{\text{type}} \subset \mathfrak{A}^{\text{graph}}.$$

Controls take values in the compact interval

$$\mathcal{A} = [0, \bar{a}].$$

The objective is

$$J(a) := \mathbb{E} \left[\Phi(L_T^{1,N}, \dots, L_T^{K,N}) + \int_0^T \frac{1}{N} \sum_{i=1}^N c_{g(i)}(a_{i,t}) dt \right],$$

where $\Phi : [0, 1]^K \rightarrow \mathbb{R}$ is bounded and nondecreasing in each coordinate. Assume further that there exists a constant

$$\underline{\theta}_k > 0$$

such that, in the terminal loss region considered in this paper, for every $r \geq 0$,

$$\Phi(\ell + re_k) - \Phi(\ell) \geq \underline{\theta}_k r.$$

The control cost $c_k \in C^2([0, \bar{a}])$, and on some compact interior interval

$$[\underline{a}, \bar{a} - \underline{a}], \quad 0 < \underline{a} < \frac{\bar{a}}{2},$$

it satisfies

$$c'_k(a) \geq \underline{c}_k > 0, \quad |c''_k(a)| \leq \bar{c}_k.$$

Assume that the type-average problem has an optimal control

$$a^{\text{type},*} \in \mathfrak{A}^{\text{type}}, \quad a_{i,t}^{\text{type},*} = \alpha_{g(i),t}^*,$$

and write

$$a^0 := a^{\text{type},*}.$$

There exist a type k , an $\mathbb{F}^{\text{graph}}$ -stopping time $\tau < T$, a window length $h > 0$ with $\tau + h \leq T$, and an event

$$\mathcal{E} \in \mathcal{F}_\tau^{\text{graph}}, \quad \mathbb{P}^{a^0}(\mathcal{E}) \geq q_k > 0,$$

such that on $\mathcal{E}$, the local graph pressure of type k has nondegenerate within-block variance. The probability space is fixed in strong form: the baseline type-level optimizer and all local perturbation controls below are constructed from the same Brownian motions, common noise, and killing-thinning random variables. Since every perturbation introduced below coincides with the baseline control before τ , the event $\mathcal{E}$ and the pressure vector $(Z_i)_{g(i)=k}$ are the same observable $\mathcal{F}_\tau^{\text{graph}}$ -measurable objects under the baseline and perturbed controls. Specifically, let

$$\bar{P}_{k,\tau} := \frac{1}{N_k} \sum_{i:g(i)=k} P_{i,\tau}^{\text{graph}}, \quad Z_i := P_{i,\tau}^{\text{graph}} - \bar{P}_{k,\tau}, \quad g(i) = k.$$

Assume that

$$\frac{1}{N_k} \sum_{i:g(i)=k} Z_i = 0,$$

and that, on $\mathcal{E}$,

$$S_k := \frac{1}{N_k} \sum_{i:g(i)=k} Z_i^2 \geq v_k > 0, \quad |Z_i| \leq R_k.$$

At the same time, the type-average optimal control lies in the interior of the control interval over this window:

$$\alpha_{k,t}^* \in [\underline{a}, \bar{a} - \underline{a}], \quad t \in [\tau, \tau + h], \quad \text{on } \mathcal{E}.$$

For any perturbation profile

$$r = (r_i)_{i:g(i)=k}, \quad r_i \in L^\infty([\tau, \tau + h]),$$

define the concatenated control a^r as follows: on the window $[\tau, \tau + h]$, for nodes of type k , set

$$a_{i,t}^r = \alpha_{k,t}^* + r_i(t), \quad g(i) = k,$$

and keep all other nodes and all other time periods equal to a^0 . Let

$$\mathcal{H}_\tau(r) := \mathbb{E} \left[\Phi(L_T^{1,N}(a^r), \dots, L_T^{K,N}(a^r)) \mid \mathcal{F}_\tau^{\text{graph}} \right]$$

denote the conditional terminal loss cost from time τ under the local perturbation r . Assume that $\mathcal{H}_\tau$ is twice Gateaux differentiable in a neighborhood of $r = 0$, and that there exists $C_H < \infty$ such that for all sufficiently small r, u ,

$$|D\mathcal{H}_\tau(0)[u]| \leq C_H \frac{1}{N_k} \sum_{i:g(i)=k} \|u_i\|_\infty,$$

and

$$|D^2\mathcal{H}_\tau(r)[u, u]| \leq C_H \frac{1}{N_k} \sum_{i:g(i)=k} \|u_i\|_\infty^2.$$

Finally, assume that the local pressure direction has strictly positive marginal benefit for terminal loss. This is the formal version of strict monotonicity of the marginal loss-reduction benefit in local pressure: on $\mathcal{E}$, reallocating an infinitesimal unit of control toward nodes with larger $P_{i,\tau}^{\text{graph}}$ strictly lowers the conditional terminal loss cost. Quantitatively,

$$-D\mathcal{H}_\tau(0)[Z] \geq \pi_k \underline{\theta}_k m_k S_k,$$

where Z denotes the perturbation profile constant in time,

$$Z_i(t) = Z_i, \quad t \in [\tau, \tau + h], \quad g(i) = k,$$

and $m_k > 0$ is the marginal benefit constant of node-level pressure information.

Theorem SM4.10 (Quantification of the information value of local pressure information). *Under Assumption SM4.9, define*

$$V^{\text{type}} := \inf_{a \in \mathfrak{A}^{\text{type}}} J(a), \quad V^{\text{graph}} := \inf_{a \in \mathfrak{A}^{\text{graph}}} J(a).$$

Then

$$V^{\text{graph}} \leq V^{\text{type}}.$$

Furthermore, if type k has nondegenerate local pressure variance on the event $\mathcal{E}$, namely

$$S_k \geq v_k > 0,$$

then there exists a constant

$$\delta_{\text{info}} > 0$$

such that

$$V^{\text{type}} - V^{\text{graph}} \geq \delta_{\text{info}}.$$

More specifically, there exists $\varepsilon_0 > 0$ such that for every

$$0 < \varepsilon \leq \varepsilon_0$$

we have

$$V^{\text{type}} - V^{\text{graph}} \geq \frac{1}{2} q_k \pi_k \theta_k m_k \varepsilon v_k.$$

Thus node-level graph-pressure information has strictly positive information value relative to type-average information.

If, in addition, there exist a constant $c_\Gamma > 0$ and a source-side new-loss vector

$$y = (y_1, \dots, y_K) \in \mathbb{R}_+^K$$

such that on $\mathcal{E}$,

$$S_k \geq c_\Gamma \text{Var}_{i:g(i)=k} \left(\sum_{\ell=1}^K y_\ell \sum_{j:g(j)=\ell} A_{ij} \right),$$

then

$$V^{\text{type}} - V^{\text{graph}} \geq \frac{1}{2} q_k \pi_k \theta_k m_k \varepsilon c_\Gamma \text{Var}_{i:g(i)=k} \left(\sum_{\ell=1}^K y_\ell \sum_{j:g(j)=\ell} A_{ij} \right).$$

This lower bound directly quantifies the information value of local pressure information as a function of within-block weighted in-exposure heterogeneity, contagion edge-weight structure, and the marginal control benefit in the kill intensity.

Proof. First, the filtration inclusion

$$\mathcal{F}_t^{\text{type}} \subset \mathcal{F}_t^{\text{graph}}, \quad 0 \leq t \leq T,$$

implies that every type-average admissible control is also a node graph-pressure admissible control. Therefore

$$\mathfrak{A}^{\text{type}} \subset \mathfrak{A}^{\text{graph}}.$$

Taking the infimum of the same objective over the two control classes gives

$$V^{\text{graph}} = \inf_{a \in \mathfrak{A}^{\text{graph}}} J(a) \leq \inf_{a \in \mathfrak{A}^{\text{type}}} J(a) = V^{\text{type}}.$$

This proves the weak information value inequality.

We now prove the strictly positive lower bound. Let

$$a^0 := a^{\text{type},*}, \quad J(a^0) = V^{\text{type}}.$$

On the event $\mathcal{E}$, define the centered pressure variables within type k by

$$Z_i = P_{i,\tau}^{\text{graph}} - \bar{P}_{k,\tau}, \quad \bar{P}_{k,\tau} = \frac{1}{N_k} \sum_{i:g(i)=k} P_{i,\tau}^{\text{graph}}.$$

By construction,

$$\frac{1}{N_k} \sum_{i:g(i)=k} Z_i = 0.$$

On $\mathcal{E}$,

$$S_k = \frac{1}{N_k} \sum_{i:g(i)=k} Z_i^2 \geq v_k, \quad S_k \leq R_k^2.$$

We construct a local perturbation control measurable with respect to graph pressure. For each

$$t \in [\tau, \tau + h]$$

and parameter $\varepsilon > 0$, define

$$F_t(b, \varepsilon) := \frac{1}{N_k} \sum_{i:g(i)=k} c_k(\alpha_{k,t}^* + b + \varepsilon Z_i) - c_k(\alpha_{k,t}^*).$$

Since

$$F_t(0, 0) = 0$$

and

$$\partial_b F_t(0, 0) = c'_k(\alpha_{k,t}^*) \geq \underline{c}_k,$$

the implicit function theorem applies uniformly over the whole window. Therefore, for sufficiently small ε , there exists a unique progressively measurable process

$$b_\varepsilon(t), \quad t \in [\tau, \tau + h],$$

such that

$$F_t(b_\varepsilon(t), \varepsilon) = 0, \quad t \in [\tau, \tau + h].$$

Moreover,

$$\partial_\varepsilon F_t(0, 0) = c'_k(\alpha_{k,t}^*) \frac{1}{N_k} \sum_{i:g(i)=k} Z_i = 0,$$

so

$$b_\varepsilon(t) = O(\varepsilon^2 S_k)$$

uniformly in $t \in [\tau, \tau + h]$. In particular, there exists

$$C_b = C_b(\underline{c}_k, \bar{c}_k, R_k) < \infty$$

such that

$$\sup_{t \in [\tau, \tau + h]} |b_\varepsilon(t)| \leq C_b \varepsilon^2 S_k \leq C_b \varepsilon^2 R_k^2.$$

Take

$$\varepsilon_{\text{adm}} := \min \left\{ 1, \frac{\underline{a}}{2(R_k + C_b R_k^2)} \right\}.$$

Whenever

$$0 < \varepsilon \leq \varepsilon_{\text{adm}},$$

we have

$$\begin{aligned} |b_\varepsilon(t) + \varepsilon Z_i| &\leq C_b \varepsilon^2 R_k^2 + \varepsilon R_k \\ &\leq \varepsilon (C_b R_k^2 + R_k) \leq \frac{\underline{a}}{2}. \end{aligned}$$

Since

$$\alpha_{k,t}^* \in [\underline{a}, \bar{a} - \underline{a}], \quad t \in [\tau, \tau + h],$$

we get

$$\alpha_{k,t}^* + b_\varepsilon(t) + \varepsilon Z_i \in [0, \bar{a}], \quad g(i) = k, \quad t \in [\tau, \tau + h].$$

Thus the following control is admissible:

$$\tilde{a}_{i,t} := \begin{cases} \alpha_{k,t}^* + b_\varepsilon(t) + \varepsilon Z_i, & \text{if } g(i) = k, \ t \in [\tau, \tau + h], \ \omega \in \mathcal{E}, \\ a_{i,t}^0, & \text{otherwise.} \end{cases}$$

Since Z_i is $\mathcal{F}_\tau^{\text{graph}}$ -measurable, τ is an $\mathbb{F}^{\text{graph}}$ -stopping time, and $b_\varepsilon(t)$ is determined pointwise by $\alpha_{k,t}^*$ and $(Z_i)_{g(i)=k}$, we have

$$\tilde{a} \in \mathfrak{A}^{\text{graph}}.$$

By the definition of $b_\varepsilon(t)$, on $\mathcal{E}$ there is exact pointwise running-cost neutralization for every $t \in [\tau, \tau + h]$:

$$\frac{1}{N_k} \sum_{i: g(i)=k} c_k(\tilde{a}_{i,t}) = c_k(\alpha_{k,t}^*).$$

On $\mathcal{E}^c$, the control is unchanged. Other types and times outside the window are also unchanged. Hence

$$\int_0^T \frac{1}{N} \sum_{i=1}^N c_{g(i)}(\tilde{a}_{i,t}) dt = \int_0^T \frac{1}{N} \sum_{i=1}^N c_{g(i)}(a_{i,t}^0) dt.$$

Thus local control reallocation does not increase the running control cost.

We now compare terminal loss costs. Let the perturbation profile be

$$r_i^\varepsilon(t) := b_\varepsilon(t) + \varepsilon Z_i, \quad g(i) = k, \quad t \in [\tau, \tau + h],$$

and set $r^\varepsilon = 0$ on $\mathcal{E}^c$. On $\mathcal{E}$, Taylor expansion gives

$$\mathcal{H}_\tau(0) - \mathcal{H}_\tau(r^\varepsilon) = -D\mathcal{H}_\tau(0)[r^\varepsilon] - \frac{1}{2}D^2\mathcal{H}_\tau(\theta r^\varepsilon)[r^\varepsilon, r^\varepsilon],$$

where $\theta = \theta(\omega) \in (0, 1)$.

Decompose

$$r_i^\varepsilon(t) = \varepsilon Z_i + b_\varepsilon(t)$$

into the pressure-direction perturbation and the common cost-correction term. By the marginal benefit condition in the local pressure direction,

$$-D\mathcal{H}_\tau(0)[\varepsilon Z] \geq \pi_k \underline{\theta}_k m_k \varepsilon S_k.$$

By boundedness of the first derivative and

$$\sup_{t \in [\tau, \tau + h]} |b_\varepsilon(t)| \leq C_b \varepsilon^2 S_k,$$

there exists $C_1 < \infty$ such that

$$|D\mathcal{H}_\tau(0)[b_\varepsilon]| \leq C_1 \varepsilon^2 S_k.$$

By boundedness of the second derivative,

$$\begin{aligned} |D^2\mathcal{H}_\tau(\theta r^\varepsilon)[r^\varepsilon, r^\varepsilon]| &\leq C_H \frac{1}{N_k} \sum_{i: g(i)=k} \|r_i^\varepsilon\|_\infty^2 \\ &\leq C_2 \varepsilon^2 S_k, \end{aligned}$$

where $C_2 < \infty$ depends only on C_H, C_b, R_k . Hence there exists

$$C_{\text{rem}} < \infty$$

such that on $\mathcal{E}$,

$$\mathcal{H}_\tau(0) - \mathcal{H}_\tau(r^\varepsilon) \geq \left(\pi_k \underline{\theta}_k m_k \varepsilon - C_{\text{rem}} \varepsilon^2 \right) S_k.$$

Take

$$\varepsilon_0 := \min \left\{ \varepsilon_{\text{adm}}, \frac{\pi_k \underline{\theta}_k m_k}{2C_{\text{rem}}} \right\}.$$

If $0 < \varepsilon \leq \varepsilon_0$, then

$$\mathcal{H}_\tau(0) - \mathcal{H}_\tau(r^\varepsilon) \geq \frac{1}{2} \pi_k \underline{\theta}_k m_k \varepsilon S_k.$$

Since $S_k \geq v_k$ on $\mathcal{E}$,

$$\mathcal{H}_\tau(0) - \mathcal{H}_\tau(r^\varepsilon) \geq \frac{1}{2} \pi_k \underline{\theta}_k m_k \varepsilon v_k \quad \text{on } \mathcal{E}.$$

On $\mathcal{E}^c$, the perturbation is zero, so the conditional terminal cost difference is zero. Taking expectation and using

$$\mathbb{P}^{a^0}(\mathcal{E}) \geq q_k,$$

we obtain

$$\mathbb{E} \left[\Phi(L_T(a^0)) - \Phi(L_T(\tilde{a})) \right] \geq \frac{1}{2} q_k \pi_k \underline{\theta}_k m_k \varepsilon v_k.$$

Combining this with the exact neutralization of running costs gives

$$J(a^0) - J(\tilde{a}) \geq \frac{1}{2} q_k \pi_k \underline{\theta}_k m_k \varepsilon v_k.$$

Since

$$J(a^0) = V^{\text{type}}$$

and

$$\tilde{a} \in \mathfrak{A}^{\text{graph}},$$

we have

$$V^{\text{graph}} \leq J(\tilde{a}) \leq V^{\text{type}} - \frac{1}{2} q_k \pi_k \underline{\theta}_k m_k \varepsilon v_k.$$

Therefore

$$V^{\text{type}} - V^{\text{graph}} \geq \frac{1}{2} q_k \pi_k \underline{\theta}_k m_k \varepsilon v_k > 0.$$

This proves the strictly positive value of local pressure information.

Finally, if there exist a constant $c_\Gamma > 0$ and a source-side new-loss vector $y \in \mathbb{R}_+^K$ such that

$$S_k \geq c_\Gamma \text{Var}_{i:g(i)=k} \left(\sum_{\ell=1}^K y_\ell \sum_{j:g(j)=\ell} A_{ij} \right),$$

then substituting this estimate into the preceding lower bound gives

$$V^{\text{type}} - V^{\text{graph}} \geq \frac{1}{2} q_k \pi_k \underline{\theta}_k m_k \varepsilon c_\Gamma \text{Var}_{i:g(i)=k} \left(\sum_{\ell=1}^K y_\ell \sum_{j:g(j)=\ell} A_{ij} \right).$$

The theorem is proved. □

Remark SM4.11 (A sufficient form of the marginal benefit condition). The condition in Assumption [SM4.9](#)

$$-D\mathcal{H}_\tau(0)[Z] \geq \pi_k \underline{\theta}_k m_k S_k$$

can be derived from more primitive conditions on the kill intensity. Suppose that within the local window, the main terminal new-death probability of node i is approximated by

$$\Psi_i(a.) = 1 - \exp \left\{ - \int_\tau^{\tau+h} \lambda_k \left(X_{i,\tau} - P_{i,\tau}^{\text{graph}}, a_t \right) dt \right\},$$

and that the kill intensity takes the exponential control form

$$\lambda_k(x, a) = \lambda_{0,k}(x) e^{-\eta_k a}, \quad \eta_k > 0.$$

Then

$$-\partial_a \lambda_k(x, a) = \eta_k \lambda_k(x, a) > 0.$$

If control has a larger marginal reduction in killing probability when pressure is higher, and if there exists $\underline{m}_k > 0$ such that on the relevant compact region

$$\frac{1}{N_k} \sum_{i:g(i)=k} Z_i \exp\{-I_i^0\} \int_\tau^{\tau+h} \left[-\partial_a \lambda_k \left(X_{i,\tau} - P_{i,\tau}^{\text{graph}}, \alpha_{k,t}^* \right) \right] dt \geq \underline{m}_k \frac{1}{N_k} \sum_{i:g(i)=k} Z_i^2,$$

where

$$I_i^0 := \int_\tau^{\tau+h} \lambda_k \left(X_{i,\tau} - P_{i,\tau}^{\text{graph}}, \alpha_{k,t}^* \right) dt,$$

and if the nonnegative contagion feedback after the local window does not offset this first-order benefit, then one may take

$$m_k = \underline{m}_k$$

or a positive lower bound obtained after subtracting second-order feedback errors. This condition states that within the same type, nodes with higher pressure have a larger marginal reduction in terminal loss from one unit of control. Therefore, reallocating a fixed cost budget from low-pressure nodes to high-pressure nodes strictly reduces terminal risk.

Lemma SM4.12 (Pressure-monotone verification in a nonsaturated exponential-killing window). *Assume, in the setting of Remark [SM4.11](#), that on the compact local window*

$$\lambda_k(x, a) = \lambda_{\text{base},k} \exp\{\gamma_k(x_{b,k} - x)\} e^{-\eta_k a}, \quad \gamma_k, \eta_k > 0,$$

that $X_{i,\tau}$ and $\alpha_{k,t}^$ remain in a compact set, and that the local cumulative intensity is nonsaturated:*

$$I_i^0 = \int_\tau^{\tau+h} \lambda_k(X_{i,\tau} - P_{i,\tau}^{\text{graph}}, \alpha_{k,t}^*) dt \leq 1 - \nu$$

for some $\nu > 0$. Then the marginal reduction of the local killing probability with respect to control is strictly increasing in local pressure. More precisely, there exists $m_{\text{loc}} > 0$, depending only on the compact region and ν , such that for any centered pressure perturbation $Z_i = P_{i,\tau}^{\text{graph}} - \bar{P}_{k,\tau}$,

$$\frac{1}{N_k} \sum_{i:g(i)=k} Z_i \exp\{-I_i^0\} \int_\tau^{\tau+h} \left[-\partial_a \lambda_k(X_{i,\tau} - P_{i,\tau}^{\text{graph}}, \alpha_{k,t}^*) \right] dt \geq m_{\text{loc}} \frac{1}{N_k} \sum_{i:g(i)=k} Z_i^2$$

whenever the compact window also gives a positive lower bound on the pressure derivative of the displayed marginal-reduction function. Thus the marginal-benefit assumption in Theorem [SM4.10](#) is verified on nonsaturated regions after subtracting any explicitly bounded second-order feedback error.

Proof. For the exponential state dependence, increasing pressure replaces x by $x - P$, so

$$\lambda_k(X - P, a) = \lambda_{\text{base},k} \exp\{\gamma_k(x_{b,k} - X + P)\} e^{-\eta_k a}$$

is strictly increasing in P . The local marginal reduction of the killing probability has the form

$$G(P) = \exp\{-I(P)\} \int_{\tau}^{\tau+h} \eta_k \lambda_k(X - P, \alpha_{k,t}^*) dt,$$

where $I(P) = \int_{\tau}^{\tau+h} \lambda_k(X - P, \alpha_{k,t}^*) dt$. In the homogeneous exponential case this is $G(P) = \eta_k I(P) e^{-I(P)}$. On the nonsaturated region $I(P) \leq 1 - \nu$, the derivative satisfies

$$G'(P) = \eta_k \gamma_k I(P) e^{-I(P)} (1 - I(P)) \geq m_{\text{loc}} > 0$$

after restricting to the compact region where $I(P)$ is bounded away from zero along the active pressure window. The heterogeneous compact case is the same calculation with the displayed positive lower bound on the pressure derivative. Applying the mean-value theorem to $G(P_i) - G(\bar{P}_{k,\tau})$, using $\sum_i Z_i = 0$, gives

$$\frac{1}{N_k} \sum_i Z_i G(P_i) = \frac{1}{N_k} \sum_i Z_i \{G(P_i) - G(\bar{P}_{k,\tau})\} \geq m_{\text{loc}} \frac{1}{N_k} \sum_i Z_i^2.$$

This is the required pressure-direction marginal-benefit inequality on the local window. The final statement follows by reducing m_{loc} by a known upper bound on second-order feedback errors. $\square$

Remark SM4.13 (Differentiability of the conditional expected terminal cost). The twice Gateaux differentiability assumption is imposed on the conditional expected terminal cost, not on individual sample-path default indicators. Individual default indicators are discontinuous as path functionals, but the regularized killing construction and conditional averaging over the killed-particle law provide the smoothed object used in the variational argument. A sufficient local regime is the following. On the control window, take bounded compact controls, assume the drift and kill intensity are twice continuously differentiable in the state and control variables with bounded derivatives on the relevant compact state region, assume the exponential kill intensity is clipped above by the bounded-killing cap used in the main model, and take a bounded twice continuously differentiable terminal penalty Φ . If the active window stays away from saturation and from degenerate zero-intensity configurations, then the compensator of the Cox killing construction and the conditional killed-particle law are twice Gateaux differentiable with respect to bounded control perturbations by the standard dominated-differentiation argument for SDE coefficients and Cox intensities. The constants in Assumption SM4.9 can then be chosen from the corresponding first- and second-variation bounds. We use this differentiability as a verification-type regularity condition for the information-value theorem; it can degrade in small samples, near control caps, or near degenerate threshold configurations.

Proposition SM4.14 (Well-posedness of the controlled multi-population contagion system). *Assume the following controlled-contagion regularity conditions. More specifically, the drift term*

$$b_k : \mathbb{R} \times \mathcal{M}_{\leq 1}(\mathbb{R})^K \times A_k \rightarrow \mathbb{R}, \quad k = 1, \dots, K,$$

is uniformly Lipschitz in the state and measure variables and satisfies a linear growth condition; the diffusion coefficients σ_k, σ_0 are bounded; the control sets A_k are compact; the contagion matrix

$$\Gamma = (\Gamma_{k\ell})_{1 \leq k, \ell \leq K}$$

is nonnegative and bounded; and the kill intensity $\lambda_k(x, a)$ is nonnegative, bounded, and uniformly Lipschitz in x , namely there exist constants

$$\Lambda < \infty, \quad L_\lambda < \infty,$$

such that for every $x, y \in \mathbb{R}$ and $a \in A_k$,

$$0 \leq \lambda_k(x, a) \leq \Lambda, \quad |\lambda_k(x, a) - \lambda_k(y, a)| \leq L_\lambda |x - y|.$$

Assume that the initial states and noises are mutually independent and satisfy

$$\max_{1 \leq k \leq K} \mathbb{E}|X_1^k(0)|^2 < \infty.$$

Admissible controls in this paper are processes $a = (a^1, \dots, a^K)$ taking values in $A_1 \times \dots \times A_K$ and progressively measurable with respect to the common-noise filtration $\mathbb{F}^0 = (\mathcal{F}_t^{W^0})_{t \in [0, T]}$.

Then the following statements hold.

(i)

For any finite particle size $(N_1, \dots, N_K)$, the controlled multi-population particle system admits a unique strong solution. More precisely, there exists a unique càdlàg process adapted to the filtration generated by the given Brownian motions and Poisson random measures,

$$(X_i^{k,N}(t), \xi_i^{k,N}(t))_{1 \leq i \leq N_k, 1 \leq k \leq K, 0 \leq t \leq T},$$

satisfying

$$dX_i^{k,N}(t) = b_k(X_i^{k,N}(t), \mu_t^{1,N}, \dots, \mu_t^{K,N}, a_t^k) dt + \sigma_k dW_i^k(t) + \sigma_0 dW^0(t) - \sum_{\ell=1}^K \Gamma_{k\ell} dL_t^{\ell,N},$$

where

$$\mu_t^{k,N} = \frac{1}{N_k} \sum_{i=1}^{N_k} \xi_i^{k,N}(t) \delta_{X_i^{k,N}(t)}, \quad L_t^{k,N} = \frac{1}{N_k} \sum_{i=1}^{N_k} (1 - \xi_i^{k,N}(t)).$$

If $\mathcal{N}_i^k(ds, du)$ are independent Poisson random measures on $[0, \infty) \times [0, \Lambda]$ with intensity $ds du$, then the killing indicator is given by

$$\xi_i^{k,N}(t) = 1 - \int_0^t \int_0^\Lambda \xi_i^{k,N}(s-) \mathbf{1}_{\{u \leq \lambda_k(X_i^{k,N}(s-), a_s^k)\}} \mathcal{N}_i^k(ds, du).$$

(ii)

The controlled McKean–Vlasov limiting system admits a unique solution. That is, there exist a unique $\mathbb{F}^0$ -progressively measurable conditional measure flow and loss flow

$$(\mu_t^k, L_t^k)_{1 \leq k \leq K, 0 \leq t \leq T}$$

and representative particle processes $(\bar{X}^k, \bar{\xi}^k)_{1 \leq k \leq K}$ such that

$$d\bar{X}^k(t) = b_k(\bar{X}^k(t), \mu_t^1, \dots, \mu_t^K, a_t^k) dt + \sigma_k dW^k(t) + \sigma_0 dW^0(t) - \sum_{\ell=1}^K \Gamma_{k\ell} dL_t^\ell,$$

and

$$\mu_t^k = \mathbb{E} \left[\bar{\xi}^k(t) \delta_{\bar{X}^k(t)} \mid \mathcal{F}_t^{W^0} \right], \quad L_t^k = 1 - \mathbb{E} \left[\bar{\xi}^k(t) \mid \mathcal{F}_t^{W^0} \right].$$

The killing indicator of the representative particle is defined by

$$\bar{\xi}^k(t) = 1 - \int_0^t \int_0^\Lambda \bar{\xi}^k(s-) \mathbf{1}_{\{u \leq \lambda_k(\bar{X}^k(s-), a_s^k)\}} \mathcal{N}^k(ds, du).$$

(iii)

For each $k = 1, \dots, K$, the limiting loss process L^k is continuous and nondecreasing on $[0, T]$. In fact,

$$L_t^k = \int_0^t \mathbb{E} \left[\lambda_k(\bar{X}^k(s), a_s^k) \bar{\xi}^k(s) \mid \mathcal{F}_s^{W^0} \right] ds, \quad 0 \leq t \leq T.$$

Therefore L^k is absolutely continuous, and

$$0 \leq L_t^k - L_s^k \leq \Lambda(t - s), \quad 0 \leq s \leq t \leq T.$$

The finite-particle loss $L^{k,N}$ is a càdlàg nondecreasing step process.

Proof. We first prove strong well-posedness of the finite-particle system. Fix a particle size $(N_1, \dots, N_K)$, and place all Brownian motions and Poisson random measures on the same complete filtered probability space. Since each particle can be killed at most once and the kill intensity satisfies

$$0 \leq \lambda_k(x, a) \leq \Lambda,$$

the total killing jump intensity in the finite system is uniformly controlled by

$$\Lambda \sum_{k=1}^K N_k.$$

Thus the total number of killing jumps is almost surely finite on any finite time interval. Given a candidate path, the Poisson thinning formula uniquely determines the killing indicators; given the killing indicators, the state equation is an SDE with Lipschitz coefficients in finite dimension and with the finite-variation contagion term

$$- \sum_{\ell=1}^K \Gamma_{k\ell} dL_t^{\ell,N}.$$

The standard piecewise construction gives a unique strong solution between consecutive killing jump times. At a killing jump time, the jump of the loss process uniquely determines the contagion shock to all state variables. Recursively concatenating these intervals gives existence of a strong solution to the finite-particle system.

We now prove pathwise uniqueness. Let

$$(X_i^{k,N}, \xi_i^{k,N}) \quad \text{and} \quad (\tilde{X}_i^{k,N}, \tilde{\xi}_i^{k,N})$$

be two solutions driven by the same initial states, Brownian motions, and Poisson random measures. Denote the corresponding empirical measures and loss processes by

$$(\mu_t^{k,N}, L_t^{k,N}), \quad (\tilde{\mu}_t^{k,N}, \tilde{L}_t^{k,N}).$$

For any test function f satisfying

$$\|f\|_\infty \leq 1, \quad \text{Lip}(f) \leq 1,$$

use the decomposition

$$\begin{aligned} \xi_i^{k,N}(t)f(X_i^{k,N}(t)) - \tilde{\xi}_i^{k,N}(t)f(\tilde{X}_i^{k,N}(t)) &= \xi_i^{k,N}(t)[f(X_i^{k,N}(t)) - f(\tilde{X}_i^{k,N}(t))] \\ &\quad + [\xi_i^{k,N}(t) - \tilde{\xi}_i^{k,N}(t)]f(\tilde{X}_i^{k,N}(t)). \end{aligned}$$

Using $\xi_i^{k,N}(t) \in \{0, 1\}$, $\|f\|_\infty \leq 1$, and $\text{Lip}(f) \leq 1$, we obtain

$$|\xi_i^{k,N}(t)f(X_i^{k,N}(t)) - \tilde{\xi}_i^{k,N}(t)f(\tilde{X}_i^{k,N}(t))| \leq |X_i^{k,N}(t) - \tilde{X}_i^{k,N}(t)| + |\xi_i^{k,N}(t) - \tilde{\xi}_i^{k,N}(t)|.$$

Hence, after taking the supremum over bounded Lipschitz test functions,

$$d_{\text{BL}}(\mu_t^{k,N}, \tilde{\mu}_t^{k,N}) \leq \frac{1}{N_k} \sum_{i=1}^{N_k} |X_i^{k,N}(t) - \tilde{X}_i^{k,N}(t)| + \frac{1}{N_k} \sum_{i=1}^{N_k} |\xi_i^{k,N}(t) - \tilde{\xi}_i^{k,N}(t)|.$$

The second term is the individual alive–dead mismatch. It cannot be replaced by the aggregate loss difference alone, since positive and negative mismatches in the two systems may cancel in the type average. The loss difference is controlled by the same nonnegative mismatch term:

$$|L_t^{k,N} - \tilde{L}_t^{k,N}| \leq \frac{1}{N_k} \sum_{i=1}^{N_k} |\xi_i^{k,N}(t) - \tilde{\xi}_i^{k,N}(t)|.$$

By synchronous Poisson thinning, the difference in killing indicators can only come from the difference between the two kill-intensity paths. Thus

$$\begin{aligned} \mathbb{E} \sup_{r \leq t} |\xi_i^{k,N}(r) - \tilde{\xi}_i^{k,N}(r)| &\leq \mathbb{E} \int_0^t \left| \lambda_k(X_i^{k,N}(s), a_s^k) - \lambda_k(\tilde{X}_i^{k,N}(s), a_s^k) \right| ds \\ &\leq L_\lambda \int_0^t \mathbb{E} |X_i^{k,N}(s) - \tilde{X}_i^{k,N}(s)| ds. \end{aligned}$$

Subtract the two state equations. The Brownian terms cancel, giving

$$\begin{aligned} |X_i^{k,N}(t) - \tilde{X}_i^{k,N}(t)| &\leq \int_0^t \left| b_k(X_i^{k,N}(s), \mu_s^{1,N}, \dots, \mu_s^{K,N}, a_s^k) \right. \\ &\quad \left. - b_k(\tilde{X}_i^{k,N}(s), \tilde{\mu}_s^{1,N}, \dots, \tilde{\mu}_s^{K,N}, a_s^k) \right| ds \\ &\quad + \sum_{\ell=1}^K \Gamma_{k\ell} |L_t^{\ell,N} - \tilde{L}_t^{\ell,N}|. \end{aligned}$$

For each type set

$$A_k(t) := \frac{1}{N_k} \sum_{i=1}^{N_k} \mathbb{E} \sup_{r \leq t} |X_i^{k,N}(r) - \tilde{X}_i^{k,N}(r)|,$$

and

$$B_k(t) := \frac{1}{N_k} \sum_{i=1}^{N_k} \mathbb{E} \sup_{r \leq t} |\xi_i^{k,N}(r) - \tilde{\xi}_i^{k,N}(r)|.$$

The preceding estimates imply

$$B_k(t) \leq L_\lambda \int_0^t A_k(s) ds,$$

while the Lipschitz property of the drift and the boundedness of Γ , together with the bounds on d_{BL} and the loss process above, give

$$A_k(t) \leq C \int_0^t A_k(s) ds + C \int_0^t \sum_{\ell=1}^K (A_\ell(s) + B_\ell(s)) ds + C \sum_{\ell=1}^K B_\ell(t).$$

Let

$$\Delta_N(t) := \sum_{k=1}^K (A_k(t) + B_k(t)).$$

Since $B_k(t) \leq C \int_0^t A_k(s) ds$, summing over k yields

$$\Delta_N(t) \leq C \int_0^t \Delta_N(s) ds.$$

By Gronwall's inequality,

$$\Delta_N(t) = 0, \quad 0 \leq t \leq T.$$

Thus the state processes and killing indicators of the two solutions are indistinguishable. The empirical measures and loss processes are consequently indistinguishable as well, so the finite-particle system has pathwise uniqueness. Combined with the piecewise construction above, this proves existence and uniqueness of a strong solution.

We next prove well-posedness of the McKean–Vlasov limiting system. Let $\mathcal{X}_T$ be the space of all

$$(\boldsymbol{\nu}, \boldsymbol{\ell}) = ((\nu^1, \dots, \nu^K), (\ell^1, \dots, \ell^K))$$

such that ν_t^k is an $\mathcal{M}_{\leq 1}(\mathbb{R})$ -valued $\mathbb{F}^0$ -progressively measurable flow, ℓ_t^k is a $[0, 1]$ -valued $\mathbb{F}^0$ -progressively measurable continuous flow, and

$$\ell_0^k = 0, \quad 0 \leq \ell_s^k \leq \ell_t^k \leq 1, \quad 0 \leq s \leq t \leq T.$$

On $\mathcal{X}_T$, define the seminorm

$$\rho_t((\boldsymbol{\nu}, \boldsymbol{\ell}), (\tilde{\boldsymbol{\nu}}, \tilde{\boldsymbol{\ell}})) := \mathbb{E} \left[\sum_{k=1}^K \sup_{0 \leq s \leq t} d_{\text{BL}}(\nu_s^k, \tilde{\nu}_s^k) + \sum_{k=1}^K \sup_{0 \leq s \leq t} |\ell_s^k - \tilde{\ell}_s^k| \right].$$

The initial second-moment condition ensures that the solution of the linearized equation below has finite second moments, so the conditional measure flow

$$\mathbb{E} [\zeta^k(t) \delta_{Y^k(t)} | \mathcal{F}_t^{W^0}]$$

is well defined and remains in the space of sub-probability measures. Specifically, the linear growth of the drift, $\ell^k \in [0, 1]$, boundedness of Γ , and bounded diffusion coefficients jointly imply

$$\mathbb{E} \sup_{0 \leq t \leq T} |Y^k(t)|^2 \leq C_T \left(1 + \mathbb{E} |X_1^k(0)|^2 \right) < \infty.$$

For any candidate flow $(\boldsymbol{\nu}, \boldsymbol{\ell}) \in \mathcal{X}_T$, define the linearized representative particle by

$$dY^k(t) = b_k(Y^k(t), \nu_t^1, \dots, \nu_t^K, a_t^k) dt + \sigma_k dW^k(t) + \sigma_0 dW^0(t) - \sum_{\ell=1}^K \Gamma_{k\ell} d\ell_t^\ell,$$

and define

$$\zeta^k(t) = 1 - \int_0^t \int_0^\Lambda \zeta^k(s-) \mathbf{1}_{\{u \leq \lambda_k(Y^k(s-), a_s^k)\}} \mathcal{N}^k(ds, du)$$

by Poisson thinning. For a fixed candidate flow, this is an SDE with Lipschitz coefficients and a given finite-variation input, and hence has a unique strong solution. Define the map $\mathcal{T}$ by

$$\mathcal{T}(\boldsymbol{\nu}, \boldsymbol{\ell}) = (\boldsymbol{\mu}, \boldsymbol{L}),$$

where

$$\mu_t^k = \mathbb{E} \left[\zeta^k(t) \delta_{Y^k(t)} \mid \mathcal{F}_t^{W^0} \right], \quad L_t^k = 1 - \mathbb{E} \left[\zeta^k(t) \mid \mathcal{F}_t^{W^0} \right].$$

Since Y^k , ζ^k , and a^k are constructed conditionally on the common noise, μ^k and L^k are $\mathbb{F}^0$ -progressively measurable. By boundedness of the kill intensity and the compensation formula,

$$L_t^k = \int_0^t \mathbb{E} \left[\lambda_k(Y^k(s), a_s^k) \zeta^k(s) \mid \mathcal{F}_s^{W^0} \right] ds,$$

so L^k is continuous, nondecreasing, and satisfies

$$0 \leq L_t^k - L_s^k \leq \Lambda(t - s).$$

Thus $\mathcal{T}$ maps $\mathcal{X}_T$ into itself.

We now prove that $\mathcal{T}$ is a contraction on a short time interval. Take two candidate flows

$$Z = (\boldsymbol{\nu}, \boldsymbol{\ell}), \quad \tilde{Z} = (\tilde{\boldsymbol{\nu}}, \tilde{\boldsymbol{\ell}}),$$

with corresponding linearized solutions

$$(Y^k, \zeta^k, \mu^k, L^k), \quad (\tilde{Y}^k, \tilde{\zeta}^k, \tilde{\mu}^k, \tilde{L}^k).$$

Couple the two systems with the same initial state, Brownian motions, and Poisson random measures. As in the finite-particle part, the state difference satisfies

$$\begin{aligned} \mathbb{E} \sup_{r \leq t} |Y^k(r) - \tilde{Y}^k(r)| &\leq C \int_0^t \mathbb{E} \sup_{u \leq s} |Y^k(u) - \tilde{Y}^k(u)| ds \\ &\quad + C \int_0^t \sum_{\ell=1}^K \mathbb{E} \sup_{u \leq s} d_{\text{BL}}(\nu_u^\ell, \tilde{\nu}_u^\ell) ds + C \sum_{\ell=1}^K \mathbb{E} \sup_{r \leq t} |\ell_r^\ell - \tilde{\ell}_r^\ell|. \end{aligned}$$

Synchronous Poisson thinning and the Lipschitz property of λ_k give

$$\mathbb{E} \sup_{r \leq t} |\zeta^k(r) - \tilde{\zeta}^k(r)| \leq C \int_0^t \mathbb{E} |Y^k(s) - \tilde{Y}^k(s)| ds.$$

Using the d_{BL} decomposition above, the output measure and output loss satisfy

$$\begin{aligned} \sup_{r \leq t} d_{\text{BL}}(\mu_r^k, \tilde{\mu}_r^k) + \sup_{r \leq t} |L_r^k - \tilde{L}_r^k| &\leq C \mathbb{E} \sup_{r \leq t} |Y^k(r) - \tilde{Y}^k(r)| \\ &\quad + C \mathbb{E} \sup_{r \leq t} |\zeta^k(r) - \tilde{\zeta}^k(r)|. \end{aligned}$$

Combining the estimates and summing over k yields

$$R(t) \leq CtD(t) + C \int_0^t R(s) ds,$$

where

$$R(t) := \rho_t(\mathcal{T}Z, \mathcal{T}\tilde{Z}), \quad D(t) := \rho_t(Z, \tilde{Z}).$$

Since $D(s) \leq D(t)$ for $s \leq t$, this is a Volterra-type Gronwall inequality for R . Fixing $t \leq T$ and applying Gronwall's inequality to R , we get

$$R(t) \leq CtD(t) \exp(Ct).$$

Thus

$$\rho_t(\mathcal{T}Z, \mathcal{T}\tilde{Z}) \leq Cte^{Ct} \rho_t(Z, \tilde{Z}).$$

Choose $\delta > 0$ sufficiently small so that

$$C\delta e^{C\delta} < 1.$$

Then $\mathcal{T}$ is a contraction on $[0, \delta]$. Banach's fixed point theorem gives a unique fixed point on $[0, \delta]$. Taking the conditional distribution induced by this fixed point at time δ as the new initial condition, and repeating the same argument on

$$[\delta, 2\delta], \quad [2\delta, 3\delta], \quad \dots,$$

we obtain a unique solution on $[0, T]$ after finitely many concatenations. The short-time contraction constant depends only on

$$T, \quad L_b, \quad L_\lambda, \quad \Lambda, \quad \|\Gamma\|, \quad \max_k \mathbb{E}|X_1^k(0)|^2,$$

and the upper bound on the diffusion coefficients, so the estimates remain uniform at every concatenation step.

Finally, we prove continuity and monotonicity of the limiting loss process. By the fixed point definition and the compensation formula for Poisson thinning,

$$L_t^k = \int_0^t \mathbb{E} \left[\lambda_k(\bar{X}^k(s), a_s^k) \bar{\xi}^k(s) \mid \mathcal{F}_s^{W^0} \right] ds.$$

Since

$$0 \leq \lambda_k(\bar{X}^k(s), a_s^k) \bar{\xi}^k(s) \leq \Lambda,$$

the right-hand side is the time integral of a nonnegative bounded process. Hence L^k is absolutely continuous and therefore continuous. For any $0 \leq s \leq t \leq T$,

$$L_t^k - L_s^k = \int_s^t \mathbb{E} \left[\lambda_k(\bar{X}^k(r), a_r^k) \bar{\xi}^k(r) \mid \mathcal{F}_r^{W^0} \right] dr \geq 0.$$

Also,

$$L_t^k - L_s^k \leq \Lambda(t - s).$$

Thus L^k is continuous and nondecreasing. The finite-particle loss process

$$L_t^{k,N} = \frac{1}{N_k} \sum_{i=1}^{N_k} \mathbf{1}_{\{\tau_i^{k,N} \leq t\}}$$

is the average of finitely many killing indicators, and is therefore a càdlàg nondecreasing step process. The proof is complete. $\square$

SM4.2 Primitive verification of the steep-killing bridge

This subsection gives a primitive verification of the steep-killing bridge. The argument derives eventual entrance of the steep-killing fixed points into a neighborhood of the absorbing fixed point, local stability of the absorbing fixed point, and uniform convergence of the driven steep-killing maps to the absorbing-boundary map from a localized regularity package: compact localization, a uniformly nondegenerate idiosyncratic diffusion coefficient near the boundary, uniform one-dimensional killed-diffusion estimates, and uniqueness of the absorbing-boundary McKean–Vlasov fixed point in the localized class.

Throughout, the common noise is retained. All estimates below may be read conditionally on the common-noise path and then integrated over the common noise. After conditioning on the common noise, the term $\sigma_0 dW_t^0$ is a given continuous forcing path and does not provide conditional smoothing. Hence the nondegeneracy assumption is placed on the idiosyncratic coefficient σ_k , not on $\sigma_k^2 + \sigma_0^2$.

State space, frozen maps, and metrics

Let $\mathbf{E} = \mathbb{R} \cup \{\partial\}$ be the alive–dead state space, where ∂ is a cemetery point. Equip $\mathbf{E}$ with any bounded metric d_∂ that agrees with $|x - y| \wedge 1$ on compact alive-state intervals and puts ∂ at positive distance from the alive region. For finite measures on $\mathbf{E}$, let d_{BL} denote the bounded-Lipschitz distance associated with d_∂ .

A measure–loss input is denoted by

$$z = (\hat{\mu}^1, \dots, \hat{\mu}^K, L^1, \dots, L^K),$$

where $\hat{\mu}_t^k$ is a probability measure on $\mathbf{E}$, its mass at ∂ equals L_t^k , and L^k is nondecreasing with values in $[0, 1]$. For $t \leq T$, set

$$\mathcal{D}_t(z, z') := \sum_{k=1}^K \mathbb{E} \left[\sup_{r \leq t} d_{\text{BL}}(\hat{\mu}_r^k, \hat{\mu}_r'^k) + \sup_{r \leq t} |L_r^k - L_r'^k| \right].$$

The expectation includes the common noise. The full-horizon metric is $\mathcal{D}_T$.

For a frozen input z , the driven type- k state process is written as $X^{k,z}$. In the localized formulation used below, it solves, up to the localization boundary,

$$dX_t^{k,z} = b_k(t, X_t^{k,z}, z_t, a_t^k) dt + \sigma_k dW_t^k + \sigma_0 dW_t^0 - \sum_{\ell=1}^K \Gamma_{k\ell} dL_t^\ell.$$

The coefficients are frozen in z ; default is then generated from the driven path. Let

$$\bar{\lambda}_k(a) := \lambda_{\text{base},k} \exp\{-\eta_k a\}.$$

For $n \geq 1$, define

$$\lambda_k^{(n)}(x, a) := \bar{\lambda}_k(a) \exp\{n(x_{b,k} - x)^+\}.$$

Given a unit exponential random variable E_k , independent of the Brownian noises and common noise, define the steep-killing clock

$$A_t^{k,n,z} := \int_0^t \bar{\lambda}_k(a_s^k) \exp\{n(x_{b,k} - X_s^{k,z})^+\} ds, \quad \tau_k^{n,z} := \inf\{t : A_t^{k,n,z} \geq E_k\}.$$

The baseline Cox clock and absorbing hitting time are

$$A_t^{k,0} := \int_0^t \bar{\lambda}_k(a_s^k) ds, \quad \tau_k^{\text{base}} := \inf\{t : A_t^{k,0} \geq E_k\}, \quad H_k^z := \inf\{t : X_t^{k,z} \leq x_{b,k}\}.$$

The absorbing-limit default time is

$$\tau_k^{\text{abs},z} := \tau_k^{\text{base}} \wedge H_k^z.$$

For $\star \in \{n, \text{abs}\}$, the map $\Phi^\star$ sends z to the lifted conditional laws and losses generated by the same driven state path and the corresponding default time:

$$\hat{\mu}_t^{k,\star,z} := \mathcal{L}(X_t^{k,z} \mathbf{1}_{\{\tau_k^{\star,z} > t\}} + \partial \mathbf{1}_{\{\tau_k^{\star,z} \leq t\}} \mid \mathcal{F}_t^{W^0}), \quad L_t^{k,\star,z} := \mathbb{P}(\tau_k^{\star,z} \leq t \mid \mathcal{F}_t^{W^0}).$$

Thus $\Phi^n(z) = (\hat{\mu}^{1,n,z}, \dots, L^{K,n,z})$ and $\Phi^{\text{abs}}(z) = (\hat{\mu}^{1,\text{abs},z}, \dots, L^{K,\text{abs},z})$.

Primitive regularity package

Assumption SM4.15 (Localized primitive regularity). The following conditions hold on $[0, T]$.

- (P1) **Compact localization.** For each type k , alive states are localized in a compact interval $I_k = [\underline{x}_k, \bar{x}_k]$, with $x_{b,k} \in (\underline{x}_k, \bar{x}_k)$. Equivalently, the original unbounded model is stopped at a large compact interval and all estimates below are applied before the localization error is removed.
- (P2) **Bounded Lipschitz frozen coefficients.** The drift $b_k(t, x, z_t, a)$ is bounded and Lipschitz in (x, z_t) , uniformly in (t, a) . The controls take values in a compact set. There are constants $0 < \underline{\lambda} \leq \bar{\lambda} < \infty$ such that

$$\underline{\lambda} \leq \bar{\lambda}_k(a_t^k) \leq \bar{\lambda} \quad \text{for all } k, t.$$

The lower bound is satisfied by the exponential regulatory-control specification on any compact control set.

- (P3) **Conditional nondegeneracy.** The idiosyncratic volatility is uniformly positive near the boundary:

$$\sigma_k \geq \underline{\sigma} > 0, \quad k = 1, \dots, K.$$

This condition is imposed on σ_k , rather than on $\sigma_k^2 + \sigma_0^2$, because the arguments are conditional on the common noise.

- (P4) **Uniform killed-diffusion estimates.** For every compact class of frozen inputs satisfying the moment and modulus bounds used below, the one-dimensional diffusion killed at $x_{b,k}$ has transition densities and survival probabilities satisfying the standard Aronson and boundary-gradient bounds, uniformly over that class. In particular, for bounded measurable f , the killed semigroup $P_{s,t}^{k,z} f(x) := \mathbb{E}_{s,x}[f(X_t^{k,z}) \mathbf{1}_{\{H_k^z > t\}}]$ satisfies

$$\|\partial_x P_{s,t}^{k,z} f\|_\infty \leq C(t-s)^{-1/2} \|f\|_\infty, \quad 0 \leq s < t \leq T,$$

with a constant C independent of z in the class. The same estimates hold for the baseline-Cox killed semigroup, since the baseline killing rate is bounded.

- (P5) **Uniform absorbing no-atom modulus.** For each compact frozen-input class considered below, the absorbing default times have a uniform distributional modulus: there is a deterministic modulus $\omega_{\text{abs}}(h) \downarrow 0$ as $h \downarrow 0$ such that

$$\sup_z \sup_{0 \leq t \leq t+h \leq T} \mathbb{P}(t < \tau_k^{\text{abs},z} \leq t+h) \leq \omega_{\text{abs}}(h).$$

This condition is required only for the absorbing default time. No uniform-in- n no-atom modulus is imposed on $\tau_k^{n,z}$, since the steep intensity has a Lipschitz and size constant that grows exponentially in n inside the localized interval. Compactness of the steep fixed-point loss paths will instead use monotonicity, Helly selection, and the continuity of the absorbing limiting loss path.

- (P6) **Absorbing fixed-point class.** The localized absorbing-boundary McKean–Vlasov map Φ^{abs} is considered on the compact class generated by (P1)–(P5). Its fixed-point uniqueness is not imposed as a separate black-box assumption; it is verified below from the Volterra stability estimate and a short-time contraction–concatenation argument.

Remark SM4.16 (Why only the absorbing modulus is imposed). The absorbing no-atom modulus in (P5) is the regularity input that controls continuity of the boundary-default map at the level of loss paths. This condition follows from one-dimensional nondegenerate killed-diffusion density estimates and bounded baseline Cox killing. It is imposed only on the absorbing default time and should not be required for $\tau_k^{n,z}$ uniformly in n : on a compact interval the bound on $\lambda_k^{(n)}$ is of order $\exp(Cn)$, so a direct density bound for the steep default time would degenerate with n . The proof below avoids such a false uniform modulus. Uniform convergence of the maps is obtained from the post-hit occupation and clock-mismatch estimate, while compactness of the fixed-point loss paths uses Helly selection and the eventual identification of every limit as the continuous absorbing fixed point.

Step 1: pointwise steep-clock convergence

Lemma SM4.17 (Pointwise convergence of steep killing clocks). *Assume (P1)–(P3). Fix a frozen input z and a type k . If the driven path is nonsticky at $x_{b,k}$, in the sense that on $\{H_k^z < T\}$ it spends positive Lebesgue time below $x_{b,k}$ on every right-neighborhood of H_k^z , then*

$$\tau_k^{n,z} \downarrow \tau_k^{\text{abs},z} \quad a.s.$$

Consequently,

$$\mathbb{E} \sup_{t \leq T} \left| \mathbf{1}_{\{\tau_k^{n,z} \leq t\}} - \mathbf{1}_{\{\tau_k^{\text{abs},z} \leq t\}} \right| \rightarrow 0.$$

Proof. The functions $x \mapsto \exp\{n(x_{b,k} - x)^+\}$ increase pointwise in n , hence $A_t^{k,n,z}$ increases in n for every t , and $\tau_k^{n,z}$ decreases in n . Also $A_t^{k,n,z} \geq A_t^{k,0}$, so $\tau_k^{n,z} \leq \tau_k^{\text{base}}$. The cumulative hazards agree before the boundary is reached: $A_t^{k,n,z} = A_t^{k,0}$ for every $t < H_k^z$, since $(x_{b,k} - X_s^{k,z})^+ = 0$ whenever $X_s^{k,z} > x_{b,k}$. Therefore no steep clock can default before $\tau_k^{\text{base}} \wedge H_k^z$. Thus

$$\tau_k^{\text{abs},z} \leq \tau_k^{n,z} \leq \tau_k^{\text{base}}.$$

If $\tau_k^{\text{base}} < H_k^z$, then the baseline clock rings before the path reaches the boundary, and all clocks give the same default time. If $H_k^z < \tau_k^{\text{base}}$, then after H_k^z the path spends positive Lebesgue time below the boundary on every interval $[H_k^z, H_k^z + \delta]$. Since $\bar{\lambda}_k \geq \underline{\lambda} > 0$, for every $\delta > 0$,

$$\int_{H_k^z}^{(H_k^z + \delta) \wedge T} \bar{\lambda}_k(a_s^k) \exp\{n(x_{b,k} - X_s^{k,z})^+\} ds \rightarrow \infty$$

by monotone convergence on a set of positive Lebesgue measure where $X_s^{k,z} < x_{b,k}$. Hence the steep clock rings before $H_k^z + \delta$ for all sufficiently large n . Letting $\delta \downarrow 0$ gives $\tau_k^{n,z} \downarrow H_k^z = \tau_k^{\text{abs},z}$ on this event. The remaining case $H_k^z = \tau_k^{\text{base}}$ is negligible under the no-atom condition; even without removing it, the monotone bounds force the same limit. Finally,

$$\sup_{t \leq T} \left| \mathbf{1}_{\{\tau_k^{n,z} \leq t\}} - \mathbf{1}_{\{\tau_k^{\text{abs},z} \leq t\}} \right| = \mathbf{1}_{\{\tau_k^{\text{abs},z} < \tau_k^{n,z}\}},$$

and the right-hand side decreases to zero. Dominated convergence completes the proof. $\square$

Step 2: uniform steep-limit continuity of the driven maps

Lemma SM4.18 (Uniform post-hit occupation). *Let $\mathcal{K}$ be a compact frozen-input class satisfying Assumption SM4.15. For each type k , each $h > 0$, and each $\varepsilon > 0$, define*

$$O_{h,\varepsilon}^{k,z} := \int_{H_k^z}^{(H_k^z + h) \wedge T} \mathbf{1}_{\{X_s^{k,z} \leq x_{b,k} - \varepsilon\}} ds.$$

Then, for every $\eta > 0$, there exist $h > 0$, $\varepsilon > 0$, and $q > 0$ such that

$$\sup_{z \in \mathcal{K}} \mathbb{P}(H_k^z < T - h, O_{h,\varepsilon}^{k,z} < q) \leq \eta.$$

Proof. Condition on the common-noise path. On the compact localized state interval, the drift and finite-variation forcing are uniformly bounded in the frozen-input class, and $\sigma_k \geq \underline{\sigma} > 0$. The boundary process therefore has the standard one-dimensional regular-boundary property uniformly over $\mathcal{K}$: after the first entrance into $(-\infty, x_{b,k}]$, the diffusion visits the strict interior $(-\infty, x_{b,k} - \varepsilon]$ during every right-neighborhood with probability tending to one as $\varepsilon \downarrow 0$. The uniform version follows from the boundary Harnack and killed-density estimates in (P4), together with compactness of $\mathcal{K}$. If the displayed quantitative assertion failed, one could find $z_m \in \mathcal{K}$, $h_m \downarrow 0$, and $\varepsilon_m \downarrow 0$ for which the post-hit occupation probability is bounded away from one. Compactness gives a subsequential limit $z_m \rightarrow z$. Stability of the driven SDE gives $X^{k,z_m} \rightarrow X^{k,z}$ uniformly on $[0, T]$ in probability. The regular-boundary property gives the standard hitting-time stability criterion for continuous one-dimensional paths: for every $\delta > 0$, survival strictly before $H_k^z - \delta$ and immediate entrance below the boundary after H_k^z , together with uniform path convergence, imply $\mathbb{P}(H_k^{z_m} < H_k^z - \delta) + \mathbb{P}(H_k^{z_m} > H_k^z + \delta) \rightarrow 0$. The absorbing no-atom modulus in (P5) removes the endpoint ambiguity uniformly. Hence $H_k^{z_m} \rightarrow H_k^z$ in probability. Passing to the limit contradicts the regular-boundary property of the nondegenerate one-dimensional diffusion driven by z . This proves the uniform occupation estimate. $\square$

Lemma SM4.19 (Uniform clock mismatch). *Let $\mathcal{K}$ be a compact frozen-input class satisfying Assumption SM4.15. Then, for every type k ,*

$$\lim_{n \rightarrow \infty} \sup_{z \in \mathcal{K}} \mathbb{E} \sup_{t \leq T} \left| \mathbf{1}_{\{\tau_k^{n,z} \leq t\}} - \mathbf{1}_{\{\tau_k^{\text{abs},z} \leq t\}} \right| = 0.$$

Proof. Since $\tau_k^{\text{abs},z} \leq \tau_k^{n,z}$, the supremum of the two indicator processes equals

$$\mathbf{1}_{\{\tau_k^{\text{abs},z} < \tau_k^{n,z}\}}.$$

Fix $\eta > 0$. Choose $h > 0$ so small that, uniformly in $z \in \mathcal{K}$, the probability that $\tau_k^{\text{abs},z}$ lies within distance h of T is at most η , using the uniform no-atom modulus in (P5). Decrease h

further so that $\mathbb{P}(|\tau_k^{\text{base}} - H_k^z| \leq h) \leq \eta$ uniformly in z . This second bound follows conditionally on the driven path from the independent exponential clock and the bounded baseline hazard: $\mathbb{P}(|\tau_k^{\text{base}} - H_k^z| \leq h \mid X^{k,z}) \leq 2\bar{\lambda}h$. By Lemma SM4.18, after decreasing h if necessary, choose $\varepsilon > 0$ and $q > 0$ such that

$$\sup_{z \in \mathcal{K}} \mathbb{P}(H_k^z < T - h, \int_{H_k^z}^{H_k^z + h} \mathbf{1}_{\{X_s^{k,z} \leq x_{b,k} - \varepsilon\}} ds < q) \leq \eta.$$

On the complementary high-probability event, if the absorbing default is caused by boundary hitting before the baseline Cox clock, then during $[H_k^z, H_k^z + h]$ the steep hazard accumulated below the strict boundary layer is bounded below by

$$\underline{\lambda} q e^{n\varepsilon}.$$

Thus the conditional probability that the same exponential clock has not rung by $H_k^z + h$ is at most

$$\exp\{-\underline{\lambda} q e^{n\varepsilon}\}.$$

If the absorbing default is caused by the baseline Cox clock at least h before the hitting time, all steep clocks default at the same baseline time. The remaining exceptional cases have probability at most a constant multiple of η . Hence

$$\sup_{z \in \mathcal{K}} \mathbb{P}(\tau_k^{\text{abs},z} < \tau_k^{n,z}) \leq C\eta + \exp\{-\underline{\lambda} q e^{n\varepsilon}\}.$$

Letting first $n \rightarrow \infty$ and then $\eta \downarrow 0$ proves the claim. $\square$

Proposition SM4.20 (Uniform steep-limit continuity on compact input classes). *Let $\mathcal{K}$ be a compact frozen-input class satisfying Assumption SM4.15. Then*

$$\lim_{n \rightarrow \infty} \sup_{z \in \mathcal{K}} \mathcal{D}_T(\Phi^n(z), \Phi^{\text{abs}}(z)) = 0.$$

Proof. For a fixed input z , the driven alive-state path is the same under $\Phi^n(z)$ and $\Phi^{\text{abs}}(z)$; only the default time differs. For any bounded-Lipschitz test function f on $\mathbb{E}$,

$$\begin{aligned} & \left| \mathbb{E} \left[f(X_t^{k,z} \mathbf{1}_{\{\tau_k^{n,z} > t\}} + \partial \mathbf{1}_{\{\tau_k^{n,z} \leq t\}}) - f(X_t^{k,z} \mathbf{1}_{\{\tau_k^{\text{abs},z} > t\}} + \partial \mathbf{1}_{\{\tau_k^{\text{abs},z} \leq t\}}) \mid \mathcal{F}_t^{W^0} \right] \right| \\ & \leq 2 \mathbb{P}(\tau_k^{\text{abs},z} \leq t < \tau_k^{n,z} \mid \mathcal{F}_t^{W^0}). \end{aligned}$$

The same bound controls the loss difference. Taking the time supremum and expectation gives

$$\mathcal{D}_T(\Phi^n(z), \Phi^{\text{abs}}(z)) \leq C \sum_{k=1}^K \mathbb{E} \sup_{t \leq T} \left| \mathbf{1}_{\{\tau_k^{n,z} \leq t\}} - \mathbf{1}_{\{\tau_k^{\text{abs},z} \leq t\}} \right|.$$

Lemma SM4.19 gives convergence to zero uniformly over $z \in \mathcal{K}$, and the asserted uniform map convergence follows. $\square$

Remark SM4.21 (Why pointwise convergence is not enough). For a moving sequence of fixed points $z^{(n)}$, pointwise convergence of $\Phi^n(z)$ to $\Phi^{\text{abs}}(z)$ at each fixed z does not imply $\Phi^n(z^{(n)}) - \Phi^{\text{abs}}(z^{(n)}) \rightarrow 0$. The compactness argument and the uniform clock-mismatch estimate above are therefore essential for verifying eventual entrance of the fixed points into a stability neighborhood.

Step 3: local stability of the absorbing-boundary fixed point

Lemma SM4.22 (Causal killed-semigroup stability). *Under Assumption SM4.15, there exist a constant $C < \infty$ and an integrable kernel*

$$g(u) := 1 + u^{-1/2}, \quad 0 < u \leq T,$$

such that for any two frozen inputs z, z' in a sufficiently small compact neighborhood of z^{abs} ,

$$\mathcal{D}_t(\Phi^{\text{abs}}(z), \Phi^{\text{abs}}(z')) \leq C \int_0^t g(t-s) \mathcal{D}_s(z, z') ds, \quad 0 \leq t \leq T.$$

Proof. Condition on the common-noise path. The difference between the two frozen equations enters only through bounded Lipschitz changes in the drift, measure argument, and finite-variation loss feedback. By causality, the output at time t depends on input perturbations only at times $s \leq t$. The killed semigroup representation gives, for bounded test functions f ,

$$P_{0,t}^{k,z} f - P_{0,t}^{k,z'} f = \int_0^t P_{0,s}^{k,z} \left[(B_s^{k,z} - B_s^{k,z'}) \partial_x P_{s,t}^{k,z'} f \right] ds + \mathcal{R}_t,$$

where $B_s^{k,z}$ denotes the frozen drift coefficient, including the absolutely continuous part of the finite-variation forcing after the standard measure approximation, and $\mathcal{R}_t$ is the limit of the corresponding finite-variation forcing term. The approximation is obtained by mollifying each nondecreasing loss path L^ℓ into smooth paths $L^{\ell,\varepsilon}$, applying the standard Duhamel formula to the smooth finite-variation drift, and then letting $\varepsilon \downarrow 0$. Since the absorbing loss paths are continuous under (P5), the mollified paths converge uniformly; the parametric stability of one-dimensional nondegenerate killed diffusions under uniform bounded-variation drift perturbations identifies the limit of the Duhamel terms with the finite-variation forcing term above. The boundary-gradient estimate in (P4) gives

$$\|\partial_x P_{s,t}^{k,z'} f\|_\infty \leq C(t-s)^{-1/2} \|f\|_\infty.$$

The Lipschitz condition on the drift and the contagion-feedback term gives

$$|B_s^{k,z} - B_s^{k,z'}| \leq C \mathcal{D}_s(z, z')$$

after summing over the finitely many types. Hence

$$\sup_{\|f\|_\infty \leq 1} |P_{0,t}^{k,z} f - P_{0,t}^{k,z'} f| \leq C \int_0^t (1 + (t-s)^{-1/2}) \mathcal{D}_s(z, z') ds.$$

The same Duhamel formula applies to the killed survival function by taking $f \equiv 1$ before killing, and therefore controls the loss component. Since the bounded-Lipschitz metric is dominated by the same class of bounded tests plus the Lipschitz part, the estimate also controls the alive-law component. Removing the conditioning on the common noise gives the stated inequality. $\square$

Proposition SM4.23 (Local stability from the Volterra structure). *Under Assumption SM4.15, the absorbing fixed point is locally stable. More precisely, choose a compact reference class $\mathcal{K}_0$ on which the constants in Lemma SM4.22 are uniform, and then choose $r_0 > 0$ so that*

$$\mathcal{U} := \{z \in \mathcal{K}_0 : \mathcal{D}_T(z, z^{\text{abs}}) \leq r_0\}$$

remains inside that class. There is a constant $C_{\text{st}} < \infty$ such that, for every $z \in \mathcal{U}$,

$$\mathcal{D}_T(z, z^{\text{abs}}) \leq C_{\text{st}} \mathcal{D}_T(z, \Phi^{\text{abs}}(z)).$$

Proof. Let

$$e(t) := \mathcal{D}_t(z, z^{\text{abs}}), \quad r(t) := \mathcal{D}_t(z, \Phi^{\text{abs}}(z)).$$

Since $z^{\text{abs}} = \Phi^{\text{abs}}(z^{\text{abs}})$, Lemma SM4.22 gives

$$\begin{aligned} e(t) &\leq r(t) + \mathcal{D}_t(\Phi^{\text{abs}}(z), \Phi^{\text{abs}}(z^{\text{abs}})) \\ &\leq r(t) + C \int_0^t g(t-s)e(s) ds. \end{aligned}$$

The kernel $g(u) = 1 + u^{-1/2}$ belongs to $L^1(0, T)$. The standard resolvent form of the Volterra–Gronwall inequality therefore yields a nonnegative resolvent kernel $G \in L^1(0, T)$, depending only on C, g, T , such that

$$e(t) \leq r(t) + \int_0^t G(t-s)r(s) ds.$$

Since $r(s) \leq r(T)$,

$$e(T) \leq \left(1 + \|G\|_{L^1(0, T)}\right) r(T).$$

Thus the desired estimate holds with

$$C_{\text{st}} := 1 + \|G\|_{L^1(0, T)}.$$

This proof is the nonlinear Volterra version of the spectral-radius argument: the linearized absorbing map is a causal Volterra operator with an integrable kernel, and hence its spectrum is $\{0\}$; equivalently, $I - D\Phi^{\text{abs}}(z^{\text{abs}})$ has a bounded resolvent. The resolvent proof avoids requiring an explicit Fréchet-derivative formula for the hitting-time functional. $\square$

Proposition SM4.24 (Localized uniqueness of the absorbing fixed point). *Under Assumption SM4.15, the absorbing-boundary McKean–Vlasov map Φ^{abs} has at most one fixed point in the localized compact class. Hence the absorbing fixed point z^{abs} , whenever it exists in that class, is unique.*

Proof. Let z and $\tilde{z}$ be two localized absorbing fixed points. On a time interval $[0, t]$, Lemma SM4.22 gives

$$\mathcal{D}_t(z, \tilde{z}) = \mathcal{D}_t(\Phi^{\text{abs}}(z), \Phi^{\text{abs}}(\tilde{z})) \leq C \int_0^t (1 + (t-s)^{-1/2}) \mathcal{D}_s(z, \tilde{z}) ds.$$

Choose $\delta > 0$ such that

$$C \int_0^\delta (1 + u^{-1/2}) du < 1.$$

Taking the supremum over $s \leq t \leq \delta$ gives

$$\sup_{s \leq \delta} \mathcal{D}_s(z, \tilde{z}) \leq \theta \sup_{s \leq \delta} \mathcal{D}_s(z, \tilde{z}), \quad \theta < 1,$$

so $z = \tilde{z}$ on $[0, \delta]$. Suppose uniqueness has been shown up to a deterministic time t_0 . Restart the absorbing driven equations from the common conditional law and common loss value at t_0 . The coefficients after t_0 satisfy the same localized bounds, and the finite-variation contagion input up to t_0 is already fixed. Applying the same short-time estimate on $[t_0, t_0 + \delta]$ yields equality on that interval. Finitely many concatenations cover $[0, T]$. This proves localized uniqueness. $\square$

Step 4: compactness and eventual entrance of steep fixed points

Lemma SM4.25 (Compactness of the steep fixed-point family). *Assume the primitive package and let $z^{(n)}$ be fixed points of Φ^n . Then every sequence of steep fixed points has a subsequence, still denoted by $z^{(n_j)}$, with the following compactness properties. The loss paths converge by Helly selection at all continuity points of nondecreasing limits $L^{k,*}$, and the lifted laws are tight in the bounded-Lipschitz path topology after this Helly extraction. If the resulting limiting loss paths are continuous, then the Helly convergence of losses is uniform on $[0, T]$, and the corresponding subsequence converges in $\mathcal{D}_T$. In particular, once the limit is identified as the continuous absorbing fixed point, the convergence is in the full $\mathcal{D}_T$ metric.*

Proof. Under compact localization the second-moment bound is immediate. In the unlocalized formulation, apply Itô's formula to $|X_t^{k,(n)}|^2$ before localization. The drift has at most linear growth with constants independent of n , the diffusion coefficient is constant, and killing only moves mass to the cemetery point. Gronwall's inequality gives

$$\sup_{n \geq 1} \sup_{t \leq T} \mathbb{E} |X_t^{k,(n)}|^2 < \infty.$$

The losses satisfy $0 \leq L_t^{k,(n)} \leq 1$ and are nondecreasing. We do not use a uniform-in- n no-atom modulus for the steep default times. Instead, Helly's selection theorem gives, for every sequence, a subsequence such that each loss path $L^{k,(n_j)}$ converges pointwise at all continuity points of a nondecreasing limit $L^{k,*}$. This supplies the loss compactness in the natural Helly topology. If a later identification step shows that $L^{k,*}$ is continuous, then Pólya's theorem for monotone functions upgrades the pointwise convergence to uniform convergence on $[0, T]$, and bounded convergence gives

$$\mathbb{E} \sup_{t \leq T} |L_t^{k,(n_j)} - L_t^{k,*}| \longrightarrow 0.$$

For the alive-state component, use tightness at fixed times and the SDE increment estimate away from the jump into the cemetery state. Brownian increments contribute order $h^{1/2}$, bounded drift contributes order h , and the finite-variation contagion feedback is controlled along convergent subsequences by the Helly convergence of the loss paths. The only discontinuity of the lifted state is the cemetery jump, whose contribution to the bounded-Lipschitz modulus is exactly controlled by the loss increment. Hence, after passing to a Helly subsequence for the losses, the lifted laws are tight in the bounded-Lipschitz path topology. Prokhorov compactness at fixed times follows from the uniform second-moment bound, and the preceding increment estimate gives the required equicontinuity on every subsequence whose losses have a continuous limit. This proves the stated subsequential compactness, and the final assertion follows from the Helly–Pólya upgrade when the limit is continuous. $\square$

Proposition SM4.26 (Eventual entrance into the stability neighborhood). *Assume the primitive package. Let $\mathcal{U}$ be the compact stability neighborhood from Proposition SM4.23. Then*

$$z^{(n)} \in \mathcal{U} \quad \text{for all sufficiently large } n.$$

In fact, $\mathcal{D}_T(z^{(n)}, z^{\text{abs}}) \rightarrow 0$.

Proof. By Lemma SM4.25, every subsequence of $\{z^{(n)}\}$ has a further subsequence, still denoted by $z^{(n_j)}$, converging in the Helly–weak path topology to some candidate limit $z^* \in \mathcal{K}_0$. The fixed-point

identity $z^{(n_j)} = \Phi^{n_j}(z^{(n_j)})$, the uniform clock-mismatch estimate on the compact class $\mathcal{K}_0$, and the parametric stability of the driven absorbing map along Helly-convergent finite-variation inputs imply

$$z^* = \Phi^{\text{abs}}(z^*)$$

in the same weak path topology. Equivalently, for every bounded Lipschitz test function and every continuity time of the limiting loss paths, the limiting lifted law and loss satisfy the absorbing driven equations. By Proposition SM4.24, $z^* = z^{\text{abs}}$.

The limit losses are therefore the absorbing fixed-point loss paths, which are continuous by (P5). Pólya's theorem upgrades the Helly convergence of $L^{k,(n_j)}$ to uniform convergence on $[0, T]$; the lifted-law convergence then upgrades to $\mathcal{D}_T$ by the SDE stability and the cemetery-jump control in Lemma SM4.25. Thus every subsequence has a further subsequence converging to z^{abs} in $\mathcal{D}_T$, and consequently the whole sequence converges to z^{abs} in $\mathcal{D}_T$. Any compact neighborhood $\mathcal{U}$ of z^{abs} inside the localized class contains all sufficiently large $z^{(n)}$. $\square$

Primitive version of Theorem SM4.16

Theorem SM4.27 (Steep-killing bridge with primitive verification). *Assume the localized primitive regularity package in Assumption SM4.15. Let $z^{(n)}$ be any sequence of McKean–Vlasov fixed points generated by the steep intensities*

$$\lambda_k^{(n)}(x, a) = \bar{\lambda}_k(a) \exp\{n(x_{b,k} - x)^+\}.$$

Let z^{abs} be the unique localized absorbing-boundary McKean–Vlasov fixed point with default time

$$\tau_k^{\text{abs}} = \tau_k^{\text{base}} \wedge H_k, \quad H_k = \inf\{t : X_t^k \leq x_{b,k}\}.$$

Then the bridge conditions are verified as follows.

First, there exists a compact stability neighborhood $\mathcal{U}$ of z^{abs} such that $z^{(n)} \in \mathcal{U}$ for all sufficiently large n . Second, the absorbing fixed point is locally stable:

$$\mathcal{D}_T(z, z^{\text{abs}}) \leq C_{\text{st}} \mathcal{D}_T(z, \Phi^{\text{abs}}(z)), \quad z \in \mathcal{U}.$$

Third, the driven maps converge uniformly on $\mathcal{U}$:

$$\lim_{n \rightarrow \infty} \sup_{z \in \mathcal{U}} \mathcal{D}_T(\Phi^n(z), \Phi^{\text{abs}}(z)) = 0.$$

Consequently,

$$\mathcal{D}_T(z^{(n)}, z^{\text{abs}}) \rightarrow 0.$$

Equivalently, for every type k ,

$$\mathbb{E} \sup_{t \leq T} d_{\text{BL}}(\hat{\mu}_t^{k,(n)}, \hat{\mu}_t^{k,\text{abs}}) \rightarrow 0, \quad \mathbb{E} \sup_{t \leq T} |L_t^{k,(n)} - L_t^{k,\text{abs}}| \rightarrow 0.$$

Proof. Uniform map convergence on compact classes is Proposition SM4.20. Local stability is Proposition SM4.23. Eventual entrance is Proposition SM4.26. For completeness, once $z^{(n)} \in \mathcal{U}$, the fixed-point residual estimate gives

$$\begin{aligned} \mathcal{D}_T(z^{(n)}, z^{\text{abs}}) &\leq C_{\text{st}} \mathcal{D}_T(z^{(n)}, \Phi^{\text{abs}}(z^{(n)})) \\ &= C_{\text{st}} \mathcal{D}_T(\Phi^n(z^{(n)}), \Phi^{\text{abs}}(z^{(n)})) \\ &\leq C_{\text{st}} \sup_{z \in \mathcal{U}} \mathcal{D}_T(\Phi^n(z), \Phi^{\text{abs}}(z)) \rightarrow 0. \end{aligned}$$

This proves the steep-killing bridge at the solution level. $\square$

Remark SM4.28 (Pure absorbing boundary). The theorem above corresponds to the intensity $\bar{\lambda}_k(a) \exp\{n(x_{b,k} - x)^+\}$, whose limit keeps the baseline Cox default mechanism before boundary hitting. A pure absorbing-boundary limit with no above-boundary Cox killing is obtained by replacing the intensity by

$$\tilde{\lambda}_k^{(n)}(x, a) := \bar{\lambda}_k(a)(\exp\{n(x_{b,k} - x)^+\} - 1).$$

The same proof applies, with $\tau_k^{\text{abs}} = H_k$, provided the lower-bound part of (P2) is imposed only below the boundary layer where the singular killing is active and the uniform no-atom modulus is stated for the hitting time H_k .

References

- [1] L. EISENBERG AND T. H. NOE, *Systemic risk in financial systems*, Manage. Sci., 47 (2001), pp. 236–249, <https://doi.org/10.1287/mnsc.47.2.236.9835>.
- [2] F. ALLEN AND D. GALE, *Financial contagion*, J. Polit. Econ., 108 (2000), pp. 1–33, <https://doi.org/10.1086/262109>.
- [3] P. GAI AND S. KAPADIA, *Contagion in financial networks*, Proc. R. Soc. A, 466 (2010), pp. 2401–2423, <https://doi.org/10.1098/rspa.2009.0410>.
- [4] D. ACEMOGLU, A. OZDAGLAR, AND A. TAHBAZ-SALEHI, *Systemic risk and stability in financial networks*, Amer. Econ. Rev., 105 (2015), pp. 564–608, <https://doi.org/10.1257/aer.20130456>.
- [5] M. ELLIOTT, B. GOLUB, AND M. O. JACKSON, *Financial networks and contagion*, Amer. Econ. Rev., 104 (2014), pp. 3115–3153, <https://doi.org/10.1257/aer.104.10.3115>.
- [6] P. GLASSERMAN AND H. P. YOUNG, *Contagion in financial networks*, J. Econ. Lit., 54 (2016), pp. 779–831, <https://doi.org/10.1257/jel.20151228>.
- [7] B. HAMBLY, S. LEDGER, AND A. SØJMARK, *A McKean–Vlasov equation with positive feedback and blow-ups*, Ann. Appl. Probab., 29 (2019), pp. 2338–2373, <https://doi.org/10.1214/18-AAP1455>.
- [8] B. HAMBLY AND A. SØJMARK, *An SPDE model for systemic risk with endogenous contagion*, Finance Stoch., 23 (2019), pp. 535–594, <https://doi.org/10.1007/s00780-019-00396-1>.
- [9] B. HAMBLY AND P. JETTKANT, *Control of McKean–Vlasov SDEs with contagion through killing at a state-dependent intensity*, Ann. Appl. Probab., 35 (2025), pp. 4381–4438, <https://doi.org/10.1214/25-AAP2226>.
- [10] V. V. ACHARYA, L. H. PEDERSEN, T. PHILIPPON, AND M. RICHARDSON, *Measuring systemic risk*, Rev. Financ. Stud., 30 (2017), pp. 2–47, <https://doi.org/10.1093/rfs/hhw088>.
- [11] T. ADRIAN AND M. K. BRUNNERMEIER, *CoVaR*, Amer. Econ. Rev., 106 (2016), pp. 1705–1741, <https://doi.org/10.1257/aer.20120555>.
- [12] L. C. G. ROGERS AND L. A. M. VERAART, *Failure and rescue in an interbank network*, Manage. Sci., 59 (2013), pp. 882–898, <https://doi.org/10.1287/mnsc.1120.1569>.

- [13] R. CIFUENTES, G. FERRUCCI, AND H. S. SHIN, *Liquidity risk and contagion*, J. Eur. Econom. Assoc., 3 (2005), pp. 556–566, <https://doi.org/10.1162/jeea.2005.3.2-3.556>.
- [14] P. GAI, A. HALDANE, AND S. KAPADIA, *Complexity, concentration and contagion*, J. Monetary Econ., 58 (2011), pp. 453–470, <https://doi.org/10.1016/j.jmoneco.2011.05.005>.
- [15] R. J. CABALLERO AND A. SIMSEK, *Fire sales in a model of complexity*, J. Finance, 68 (2013), pp. 2549–2587, <https://doi.org/10.1111/jofi.12087>.
- [16] R. M. MAY AND N. ARINAMINPATHY, *Systemic risk: The dynamics of model banking systems*, J. R. Soc. Interface, 7 (2010), pp. 823–838, <https://doi.org/10.1098/rsif.2009.0359>.
- [17] M. BOSS, H. ELSINGER, M. SUMMER, AND S. THURNER, *Network topology of the interbank market*, Quant. Finance, 4 (2004), pp. 677–684, <https://doi.org/10.1080/14697680400020325>.
- [18] E. NIER, J. YANG, T. YORULMAZER, AND A. ALENTORN, *Network models and financial stability*, J. Econom. Dynam. Control, 31 (2007), pp. 2033–2060, <https://doi.org/10.1016/j.jedc.2007.01.014>.
- [19] R. CONT, A. MOUSSA, AND E. B. SANTOS, *Network structure and systemic risk in banking systems*, in Handbook on Systemic Risk, J.-P. Fouque and J. A. Langsam, eds., Cambridge University Press, 2013, pp. 327–368, <https://doi.org/10.1017/CB09781139151184.018>.
- [20] S. BATTISTON, D. DELLI GATTI, M. GALLEGATI, B. GREENWALD, AND J. E. STIGLITZ, *Liaisons dangereuses: Increasing connectivity, risk sharing, and systemic risk*, J. Econom. Dynam. Control, 36 (2012), pp. 1121–1141, <https://doi.org/10.1016/j.jedc.2012.04.001>.
- [21] P. GLASSERMAN AND H. P. YOUNG, *How likely is contagion in financial networks?*, J. Banking Finance, 50 (2015), pp. 383–399, <https://doi.org/10.1016/j.jbankfin.2014.02.006>.
- [22] F. CACCIOLI, J. D. FARMER, N. FOTI, AND D. ROCKMORE, *Overlapping portfolios, contagion, and financial stability*, J. Econom. Dynam. Control, 51 (2015), pp. 50–63, <https://doi.org/10.1016/j.jedc.2014.09.041>.
- [23] A. CAPPONI, P.-C. CHEN, AND D. D. YAO, *Liability concentration and systemic losses in financial networks*, Oper. Res., 64 (2016), pp. 1121–1134, <https://doi.org/10.1287/opre.2015.1402>.
- [24] A. GANDY AND L. A. M. VERAART, *A bayesian methodology for systemic risk assessment in financial networks*, Manage. Sci., 63 (2017), pp. 4428–4446, <https://doi.org/10.1287/mnsc.2016.2546>.
- [25] M. DEMIRER, F. X. DIEBOLD, L. LIU, AND K. YILMAZ, *Estimating global bank network connectedness*, J. Appl. Econometrics, 33 (2018), pp. 1–15, <https://doi.org/10.1002/jae.2585>.
- [26] H. AMINI, R. CONT, AND A. MINCA, *Resilience to contagion in financial networks*, Math. Finance, 26 (2016), pp. 329–365, <https://doi.org/10.1111/mafi.12051>.
- [27] J. P. GLEESON, *High-accuracy approximation of binary-state dynamics on networks*, Phys. Rev. Lett., 107 (2011), p. 068701, <https://doi.org/10.1103/PhysRevLett.107.068701>.

- [28] T. R. HURD, J. P. GLEESON, AND S. MELNIK, *A framework for analyzing contagion in assortative banking networks*, PLoS ONE, 12 (2017), p. e0170579, <https://doi.org/10.1371/journal.pone.0170579>.
- [29] J.-P. FOUQUE AND J. A. LANGSAM, eds., *Handbook on Systemic Risk*, Cambridge University Press, 2013, <https://doi.org/10.1017/CB09781139151184>.
- [30] J.-P. FOUQUE AND T. ICHIBA, *Stability in a model of interbank lending*, SIAM J. Financial Math., 4 (2013), pp. 784–803, <https://doi.org/10.1137/110841096>.
- [31] J. GARNIER, G. PAPANICOLAOU, AND T.-W. YANG, *Large deviations for a mean field model of systemic risk*, SIAM J. Financial Math., 4 (2013), pp. 151–184, <https://doi.org/10.1137/12087387X>.
- [32] S. LEDGER AND A. SØJMARK, *Uniqueness for contagious McKean–Vlasov systems in the weak feedback regime*, Bull. London Math. Soc., 52 (2020), pp. 448–463, <https://doi.org/10.1112/blms.12337>.
- [33] F. DELARUE, J. INGLIS, S. RUBENTHALER, AND E. TANRÉ, *Particle systems with a singular mean-field self-excitation. application to neuronal networks*, Stochastic Process. Appl., 125 (2015), pp. 2451–2492, <https://doi.org/10.1016/j.spa.2015.01.007>.
- [34] A.-S. SZNITMAN, *Topics in propagation of chaos*, in École d’Été de Probabilités de Saint-Flour XIX—1989, vol. 1464 of Lecture Notes in Mathematics, Springer, Berlin, 1991, pp. 165–251, <https://doi.org/10.1007/BFb0085169>.
- [35] A. BUDHIRAJA, P. DUPUIS, AND M. FISCHER, *Large deviation properties of weakly interacting processes via weak convergence methods*, Ann. Probab., 40 (2012), pp. 74–102, <https://doi.org/10.1214/10-AOP616>.
- [36] Z. FEINSTEIN AND A. SØJMARK, *Dynamic default contagion in heterogeneous interbank systems*, SIAM J. Financial Math., 12 (2021), pp. SC83–SC97, <https://doi.org/10.1137/20M1376765>.
- [37] Z. FEINSTEIN AND A. SØJMARK, *Contagious McKean–Vlasov systems with heterogeneous impact and exposure*, Finance Stoch., 27 (2023), pp. 663–711, <https://doi.org/10.1007/s00780-023-00504-2>.
- [38] M. HUANG, R. P. MALHAMÉ, AND P. E. CAINES, *Large population stochastic dynamic games: Closed-loop McKean–Vlasov systems and the Nash certainty equivalence principle*, Commun. Inf. Syst., 6 (2006), pp. 221–252, <https://doi.org/10.4310/CIS.2006.v6.n3.a5>.
- [39] R. A. CARMONA, J.-P. FOUQUE, AND L.-H. SUN, *Mean field games and systemic risk*, Commun. Math. Sci., 13 (2015), pp. 911–933, <https://doi.org/10.4310/CMS.2015.v13.n4.a4>.
- [40] R. CARMONA AND F. DELARUE, *Probabilistic Theory of Mean Field Games with Applications I–II*, Springer, 2018.
- [41] G. FABBRI, F. GOZZI, AND A. SWIECH, *Stochastic Optimal Control in Infinite Dimension*, Springer, 2017, <https://doi.org/10.1007/978-3-319-53067-3>.
- [42] M. G. CRANDALL, H. ISHII, AND P.-L. LIONS, *User’s guide to viscosity solutions of second order partial differential equations*, Bull. Amer. Math. Soc., 27 (1992), pp. 1–67, <https://doi.org/10.1090/S0273-0979-1992-00266-5>.

- [43] A. COSSO, F. GOZZI, I. KHARROUBI, H. PHAM, AND M. ROSESTOLATO, *Master bellman equation in the Wasserstein space: Uniqueness of viscosity solutions*, Trans. Amer. Math. Soc., 377 (2024), pp. 31–83, <https://doi.org/10.1090/tran/8986>.
- [44] W. GANGBO, A. R. MÉSZÁROS, C. MOU, AND J. ZHANG, *Mean field games master equations with nonseparable hamiltonians and displacement monotonicity*, Ann. Probab., 50 (2022), pp. 2178–2217, <https://doi.org/10.1214/22-AOP1580>.
- [45] J.-F. CHASSAGNEUX, D. CRISAN, AND F. DELARUE, *Numerical method for FBSDEs of McKean–Vlasov type*, Ann. Appl. Probab., 29 (2019), pp. 1640–1684, <https://doi.org/10.1214/18-AAP1429>.
- [46] J. HAN, A. JENTZEN, AND W. E, *Solving high-dimensional partial differential equations using deep learning*, Proc. Natl. Acad. Sci. USA, 115 (2018), pp. 8505–8510, <https://doi.org/10.1073/pnas.1718942115>.
- [47] L. LOVASZ, *Large Networks and Graph Limits*, American Mathematical Society, 2012.
- [48] I. BENJAMINI AND O. SCHRAMM, *Recurrence of distributional limits of finite planar graphs*, Electron. J. Probab., 6 (2001), pp. 1–13, <https://doi.org/10.1214/EJP.v6-96>.
- [49] C. BORGS, J. T. CHAYES, H. COHN, AND N. HOLDEN, *Sparse exchangeable graphs and their limits via Graphon processes*, J. Mach. Learn. Res., 18 (2018), pp. 1–71. JMLR article 210; available at <https://jmlr.org/papers/v18/16-421.html>.
- [50] S. DELATTRE, G. GIACOMIN, AND E. LUÇON, *A note on dynamical models on random graphs and Fokker–Planck equations*, J. Stat. Phys., 165 (2016), pp. 785–798, <https://doi.org/10.1007/s10955-016-1652-3>.
- [51] R. I. OLIVEIRA AND G. H. REIS, *Interacting diffusions on random graphs with diverging average degrees: Hydrodynamics and large deviations*, J. Stat. Phys., 176 (2019), pp. 1057–1087, <https://doi.org/10.1007/s10955-019-02332-1>.
- [52] R. I. OLIVEIRA, G. H. REIS, AND L. M. STOLERMAN, *Interacting diffusions on sparse graphs: Hydrodynamics from local weak limits*, Electron. J. Probab., 25 (2020), pp. 1–35, <https://doi.org/10.1214/20-EJP505>.
- [53] F. PARISE AND A. OZDAGLAR, *Graphon games: A statistical framework for network games and interventions*, Econometrica, 91 (2023), pp. 191–225, <https://doi.org/10.3982/ECTA17564>.
- [54] P. E. CAINES AND M. HUANG, *Graphon mean field games and their equations*, SIAM J. Control Optim., 59 (2021), pp. 4373–4399, <https://doi.org/10.1137/20M136373X>.
- [55] R. CARMONA, D. B. COONEY, C. V. GRAVES, AND M. LAURIÈRE, *Stochastic Graphon games: I. the static case*, Math. Oper. Res., 47 (2022), pp. 750–778, <https://doi.org/10.1287/moor.2021.1148>.
- [56] A. AURELL, R. CARMONA, G. DAYANIKLI, AND M. LAURIÈRE, *Finite state Graphon games with applications to epidemics*, Dyn. Games Appl., 12 (2022), pp. 49–81, <https://doi.org/10.1007/s13235-021-00410-2>.
- [57] E. BAYRAKTAR, S. CHAKRABORTY, AND R. WU, *Graphon mean field systems*, Ann. Appl. Probab., 33 (2023), pp. 3587–3619, <https://doi.org/10.1214/22-AAP1901>.

- [58] D. LACKER AND A. SORET, *A case study on stochastic games on large graphs in mean field and sparse regimes*, Math. Oper. Res., 47 (2022), pp. 1530–1565, <https://doi.org/10.1287/moor.2021.1179>.
- [59] R. J. BENDEL, *Sharp transitions and systemic risk in sparse financial networks*, arXiv preprint arXiv:2601.04096, (2026), <https://doi.org/10.48550/arXiv.2601.04096>.
- [60] EUROPEAN BANKING AUTHORITY, *2024 eu-wide transparency exercise*, 2024. Public data release, <https://www.eba.europa.eu/risk-and-data-analysis/risk-analysis/eu-wide-transparency-exercise/2024-eu-wide-transparency-exercise>.
- [61] EUROPEAN BANKING AUTHORITY, *Pillar 3 data hub*, 2026. European Data Access Portal disclosure platform, <https://www.eba.europa.eu/risk-and-data-analysis/pillar-3-data-hub>.
- [62] NATURAL EARTH, *Natural earth vector map data*, 2024. Public-domain map data, <https://www.naturalearthdata.com/>.
- [63] A. W. VAN DER VAART AND J. A. WELLNER, *Weak Convergence and Empirical Processes: With Applications to Statistics*, Springer Series in Statistics, Springer, New York, 1996, <https://doi.org/10.1007/978-1-4757-2545-2>.