

Lipschitzian SLLNs for random functions

Lai Tian*

Johannes O. Royset*

July 22, 2026

Abstract

We prove strong laws of large numbers for locally Lipschitz functions in the Lipschitz pseudometric. Our results hold under either a topological or a model-theoretic condition, with the latter encompassing functions jointly definable in o-minimal structures but extending substantially beyond this class. Applications include uniform convergence of limiting and Clarke subdifferentials and finite-sample identification of solutions. Consequently, we identify broad classes of functions for which the failure phenomena revealed by our previous negative results [Tian and Royset, arXiv:2511.16568, 2025] do not occur.

1 Introduction

In stochastic optimization and statistical learning, empirical approximations are expected to preserve the landscape of the population problem. This principle is well understood for global solutions and, in nonconvex smooth settings, for stationary points. Our goal is to develop an analogous consistency theory for stationary points of nonconvex nonsmooth problems.

Let $(\Xi, \mathcal{A})$ be a measurable space and let $\xi, \xi^1, \xi^2, \dots$ be independent and identically distributed (iid) Ξ -valued random variables on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Throughout, we use boldface notation for random variables. For each $\nu \in \mathbb{N} = \{1, 2, \dots\}$, consider the problems

$$\text{minimize}_{x \in X} Ef(x) = \mathbb{E}[f(\xi, x)], \quad (1)$$

$$\text{minimize}_{x \in X} E^\nu f(x) = \frac{1}{\nu} \sum_{i=1}^{\nu} f(\xi^i, x). \quad (2)$$

Here, $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ is $\mathcal{A}$ -measurable in ξ for every $x \in \mathbb{R}^d$, and each slice $f(\xi, \cdot)$ is a possibly nonconvex and nonsmooth Lipschitz function on $X \subset \mathbb{R}^d$. We refer to (1) as the *population* problem and to (2) as its *sample-average approximation* (SAA) problem. Epigraphical and uniform strong laws of large numbers (SLLNs) establish consistency of global solutions for the SAA problem; see, e.g., [3] and [49, Section 9.2.6]. For stationary points, uniform laws for gradients yield an analogous consistency theory in the nonconvex smooth setting; see, e.g., [49, p. 450] and [38, 24]. When the function $f(\xi, \cdot)$ may be nonconvex and nonsmooth, however, even the appropriate formulation of consistency concepts for stationary points requires care.

For ease of exposition, assume temporarily that $X = \mathbb{R}^d$ and consider stationarity defined through the Clarke subdifferential [42, Equation 8(32)]. Whenever well-defined, a point $x \in \mathbb{R}^d$

*Daniel J. Epstein Department of Industrial and Systems Engineering, University of Southern California, Los Angeles, CA. Emails: {laitian, royset}@usc.edu

is Clarke stationary for the population problem if $0 \in \bar{\partial}Ef(x)$; the analogous definition applies to the SAA problem. The desired conclusion is that, if $\mathbb{P}$ -almost surely ($\mathbb{P}$ -a.s.), $\mathbf{x}^\nu \rightarrow \mathbf{x}$ and $\text{dist}(0, \bar{\partial}E^\nu f(\mathbf{x}^\nu)) \rightarrow 0$, then $0 \in \bar{\partial}Ef(\mathbf{x})$, $\mathbb{P}$ -a.s. However, most existing consistency results do not handle the two set-valued mappings $\bar{\partial}E^\nu f$ and $\bar{\partial}Ef$; see, e.g., [47, 60, 40, 13]. Instead, these works interchange expectation and subdifferentiation and study $\frac{1}{\nu} \sum_{i=1}^\nu \bar{\partial}_x f(\boldsymbol{\xi}^i, \cdot)$ and $\mathbb{E}[\bar{\partial}_x f(\boldsymbol{\xi}, \cdot)]$, where $\bar{\partial}_x f(\boldsymbol{\xi}, \cdot) = \bar{\partial}(f(\boldsymbol{\xi}, \cdot))$ denotes the partial Clarke subdifferential and the expectation is understood in the Aumann sense [4]. The solution concept defined by the zero sets of $\frac{1}{\nu} \sum_{i=1}^\nu \bar{\partial}_x f(\boldsymbol{\xi}^i, \cdot)$ and $\mathbb{E}[\bar{\partial}_x f(\boldsymbol{\xi}, \cdot)]$ is commonly referred to as *weak stationarity*; see, e.g., [60, 13]. Consistency of weak stationarity typically holds under mild assumptions [47, 40, 60]. By standard interchange rules [16, Theorem 2.7.2], one has $\bar{\partial}E^\nu f(x) \subset \frac{1}{\nu} \sum_{i=1}^\nu \bar{\partial}_x f(\boldsymbol{\xi}^i, x)$ and $\bar{\partial}Ef(x) \subset \mathbb{E}[\bar{\partial}_x f(\boldsymbol{\xi}, x)]$, while the reverse inclusions generally fail unless suitable subdifferential regularity conditions hold. Thus, *stationarity*, defined through the subdifferential of the expectation function, is a sharper solution concept than weak stationarity.

Indeed, this distinction between weak stationarity and stationarity can be decisive. For illustration, let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be a 1-Lipschitz function satisfying $\bar{\partial}\phi(x) = [-1, 1]$ for every $x \in \mathbb{R}$; see, e.g., [41, p. 129]. Let $\Xi = \{-1, 1\}$ with equal probabilities, and define $f(\xi, x) = \frac{1}{2}x + \xi\phi(x)$. Then $\bar{\partial}Ef(x) = \{\frac{1}{2}\}$ and $\bar{\partial}_x f(\xi, x) = [-\frac{1}{2}, \frac{3}{2}]$ for every $\xi \in \Xi$ and $x \in \mathbb{R}$. Thus, any $x \in \mathbb{R}$ is weakly stationary for both the population and SAA problems, because $0 \in \mathbb{E}[\bar{\partial}_x f(\boldsymbol{\xi}, x)] = \frac{1}{\nu} \sum_{i=1}^\nu \bar{\partial}_x f(\boldsymbol{\xi}^i, x)$ for any $\nu \in \mathbb{N}$. But the population objective has no stationary point. Consequently, consistency of weak stationarity may hold under mild assumptions yet remain vacuous. By contrast, consistency of stationarity is a stronger concept but can fail under standard assumptions; see [53, Section 2.3].

The desired conclusion can be sharpened further by using finer notions of subdifferentiation. For a locally Lipschitz function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, the Clarke subdifferential satisfies $\bar{\partial}h(x) = \text{con } \partial h(x)$, where ∂h denotes the limiting subdifferential [42, Definition 8.3]. This convexification may produce Clarke stationary points that are not limiting stationary. For example, $\partial(-|\cdot|)(0) = \{-1, 1\}$, while $\bar{\partial}(-|\cdot|)(0) = [-1, 1]$. Thus, the strict local maximizer 0 is Clarke stationary but not limiting stationary. In general, limiting stationarity is a sharper solution concept than Clarke stationarity. They are equivalent when h is a subdifferentially regular function [42, Definition 7.25].

In this paper, we establish SAA consistency results for nonconvex nonsmooth problems that are sharper in two respects: We compare the subdifferentials of the population and SAA objectives without interchanging expectation and subdifferentiation, and we treat limiting subdifferentials rather than merely their Clarke convexifications. We achieve this by establishing a new uniform convergence result for limiting subdifferentials. Existing results concern enlarged Clarke subdifferentials [47], or restricting to subdifferentially regular functions for which the limiting and Clarke subdifferentials coincide; see [47, 20, 43]. To the best of our knowledge, no previous uniform law compares $\bar{\partial}E^\nu f$ and $\bar{\partial}Ef$ for an irregular class of locally Lipschitz random functions. Consequently, no comparable consistency theory has been available for limiting stationary points.

Our approach is to compare the population and SAA functions before passing to subdifferentials. We establish the following *Lipschitzian SLLN*:

$$d_X(E^\nu f, Ef) \rightarrow 0, \quad \mathbb{P}\text{-a.s.}, \quad (3)$$

where $X \subset \mathbb{R}^d$ and $d_X(\cdot, \cdot)$ denotes the Lipschitz pseudometric; see Section 2.1 for the definition. Informally, $d_X(\cdot, \cdot)$ controls both the function values and the global Lipschitz modulus of the function $E^\nu f - Ef$ on the set X . This makes it a natural mode of convergence for our purposes. Indeed,

if h and g are locally Lipschitz on an open set $X \subset \mathbb{R}^d$, then

$$\sup_{x \in X} d(\bar{\partial}h(x), \bar{\partial}g(x)) \leq \sup_{x \in X} d(\partial h(x), \partial g(x)) \leq d_X(h, g), \quad (4)$$

where $d(A, B)$ denotes the Hausdorff distance between nonempty sets A and B ; see Proposition 5.2. Thus, we obtain simultaneous uniform control of limiting and Clarke subdifferentials and, consequently, the desired consistency of stationary points. Section 5 discusses further applications and properties of the Lipschitzian SLLN. Notably, the work [53] shows that the uniform law corresponding to (4), with $h = E^\nu f$ and $g = Ef$, cannot hold in general even when every slice $f(\xi, \cdot)$ is convex and satisfies the standard integrability and Lipschitz assumptions. Hence, additional assumptions are necessary for the Lipschitzian law in (3) to hold.

We prove (3) under two distinct structural conditions, together with mild integrability and Lipschitz assumptions. The first is a topological separability condition; see Assumption 3.2. It holds, in particular, when either the distribution is discrete or the slices are continuously differentiable on a compact set; see Section 4.1. The second is a model-theoretic definability condition; see Assumption 3.7. It encompasses many nonsmooth functions arising in practice, including losses from modern learning models [5]; see Section 4.2. For illustration, under mild integrability and Lipschitz assumptions, the Lipschitzian law (3) holds if either of the following conditions is satisfied (cf. the more general results in Theorems 3.3, 3.8 and 4.10):

- (i) The family $\{f(\xi, \cdot)|_X \mid \xi \in \Xi\}$ is countable.
- (ii) For every $\xi \in \Xi$, the slice $f(\xi, \cdot)$ is definable in the real exponential field $\mathcal{R}_{\text{exp}}$.

The first condition arises naturally for discrete distributions, while the second is satisfied by the training losses of many modern deep learning models; see Section 2.3 for definitions and Section 4 for details. Despite their fundamentally different nature, both conditions ensure the Lipschitzian law and thereby rule out the failure phenomenon exhibited in [53].

In addition to establishing the Lipschitzian law (3), we study its convergence rate, obtaining the canonical rate $O_{\mathbb{P}}(\nu^{-1/2})$ in one setting and rates that can be arbitrarily slow in others. Beyond uniform convergence of subdifferentials, we also apply (3) to identify solutions to the population problem using only a finite number of samples, extending existing results [46, 32, 48] by relaxing convexity and distributional assumptions.

The remainder of the paper is organized as follows. Section 2 reviews Lipschitz spaces, VC classes, and the model-theoretic notions used throughout. Section 3 introduces the two structural assumptions and establishes the Lipschitzian SLLNs and their convergence rates. Section 4 provides practical conditions ensuring separability and definability. Section 5 discusses applications and related work, including uniform SLLNs for subdifferentials and finite-sample identification of solutions.

Notation. We use mostly standard notation. The sets of positive integers, integers, and rational numbers are denoted by $\mathbb{N} = \{1, 2, \dots\}$, $\mathbb{Z}$, and $\mathbb{Q}$, respectively. For $\nu \in \mathbb{N}$, write $[\nu] = \{1, \dots, \nu\}$, and let $\bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty, \infty\}$. For a set S , we denote its cardinality by $|S|$ and its power set by 2^S . For disjoint sets A and B , we write the union $A \cup B$ as $A \sqcup B$. Let $\|\cdot\|$ denote a norm on a Euclidean space, $\|\cdot\|_*$ its dual norm, $\|\cdot\|_2$ the Euclidean norm, and $\mathbb{B} = \{y \mid \|y\| \leq 1\}$. A function $g : \mathbb{R}^d \rightarrow \mathbb{R}$ is L -Lipschitz on $X \subset \mathbb{R}^d$ if $|g(x) - g(y)| \leq L\|x - y\|$ for all $x, y \in X$, and simply L -Lipschitz when $X = \mathbb{R}^d$. We denote the restriction of g to X by $g|_X$ and the graph of g

by $\text{gph } g$. For nonempty sets $C, D \subset \mathbb{R}^d$ and $x \in \mathbb{R}^d$, define $\text{dist}(x, C) = \inf_{z \in C} \|x - z\|$. The excess of C relative to D is $\text{exs}(C; D) = \sup_{z \in C} \text{dist}(z, D)$, and the Hausdorff distance between C and D is $d(C, D) = \max\{\text{exs}(C; D), \text{exs}(D; C)\}$. The convex hull and real linear span of a set S are denoted by $\text{con } S$ and $\text{span } S$, respectively. For a set C , define $\mathbf{1}_C(x) = 1, \iota_C(x) = 0$ if $x \in C$, and $\mathbf{1}_C(x) = 0, \iota_C(x) = \infty$ otherwise. For $t \in \mathbb{R}$, define $\text{sgn}(t) = t/|t|$ if $t \neq 0$ and $\text{sgn}(t) = 0$ otherwise. The Borel σ -algebra on a topological set S is denoted by $\mathcal{B}(S)$. A sequence of random variables $\{\mathbf{y}^\nu\}_{\nu \in \mathbb{N}} = O_{\mathbb{P}}(r^\nu)$ if $\{\mathbf{y}^\nu/r^\nu\}_{\nu \in \mathbb{N}}$ is bounded in probability.

2 Preliminaries

In this section, we set the stage and recall concepts and results used in the sequel. Let $(\Xi, \mathcal{A})$ be a measurable space, and let $\boldsymbol{\xi}, \boldsymbol{\xi}^1, \boldsymbol{\xi}^2, \dots$ be independent and identically distributed (iid) Ξ -valued random variables. For an $\mathcal{A}$ -measurable function $h : \Xi \rightarrow \mathbb{R}$, when the underlying random variable is clear from the context, we write its expectation $\mathbb{E}[h(\boldsymbol{\xi})]$ as $\mathbb{E}h$ and, by a slight abuse of notation, denote the sample average $\frac{1}{\nu} \sum_{i=1}^\nu h(\boldsymbol{\xi}^i)$ by $\mathbb{E}^\nu[h(\boldsymbol{\xi})]$ or $\mathbb{E}^\nu h$.

For a function $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ such that $f(\cdot, x)$ is $\mathcal{A}$ -measurable for every $x \in \mathbb{R}^d$, its expectation function $Ef : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$ and its sample-average approximation $E^\nu f : \mathbb{R}^d \rightarrow \mathbb{R}$ are given by

$$Ef(x) = \mathbb{E}[f(\boldsymbol{\xi}, x)], \quad E^\nu f(x) = \frac{1}{\nu} \sum_{i=1}^\nu f(\boldsymbol{\xi}^i, x),$$

adopting the convention $\infty - \infty = \infty$ if needed. As $\nu \rightarrow \infty$, we are interested in the convergence of $E^\nu f$ to Ef in suitable topologies and in the consequences of such convergence. Classical examples include epigraphical convergence [42, Section 7B] and uniform convergence [42, Section 7C]. Results of this kind are referred to in the literature as epigraphical SLLNs [3] and uniform SLLNs [49, Section 9.2.6].

In this paper, we focus on a stronger mode of convergence, defined through a Lipschitz space and Lipschitz metric recalled below.

2.1 Lipschitz space, norm, and pseudometric

For a real-valued function $h : Y \rightarrow \mathbb{R}$ and nonempty sets $X \subset Y \subset \mathbb{R}^d$, if X is not singleton, the *global Lipschitz modulus* $\text{g-lip}_X h \in \overline{\mathbb{R}}$ of the function h on the set X can be written as

$$\text{g-lip}_X h = \sup_{x, y \in X, x \neq y} \frac{h(x) - h(y)}{\|x - y\|}.$$

For singleton X , we define by convention $\text{g-lip}_X h = 0$. We may write $\text{g-lip}_X h$ as $\text{g-lip } h$ when $X = \text{dom } h$. The *local Lipschitz modulus* of h at x is defined by [42, Definition 9.1]

$$\text{lip } h(x) = \limsup_{y, z \rightarrow x, y \neq z} \frac{h(y) - h(z)}{\|y - z\|}.$$

The global Lipschitz modulus controls the local Lipschitz modulus at every interior point of X , as stated below. The proof is elementary and omitted.

Lemma 2.1. *If $X \subset \mathbb{R}^d$ and $h : \mathbb{R}^d \rightarrow \mathbb{R}$, then for every $x \in \text{int } X$, $\text{lip } h(x) \leq \text{g-lip}_X h$.*

When $0 \in X$, consider the *space of Lipschitz functions* $\text{Lip}(X)$ on X defined as

$$\text{Lip}(X) = \{h : X \rightarrow \mathbb{R} \mid \text{g-lip } h < \infty\},$$

which is endowed with the following Lipschitz norm (see [7] and [6, Section 4]):

$$\|h\|_{\text{lip}} = \max\{|h(0)|, \text{g-lip } h\}.$$

The space $(\text{Lip}(X), \|\cdot\|_{\text{lip}})$ is a Banach space [6, Proposition 4.1]. The point $0 \in X$ in the definition above can be replaced by any point in X without changing the topology [6, p. 27]. We make such a choice for notational simplicity. For $0 \in X \subset \mathbb{R}^d$ and functions $h, g : \mathbb{R}^d \rightarrow \mathbb{R}$ that are real-valued on X , the Lipschitz norm $\|\cdot\|_{\text{lip}}$ on $\text{Lip}(X)$ also gives rise to a Lipschitz (extended) pseudometric between h and g as follows

$$d_X(h, g) = \|(h - g)|_X\|_{\text{lip}} = \max\{|h(0) - g(0)|, \text{g-lip } (h|_X - g|_X)\}.$$

When $(h - g)|_X \in \text{Lip}(X)$, one has $d_X(h, g) < \infty$.

2.2 VC-class of sets and functions

Let $\mathcal{C}$ be a collection of subsets of a set S and $\{x_1, \dots, x_n\}$ be an arbitrary finite subset of S . We say the collection $\mathcal{C}$ *picks out* a subset A from S if there exists $C \in \mathcal{C}$ such that $A = C \cap \{x_1, \dots, x_n\}$. The collection $\mathcal{C}$ is said to *shatter* $\{x_1, \dots, x_n\}$ if each of its 2^n subsets can be picked out. The *Vapnik–Chervonenkis dimension* (VC-dimension) $\text{VC}(\mathcal{C})$ of the class $\mathcal{C}$ is the largest $n \in \mathbb{N}$ such that some $\{x_1, \dots, x_n\} \subset S$ can be shattered by $\mathcal{C}$; see [54, Section 2.6.1]. Formally, one has

$$\text{VC}(\mathcal{C}) = \sup \left\{ n \in \mathbb{N} \mid 2^n = \max_{x_1, \dots, x_n \in S} |\{C \cap \{x_1, \dots, x_n\} \mid C \in \mathcal{C}\}| \right\},$$

which can be infinity if $\mathcal{C}$ shatters sets of arbitrarily large size and we write $\text{VC}(\mathcal{C}) = \infty$. If $\text{VC}(\mathcal{C}) < \infty$, we call the collection $\mathcal{C}$ a VC-class.

VC-classes are stable under certain operations; see [54, Section 2.6.6]. The following properties will be useful for us.

Lemma 2.2 (permanence). *Given $G : X \rightarrow Y$ and VC-classes $\mathcal{C}, \mathcal{D} \subset 2^Y$, the following hold.*

- (a) *The family $\{C \cup D \mid C \in \mathcal{C}, D \in \mathcal{D}\}$ is a VC-class; see [54, Lemma 2.6.19(iii)].*
- (b) *The family $\mathcal{C} \cup \mathcal{D}$ is a VC-class; see [45, Exercise 6.11].*
- (c) *The family $G^{-1}(\mathcal{C}) = \{G^{-1}(C) \mid C \in \mathcal{C}\}$ is a VC-class; see [54, Lemma 2.6.19(v)].*

We define similar concepts for functions by considering their subgraphs (or, hypographs); see [54, Section 2.6.2]. For a collection of functions $\mathcal{H} = \{h_i : \Xi \rightarrow \mathbb{R} \mid i \in I\}$, we call $\mathcal{H}$ a VC-subgraph class if the collection $\{(\xi, r) \in \Xi \times \mathbb{R} \mid r < h_i(\xi)\} \mid i \in I\}$ is a VC-class.

2.3 Definability, NIP, and o-minimality

For general introductions to model theory, see [14]. A first-order language $\mathcal{L}$ consists of relation, function, and constant symbols. An $\mathcal{L}$ -structure $\mathcal{M}$ consists of a nonempty universe M together with interpretations of all symbols of $\mathcal{L}$ on M . An $\mathcal{L}$ -formula is built from atomic formulas using logical connectives and quantifiers. A set $S \subset M^d$ is said to be *definable* in the structure $\mathcal{M}$ if there exist an $\mathcal{L}$ -formula $\varphi(x; y)$, with variables $x = (x_1, \dots, x_d)$ and $y = (y_1, \dots, y_m)$, and parameter $b \in M^m$ such that

$$S = \varphi(M^d; b) = \{a \in M^d \mid \mathcal{M} \models \varphi(a; b)\}.$$

Here, $\mathcal{M} \models \varphi(a; b)$ means that the formula $\varphi(a; b)$ holds in $\mathcal{M}$. We may simply write $\varphi(a; b)$ when $\mathcal{M}$ is clear from context. The semicolon in $\varphi(x; y)$ has no syntactic significance; it is used only to distinguish the object variables x from the parameter variables y . To emphasize the dependence, we may say that the set S is definable in the structure $\mathcal{M}$ by the formula ϕ with parameter b . A function is called definable if its graph is definable. Let $\mathcal{M}$ and $\mathcal{M}'$ have the same universe. If the language of $\mathcal{M}'$ extends that of $\mathcal{M}$ and the interpretations of the common symbols agree, then $\mathcal{M}'$ is an *expansion* of $\mathcal{M}$, and $\mathcal{M}$ is a *reduct* of $\mathcal{M}'$. In this paper, we will only consider structures that are expansions of the real ordered field $\mathcal{R} = (\mathbb{R}, +, \cdot, <, 0, 1)$.

For illustration, we examine $\mathcal{R}$ more closely. The structure $\mathcal{R}$ consists of the universe $\mathbb{R}$, the functions $\{+, \cdot\}$, the relation $\{<\}$, and the constants $\{0, 1\}$. Hence, the language of $\mathcal{R}$ consists of the symbols $(+, \cdot, <, 0, 1)$. A formula in that language is built inductively from atomic formulas using the logical connectives $\neg, \wedge, \vee$, together with the quantifiers $\exists, \forall$. For example, define a formula $\phi(x; y) = (x_1 > y) \wedge (x_2 > y)$, where $x = (x_1, x_2) \in \mathbb{R}^2$ is the object variable and $y \in \mathbb{R}$ is the parameter variable. For every $b \in \mathbb{R}$, one has $\phi(\mathbb{R}^2; b) = (b, \infty)^2$. Thus, a single formula ϕ defines a family of sets as the parameter b varies; in particular, the choice $b = 0$ defines $(0, \infty)^2$. Notably, by the Tarski–Seidenberg theorem [19, Theorem 2.6], the sets definable in the structure $\mathcal{R}$ are precisely the semialgebraic sets, namely, finite Boolean combinations of sets defined by polynomial equalities and inequalities with real coefficients; see [19, Section 2.1.1].

Functions definable in an arbitrary expansion of $\mathcal{R}$ can be quite wild. We will focus on expansions that have the *non-independence property* (NIP); see [50, Definition 2.1].

Definition 2.3 ([34, Definitions 1.1 and 1.2]). *A formula $\phi(x_1, \dots, x_m; y_1, \dots, y_k)$ has the independence property (IP) with respect to a structure $\mathcal{M}$ with universe M if for every $n \in \mathbb{N}$ there is a sequence $(a_1, \dots, a_n)$ of elements from M^k so that for every $I \subset [n]$ there is $c_I \in M^m$ with*

$$\mathcal{M} \models \phi(c_I; a_i) \iff i \in I.$$

A formula has IP if it has the independence property, and has NIP otherwise. A structure $\mathcal{M}$ has NIP if every formula in its language has NIP with respect to $\mathcal{M}$; otherwise, it has IP.

Our focus is on functions and sets definable in NIP expansions of $\mathcal{R}$. A key consequence of such definability is the following observation.

Lemma 2.4 ([34, Proposition 1.3]). *For a structure $\mathcal{M}$ with universe M and a formula $\phi(x; y)$ with $x \in M^m$ and $y \in M^k$ in the language of $\mathcal{M}$, the family*

$$\{\{c \in M^m \mid \mathcal{M} \models \phi(c; b)\} \mid b \in M^k\}$$

is a VC-class if and only if the formula ϕ has NIP.

VC-classes play a central role in statistical learning theory [45, Chapter 6]. Motivated by this connection, definability in NIP structures has been studied extensively in learning theory [37, 31, 36, 15] and, more recently, in stochastic programming [39]. These works focus primarily on uniform convergence of function values, which is insufficient to control first-order objects such as subdifferentials.

Another important concept in model theory is that of an *o-minimal* structure; see [56] and [18, Definition 1.4]. An expansion of $\mathcal{R}$ is o-minimal if every definable subset of $\mathbb{R}$ is a finite union of points and intervals. Such structures have been widely used in optimization to exclude pathological behavior; see [28] and the references therein. The real ordered field $\mathcal{R}$ introduced above and the real exponential field $\mathcal{R}_{\text{exp}} = (\mathcal{R}, \exp)$ are both o-minimal structures; see [18, Exercise 1.7] and [59].

Proposition 2.5 ([50, Example 2.12]). *Every o-minimal structure has NIP.*

Conversely, definability of the integers $\mathbb{Z}$ provides a simple sufficient condition for IP and can be useful in constructing counterexamples.

Proposition 2.6 ([33, Observation 2.6 and footnote 4]). *If the set $\mathbb{Z}$ is definable in an expansion $\mathcal{M}$ of $\mathcal{R}$, then $\mathcal{M}$ has IP.*

3 SLLNs in Lipschitz Pseudometric

In this section, we present our general results for the convergence of $E^\nu f$ to Ef in terms of the Lipschitz pseudometric. We begin with a blanket assumption.

Assumption 3.1 (blanket). *Given a probability space $(\Xi, \mathcal{A}, P)$, for a function $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$, a random variable ξ with law P , and a set $X \subset \mathbb{R}^d$ with $0 \in X$, assume the following hold.*

- (i) *For any $x \in \mathbb{R}^d$, the function $f(\cdot, x)$ is $\mathcal{A}$ -measurable.*
- (ii) *$\mathbb{E}[|f(\xi, 0)|] < \infty$.*
- (iii) *There exists an $\mathcal{A}$ -measurable $L : \Xi \rightarrow [0, \infty)$ with $\mathbb{E}L < \infty$ such that, for any $\xi \in \Xi$, the function $f(\xi, \cdot)$ is $L(\xi)$ -Lipschitz on X .*

Assumption 3.1 imposes mild conditions ensuring well-definedness; see also [47, 43] for similar assumptions. The condition $0 \in X$ is without loss of generality. Indeed, for any $x_0 \in X$, one may replace X by $X - \{x_0\}$ and f by the translated function $(\xi, x) \mapsto f(\xi, x + x_0)$. Assumption 3.1 requires no compactness, openness, or measurability assumption on X .

3.1 Separability

Our first result applies to functions satisfying the following topological condition.

Assumption 3.2 (separability). *Given $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $X \subset \mathbb{R}^d$ with $0 \in X$, assume the collection $\{f(\xi, \cdot)|_X \mid \xi \in \Xi\}$ lies in a separable subspace of $(\text{Lip}(X), \|\cdot\|_{\text{lip}})$.*

Recall that a topological space is separable if it contains a countable dense subset. Under Assumption 3.1, a straightforward sufficient condition for Assumption 3.2 is Ξ being countable, which is natural when the random variable ξ has a discrete distribution. Separability, possibly in

different forms, has already been used to establish SLLNs for random variables taking values in infinite-dimensional spaces; see, e.g., [35, 22, 52]. We defer the discussion on functions satisfying Assumption 3.2 to Section 4.1.

Theorem 3.3 (separability). *Let $(\Xi, \mathcal{A})$ be a measurable space and $\xi, \xi^1, \xi^2, \dots$ be Ξ -valued iid random variables on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Suppose that $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $X \subset \mathbb{R}^d$ satisfy Assumptions 3.1 and 3.2. Then*

$$\lim_{\nu \rightarrow \infty} d_X(E^\nu f, Ef) = 0, \quad \mathbb{P}\text{-a.s.}$$

In essence, Theorem 3.3 follows from a strong law for separable Banach-valued random variables; see, e.g., [35, Chapter 7]. The main task is to verify its applicability. To this end, Assumption 3.2 allows us to regard the mapping $\xi \mapsto f(\xi, \cdot)|_X$ as a random variable taking values in a separable subspace of $(\text{Lip}(X), \|\cdot\|_{\text{lip}})$. The proof then concludes by identifying its Bochner integral with the corresponding pointwise integral. We defer the proof of Theorem 3.3 to Section 3.3.1.

As mentioned above, we refer to such convergence of $E^\nu f$ to Ef in the Lipschitz pseudometric as a Lipschitzian SLLN. Indeed, Theorem 3.3 generalizes Kolmogorov's strong law of large numbers.

Example 3.4 (Kolmogorov's SLLN). *Suppose that $(\Xi, \mathcal{A}) = (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ and $\xi, \xi^1, \xi^2, \dots$ be Ξ -valued iid random variables on the complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with $\mathbb{E}[|\xi|] < \infty$. Let $d = 1, X = \mathbb{R}$, $f(\xi, x) = \xi$ for all $x \in \mathbb{R}$. Hence, Assumptions 3.1 and 3.2 hold trivially by the separability of $\mathbb{R}$, and Theorem 3.3 reduces to Kolmogorov's SLLN.*

In contrast to Assumption 3.1, one may not argue that Assumption 3.2 is a mild requirement. Indeed, as shown in the following example, adapted from [40, Example 3.4], separability can fail for a simple, convex $f : \mathbb{R}^2 \rightarrow \mathbb{R}$.

Example 3.5 (nonseparability). *Let $d = 1, X = \mathbb{R}, \Xi = [0, 1], \mathcal{A} = \mathcal{B}(\Xi)$, and $f(\xi, x) = |x - \xi|$. Then Assumption 3.1 holds. Moreover, for any distinct $\xi, \xi' \in \Xi$, one has $\|f(\xi, \cdot) - f(\xi', \cdot)\|_{\text{lip}} = 2$. Hence, the family $\{f(\xi, \cdot)|_X \mid \xi \in \Xi\}$ is not contained in any separable subspace of $\text{Lip}(X)$.*

Therefore, without Assumption 3.2, the applicability of the Banach-valued strong law is not automatic. One may wonder whether the lack of separability reflects a genuine obstruction. The following result, as an easy corollary of [53, Theorem 3], shows that the conclusion of Theorem 3.3 might fail without the separability assumption.

Proposition 3.6 (failure). *Let $X = [-2, 2]^2 \subset \mathbb{R}^2$ and $\xi, \xi^1, \xi^2, \dots$ be iid random variables on the complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$, each uniformly distributed on $\Xi = [0, 1]$. There exist a function $f : \Xi \times \mathbb{R}^2 \rightarrow \mathbb{R}$ and constant $c > 0$ such that the following hold.*

- (a) *Assumption 3.1 holds, and $f(\xi, \cdot)$ is convex for every $\xi \in \Xi$.*
- (b) *$\mathbb{P}$ -a.s., one has $\liminf_{\nu \rightarrow \infty} d_X(E^\nu f, Ef) \geq c$.*

Proof. Let f be the bivariate convex construction in [53, Theorem 3], which satisfies Assumption 3.1. Let $c > 0$ be such that $\|\cdot\|_* \geq 2c\|\cdot\|_2$, where $\|\cdot\|_*$ is the dual norm of $\|\cdot\|$. From [53, Section 3.4, Step 5], there exist random points $\{\mathbf{p}^\nu\}_\nu \in \{(0, 0)\} \cup \{(0, k^{-1})\}_{k \in \mathbb{N}} \subset X$ such that, $\mathbb{P}$ -a.s., $E^\nu f$ and Ef are both continuously differentiable at $\mathbf{p}^\nu$ and

$$\liminf_{\nu \rightarrow \infty} d_X(E^\nu f, Ef) \geq \liminf_{\nu \rightarrow \infty} 2c\|\nabla E^\nu f(\mathbf{p}^\nu) - \nabla Ef(\mathbf{p}^\nu)\|_2 \geq c,$$

where the last inequality is from [53, Theorem 3]. □

3.2 Definability

As shown in Proposition 3.6, Assumption 3.1 alone does not guarantee convergence in the Lipschitz pseudometric. Conversely, Assumption 3.2 can be restrictive and, as illustrated by Example 3.5, excludes even simple convex semialgebraic functions. In this subsection, we introduce an alternative structural assumption that avoids both the Banach-valued strong law and the associated separability requirement.

Assumption 3.7 (piecewise uniform definability). *Let $\{q_k\}_{k \in \mathbb{N}}$ be a dense subset of $X \subset \mathbb{R}^d$. Given a measurable space $(\Xi, \mathcal{A})$ and a function $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$, assume that, for some index set $I \subset \mathbb{N}$ and every $\ell \in I$, there exist*

- (i) *a universally measurable set $\Xi_\ell \subset \Xi$ relative to $\mathcal{A}$,*
- (ii) *an NIP expansion $\mathcal{M}_\ell$ of $\mathcal{R}$, and*
- (iii) *a formula $\phi_\ell(x, r; z)$ with $x \in \mathbb{R}^d$, $r \in \mathbb{R}$, and $z \in \mathbb{R}^{k_\ell}$ in the language of $\mathcal{M}_\ell$*

such that $\cup_{\ell \in I} \Xi_\ell = \Xi$ and, for any $\xi \in \Xi_\ell$, there exists parameter $z \in \mathbb{R}^{k_\ell}$ such that, for each $k \in \mathbb{N}$,

$$r = f(\xi, q_k) \iff \mathcal{M}_\ell \models \phi_\ell(q_k, r; z). \quad (5)$$

For $\ell \in I$ and $\xi \in \Xi_\ell$, if $f(\xi, \cdot)$ is definable by ϕ_ℓ , then (5) follows immediately by examining only $\{f(\xi, q_k)\}_{k \in \mathbb{N}}$. Recall that a set $S \subset \Xi$ is universally measurable relative to $\mathcal{A}$ if it is measurable with respect to every completion of $\mathcal{A}$. The set X need not be definable or measurable. Although Assumption 3.7 may initially appear restrictive, it naturally generalizes joint definability in a fixed o-minimal structure; see Section 4.2.1. Here, countable covers $\{\Xi_\ell\}_{\ell \in I}$ of Ξ are standard tools for establishing uniform convergence in empirical process theory [57, 58] and learning theory [45, Chapter 7]. In Section 4.2.3, we show that, when the languages of the structures $\{\mathcal{M}_\ell\}_{\ell \in I}$ are countable, uniform definability in Assumption 3.7 can be replaced by pointwise definability; see Assumption 4.9.

Assumption 3.7 encompasses a broad class of functions arising in practice and we defer a detailed discussion to Section 4.2. The construction in Example 3.5 satisfies Assumption 3.7 because f is jointly semialgebraic. Thus, Assumption 3.7 may hold even when the separability condition fails.

Under the definability condition in Assumption 3.7, we obtain the following result.

Theorem 3.8 (piecewise uniform definability). *Let $(\Xi, \mathcal{A})$ be a measurable space and $\xi, \xi^1, \xi^2, \dots$ be Ξ -valued iid random variables on the complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Suppose that $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $X \subset \mathbb{R}^d$ satisfy Assumptions 3.1 and 3.7. Then*

$$\lim_{\nu \rightarrow \infty} d_X(E^\nu f, Ef) = 0, \quad \mathbb{P}\text{-a.s.}$$

Unlike in Theorem 3.3, the mapping $\xi \mapsto f(\xi, \cdot)|_X$ need not take values in a separable Banach space. Instead, we reduce convergence in the Lipschitz pseudometric to the Glivenko–Cantelli property of a class of difference quotients indexed by $\{q_k\}_{k \in \mathbb{N}}$. The cover $\{\Xi_\ell\}_{\ell \in I}$ in Assumption 3.7 induces a countable partition of Ξ . On each piece, the restrictions of the slices $\{f(\xi, \cdot) \mid \xi \in \Xi_\ell\}$ to $\{q_k\}_{k \in \mathbb{N}}$ are uniformly definable in an NIP structure; consequently, the corresponding collection of difference quotients forms a VC-subgraph class. The desired convergence then follows by combining the corresponding estimates across the partition. Notably, the norm $\|\cdot\|$ used to define the

pseudometric d_X need not be definable; in particular, our analysis is not restricted to the Euclidean norm. We defer the proof of Theorem 3.8 to Section 3.3.2.

Under stronger integrability assumptions and the additional assumption that $|I| < \infty$, we obtain the following convergence rate.

Theorem 3.9 (rate of convergence). *Suppose that $(\Xi, \mathcal{A})$, $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$, and $X \subset \mathbb{R}^d$ satisfy Assumptions 3.1 and 3.7 with $|I| < \infty$. Let $\xi, \xi^1, \xi^2, \dots$ be Ξ -valued iid random variables on the complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. If, in addition, $\mathbb{E}[|f(\xi, 0)|^2] < \infty$ and $\mathbb{E}[L^2] < \infty$, then*

$$d_X(E^\nu f, Ef) = O_{\mathbb{P}}(\nu^{-1/2}).$$

In other words, for any $\delta > 0$, there exist a constant $C_\delta < \infty$ and $\nu_0 \in \mathbb{N}$ such that for all $\nu \geq \nu_0$, with probability at least $1 - \delta$, one has $d_X(E^\nu f, Ef) \leq C_\delta \nu^{-1/2}$.

Theorem 3.9 may be viewed as a uniform tightness result for the sequence $\{d_X(E^\nu f, Ef)\}_\nu$. Its proof is deferred to Section 3.3.3. Under the stated assumptions, we do not have uniform control of the constants C_δ because of the qualitative nature of Lemma 2.4. Nevertheless, C_δ are related to the dimension d and the VC-dimensions of certain function classes. Explicit bounds may be obtainable in more structured settings; we do not pursue this direction here. We emphasize that the finite-pieces requirement in Theorem 3.9 cannot be removed without loss of the rate of convergence.

Proposition 3.10. *Let $\Xi = \mathbb{N}$, $\mathcal{A} = 2^{\mathbb{N}}$, $d = 1$, and $X = [0, 1]$. There exists a function $f : \Xi \times \mathbb{R} \rightarrow \mathbb{R}$ satisfying Assumptions 3.1, 3.2, and 3.7, such that $L(\xi) = 1$ and $f(\xi, \cdot)$ is semialgebraic for every $\xi \in \Xi$, with the following property. For every sequence $r^\nu \in (0, 1) \downarrow 0$, there exists a law P on Ξ such that, for any iid sequence $\xi, \xi^1, \xi^2, \dots$ with law P on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$,*

$$d_X(E^\nu f, Ef) \neq O_{\mathbb{P}}(r^\nu).$$

The proof of Proposition 3.10 constructs a semialgebraic basis that relates the Lipschitz pseudometric to the *missing mass problem* studied in [8], whose convergence can be arbitrarily slow; see Section 3.3.4 for details. Since $L(\xi) \equiv 1$, the square-integrability condition in Theorem 3.9 holds trivially. Nevertheless, the convergence in Theorems 3.3 and 3.8 can be arbitrarily slow. By contrast, for sample averages of $\mathbb{R}^m$ -valued random variables, finite second moments yield a central limit theorem and hence the canonical rate $O_{\mathbb{P}}(\nu^{-1/2})$. This contrast highlights the fundamental difference between averaging random vectors in a Euclidean space and averaging random functions in the Lipschitz space.

3.3 Proofs

We begin with some preparatory moves. Let $(\Xi, \mathcal{A})$ be a measurable space. Suppose that the function $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ and the set $X \subset \mathbb{R}^d$ satisfy Assumption 3.1. Let $D = \{q_k\}_{k \in \mathbb{N}}$ be a countable dense subset of X . If X is a singleton, then necessarily $X = \{0\}$ and $d_X(E^\nu f, Ef) = |E^\nu f(0) - Ef(0)|$ and the results in Section 3 reduce directly to the classical SLLN. Hence, in what follows, we assume that X contains at least two distinct points. For every $x \in X$, one has $|Ef(x)| \leq \mathbb{E}[|f(\xi, x)|] \leq \mathbb{E}[|f(\xi, 0)|] + (\mathbb{E}L)\|x\|$. Since $\mathbb{E}[|f(\xi, 0)|] < \infty$ and $\mathbb{E}L < \infty$, the function $Ef : \mathbb{R}^d \rightarrow \mathbb{R}$ is real-valued on X . For any $x, y \in X, x \neq y$, let a function $g_{x,y} : \Xi \rightarrow \mathbb{R}$ be

$$g_{x,y}(\xi) = \frac{f(\xi, x) - f(\xi, y)}{\|x - y\|}.$$

For any $\mathcal{A}$ -measurable set $S \subset \Xi$, define $\mathcal{G}_S = \{g_{x,y}\mathbf{1}_S \mid x, y \in D, x \neq y\}$. Observe the identity

$$\text{g-lip}_X(E^\nu f - Ef) = \sup_{x,y \in X, x \neq y} \frac{|(E^\nu f(x) - Ef(x)) - (E^\nu f(y) - Ef(y))|}{\|x - y\|} = \sup_{g \in \mathcal{G}_\Xi} |\mathbb{E}^\nu g - \mathbb{E}g|.$$

To see the last equality, fix $x, y \in X, x \neq y$. Let $x_k, y_k \in D$ be such that $(x_k, y_k) \rightarrow (x, y)$. Since $x \neq y$, for all sufficiently large $k \in \mathbb{N}$, one has $x_k \neq y_k$. Then, for any $\xi \in \Xi$, one has $f(\xi, x_k) \rightarrow f(\xi, x)$, $f(\xi, y_k) \rightarrow f(\xi, y)$, and $\|x_k - y_k\| \rightarrow \|x - y\| > 0$, so that we get $g_{x_k, y_k} \rightarrow g_{x, y}$ pointwise and the claim follows by the dominated convergence theorem. Thus, the convergence of $\text{g-lip}_X(E^\nu f - Ef)$ is governed by the empirical process indexed by $\mathcal{G}_\Xi$. Let $P = \mathbb{P} \circ \boldsymbol{\xi}^{-1}$ be the law of $\boldsymbol{\xi}$. Throughout the proofs, we work on appropriate probability-one events on which all relevant almost sure limits hold. Such events exist because only countably many almost sure statements are used and $\mathcal{F}$ is $\mathbb{P}$ -complete; for notational simplicity, we do not name them explicitly.

3.3.1 Proof of Theorem 3.3

Let B be the $\|\cdot\|_{\text{lip}}$ -closure of the separable subspace in Assumption 3.2, hence $(B, \|\cdot\|_{\text{lip}})$ is a separable Banach space. Let $F : \Xi \rightarrow B$ be the mapping $\xi \mapsto f(\xi, \cdot)|_X \in B$. We will view $F(\boldsymbol{\xi}) : \Omega \rightarrow B$ as a B -valued random variable. To this end, it suffices to show that F is $\mathcal{A}/\mathcal{B}(B)$ -measurable. By restricting the defining supremum to the countable set D , one sees that the function $\xi \mapsto \text{g-lip}(f(\xi, \cdot)|_X - b)$ is $\mathcal{A}$ -measurable for any fixed and deterministic $b \in B \subset \text{Lip}(X)$. Since $\xi \mapsto |f(\xi, 0) - b(0)|$ is $\mathcal{A}$ -measurable, the function $\xi \mapsto \|f(\xi, \cdot)|_X - b\|_{\text{lip}}$ is $\mathcal{A}$ -measurable. Let $\{b_k\}_{k \in \mathbb{N}}$ be a countable dense subset of B . For each $k \in \mathbb{N}$ and any $r \in \mathbb{Q}$, we have

$$F^{-1}(\{b \in B \mid \|b - b_k\|_{\text{lip}} < r\}) = \{\xi \in \Xi \mid \|f(\xi, \cdot)|_X - b_k\|_{\text{lip}} < r\} \in \mathcal{A},$$

so that F is $\mathcal{A}/\mathcal{B}(B)$ -measurable. Since B is separable, by [1, Lemma 11.37], $F(\boldsymbol{\xi}), F(\boldsymbol{\xi}^1), F(\boldsymbol{\xi}^2), \dots$ are strongly measurable B -valued iid random variables on $(\Omega, \mathcal{F}, \mathbb{P})$. For any $\xi \in \Xi$, we have

$$\|F(\xi)\|_{\text{lip}} = \max\{|f(\xi, 0)|, \text{g-lip}(f(\xi, \cdot)|_X)\} \leq |f(\xi, 0)| + L(\xi).$$

Therefore, $\mathbb{E}[\|F(\boldsymbol{\xi})\|_{\text{lip}}] \leq \mathbb{E}[|f(\boldsymbol{\xi}, 0)|] + \mathbb{E}[L(\boldsymbol{\xi})] < \infty$, so that $F(\boldsymbol{\xi})$ is Bochner integrable from [1, Theorem 11.44]; see also [35, p. 42]. Using [35, Corollary 7.10], one has

$$\left\| \frac{1}{\nu} \sum_{i=1}^{\nu} F(\boldsymbol{\xi}^i) - \mathbb{E}[F(\boldsymbol{\xi})] \right\|_{\text{lip}} \rightarrow 0, \quad \mathbb{P}\text{-a.s.},$$

where $\mathbb{E}[F(\boldsymbol{\xi})] \in B$ denotes the Bochner integral of $F(\boldsymbol{\xi})$. Now, we relate the above convergence to that of $E^\nu f$ and Ef . For any $x \in X$ and $\nu \in \mathbb{N}$, we have

$$\left(\frac{1}{\nu} \sum_{i=1}^{\nu} F(\boldsymbol{\xi}^i) \right)(x) = \frac{1}{\nu} \sum_{i=1}^{\nu} f(\boldsymbol{\xi}^i, x) = E^\nu f(x).$$

For each $x \in X$, define the evaluation map $\text{ev}_x : B \rightarrow \mathbb{R}$ by $\text{ev}_x(b) = b(x)$ for $b \in B$. This map is linear. Meanwhile, for any $b \in B$, one has

$$|\text{ev}_x(b)| \leq |b(0)| + \text{g-lip}(b)\|x\| \leq (1 + \|x\|)\|b\|_{\text{lip}},$$

hence the linear map ev_x is bounded. From [1, Lemma 11.45], for any $x \in X$,

$$\mathbb{E}[F(\boldsymbol{\xi})](x) = \text{ev}_x(\mathbb{E}[F(\boldsymbol{\xi})]) = \mathbb{E}[\text{ev}_x(F(\boldsymbol{\xi}))] = \mathbb{E}[f(\boldsymbol{\xi}, x)] = Ef(x).$$

Therefore, as $\nu \rightarrow \infty$, we obtain

$$d_X(E^\nu f, Ef) = \|(E^\nu f - Ef)|_X\|_{\text{lip}} = \left\| \frac{1}{\nu} \sum_{i=1}^{\nu} F(\boldsymbol{\xi}^i) - \mathbb{E}[F(\boldsymbol{\xi})] \right\|_{\text{lip}} \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

3.3.2 Proof of Theorem 3.8

Let $\{\Xi_\ell\}_{\ell \in I}$ be the cover of Ξ in Assumption 3.7. After reindexing and setting $\Xi_\ell = \emptyset$ for unused indices, we may assume without loss of generality that $I = \mathbb{N}$. Due to universal measurability, each Ξ_ℓ is measurable on the P -completion $\mathcal{A}^P$ of $\mathcal{A}$. For any $\ell \in \mathbb{N}$, there exist $C_\ell \subset \Xi_\ell \subset C'_\ell$ such that $C_\ell, C'_\ell \in \mathcal{A}$ and $P(C'_\ell \setminus C_\ell) = 0$; see [17, Section 1.5]. Let $B_\ell = C_\ell \setminus (\cup_{k < \ell} C_k)$ and $A_\ell = \cup_{k \in [\ell]} C_k = \sqcup_{k \in [\ell]} B_k$, so that $\{A_\ell\}_\ell$ and $\{B_\ell\}_\ell$ are both $\mathcal{A}$ -measurable. One has $P(\cup_\ell C_\ell) = 1$, since

$$1 = P(\Xi) = P(\cup_\ell \Xi_\ell) \leq P(\cup_\ell C'_\ell) \leq P(\cup_\ell C_\ell) + P(\cup_\ell (C'_\ell \setminus C_\ell)) = P(\cup_\ell C_\ell).$$

Therefore, $P(A_\ell) \uparrow 1$. A key ingredient of our proof is the following VC-subgraph lemma.

Lemma 3.11. *For each $\ell \in \mathbb{N}$, the family $\mathcal{G}_{B_\ell}$ is a VC-subgraph class.*

Proof. For any $x, y \in \mathbb{R}^d$, let

$$S_{x,y}^\ell = \{(\xi, t) \in B_\ell \times \mathbb{R} \mid t\|x - y\| < f(\xi, x) - f(\xi, y)\}.$$

Define the following formula in the language of $\mathcal{M}_\ell$:

$$\eta_\ell(z, t; u, v, a) = (\exists r_1, r_2) \quad \phi_\ell(u, r_1; z) \wedge \phi_\ell(v, r_2; z) \wedge ta < r_1 - r_2$$

with $r_1, r_2 \in \mathbb{R}, z \in \mathbb{R}^{k_\ell}, t \in \mathbb{R}, u \in \mathbb{R}^d, v \in \mathbb{R}^d, a \in \mathbb{R}$. Since $\mathcal{M}_\ell$ has NIP, every formula in its language has NIP, so that η_ℓ has NIP. Using Lemma 2.4, we know that the family

$$\left\{ \{(z, t) \in \mathbb{R}^{k_\ell+1} \mid \mathcal{M}_\ell \models \eta_\ell(z, t; x, y, a)\} \mid x, y \in \mathbb{R}^d, a \in \mathbb{R} \right\}$$

is a VC-class. Hence, its subfamily

$$\left\{ Z_{x,y}^\ell = \{(z, t) \in \mathbb{R}^{k_\ell+1} \mid \mathcal{M}_\ell \models \eta_\ell(z, t; x, y, \|x - y\|)\} \mid x, y \in D, x \neq y \right\}$$

is also a VC-class. For each $\ell \in \mathbb{N}$, consider the following mapping

$$\Psi_\ell : B_\ell \times \mathbb{R} \longrightarrow \mathbb{R}^{k_\ell} \times \mathbb{R}, \quad (\xi, t) \longmapsto (z_{\ell, \xi}, t).$$

For any $x, y \in D, x \neq y$, by Assumption 3.7 and the inclusions $B_\ell \subset C_\ell \subset \Xi_\ell$, we observe

$$\begin{aligned} S_{x,y}^\ell &= \{(\xi, t) \in B_\ell \times \mathbb{R} \mid r_1 = f(\xi, x), r_2 = f(\xi, y), a = \|x - y\|, ta < r_1 - r_2\} \\ &= \{(\xi, t) \in B_\ell \times \mathbb{R} \mid \mathcal{M}_\ell \models \phi_\ell(x, r_1; z_{\ell, \xi}) \wedge \phi_\ell(y, r_2; z_{\ell, \xi}), a = \|x - y\|, ta < r_1 - r_2\} \\ &= \{(\xi, t) \in B_\ell \times \mathbb{R} \mid \mathcal{M}_\ell \models \eta_\ell(z_{\ell, \xi}, t; x, y, \|x - y\|)\} = \Psi_\ell^{-1}(Z_{x,y}^\ell). \end{aligned}$$

By Lemma 2.2, the family $\{S_{x,y}^\ell \mid x, y \in D, x \neq y\}$ is a VC-class. Note that

$$\left\{ \{(\xi, t) \mid t < g_{x,y}(\xi) \mathbf{1}_{B_\ell}(\xi)\} \mid x, y \in D, x \neq y \right\} = \left\{ S_{x,y}^\ell \cup (B_\ell^c \times (-\infty, 0)) \mid x, y \in D, x \neq y \right\}.$$

Since a class containing only one set is trivially a VC-class, by Lemma 2.2, the above collection is a VC-class. Therefore, the collection $\mathcal{G}_{B_\ell}$ is a VC-subgraph class. $\square$

We are now ready to prove Theorem 3.8, following an argument similar to that of [58, Theorem 4]. For any distinct $x, y \in D$, the function $g_{x,y}$ is $\mathcal{A}$ -measurable, since $f(\cdot, z)$ is $\mathcal{A}$ -measurable for every $z \in \mathbb{R}^d$. For any $\mathcal{A}$ -measurable S , the function $g_{x,y}\mathbf{1}_S$ is $\mathcal{A}$ -measurable as well. Thus, by countability of D , the families $\mathcal{G}_{A_\ell}$ and $\mathcal{G}_{B_\ell}$ are both pointwise measurable [54, Example 2.3.4], hence P -measurable in the sense of [54, Definition 2.3.3]. Since $f(\xi, \cdot)$ is $L(\xi)$ -Lipschitz on X , one has $|g_{x,y}(\xi)\mathbf{1}_S(\xi)| \leq |g_{x,y}(\xi)| \leq L(\xi)$ for any $x, y \in X, x \neq y$. By $\mathbb{E}L < \infty$, $\mathcal{G}_{B_\ell}$ has an integrable envelope L . From Lemma 3.11, the family $\mathcal{G}_{B_\ell}$ is a VC-subgraph class. By the control of uniform entropy from [54, Theorem 2.6.7] and then invoking [54, Theorem 2.8.1], the class $\mathcal{G}_{B_\ell}$ is P -Glivenko–Cantelli. Hence, for each $k \in \mathbb{N}$,

$$\lim_{\nu \rightarrow \infty} \sup_{g \in \mathcal{G}_{B_k}} |\mathbb{E}^\nu g - \mathbb{E}g| = 0, \quad \mathbb{P}\text{-a.s.}$$

Since $\mathbf{1}_{A_\ell} = \sum_{k=1}^\ell \mathbf{1}_{B_k}$, for any $\ell \in \mathbb{N}$, $\mathbb{P}$ -a.s., one has

$$\limsup_{\nu \rightarrow \infty} \sup_{g \in \mathcal{G}_{A_\ell}} |\mathbb{E}^\nu g - \mathbb{E}g| \leq \sum_{k=1}^\ell \limsup_{\nu \rightarrow \infty} \sup_{g \in \mathcal{G}_{B_k}} |\mathbb{E}^\nu g - \mathbb{E}g| = 0.$$

For any $\ell \in \mathbb{N}$ and $g \in \mathcal{G}_\Xi$, observe that

$$\begin{aligned} |\mathbb{E}^\nu g - \mathbb{E}g| &\leq |\mathbb{E}^\nu[g\mathbf{1}_{A_\ell}] - \mathbb{E}[g\mathbf{1}_{A_\ell}]| + |\mathbb{E}^\nu[g\mathbf{1}_{A_\ell^c}] - \mathbb{E}[g\mathbf{1}_{A_\ell^c}]| \\ &\leq |\mathbb{E}^\nu[g\mathbf{1}_{A_\ell}] - \mathbb{E}[g\mathbf{1}_{A_\ell}]| + \mathbb{E}^\nu[L\mathbf{1}_{A_\ell^c}] + \mathbb{E}[L\mathbf{1}_{A_\ell^c}]. \end{aligned}$$

Therefore, for any $\ell \in \mathbb{N}$,

$$\text{g-lip}_X(E^\nu f - Ef) = \sup_{g \in \mathcal{G}_\Xi} |\mathbb{E}^\nu g - \mathbb{E}g| \leq \sup_{g \in \mathcal{G}_{A_\ell}} |\mathbb{E}^\nu g - \mathbb{E}g| + \mathbb{E}^\nu[L\mathbf{1}_{A_\ell^c}] + \mathbb{E}[L\mathbf{1}_{A_\ell^c}].$$

Using $\mathbb{E}[L] < \infty$, by SLLN, we have $\mathbb{E}^\nu[L\mathbf{1}_{A_\ell^c}] \rightarrow \mathbb{E}[L\mathbf{1}_{A_\ell^c}]$, $\mathbb{P}$ -a.s., as $\nu \rightarrow \infty$. Since $P(A_\ell^c) \downarrow 0$ and $\mathbb{E}[L] < \infty$, by dominated convergence theorem, we have $\mathbb{E}[L\mathbf{1}_{A_\ell^c}] \rightarrow 0$. Then, $\mathbb{P}$ -a.s., we have

$$\limsup_{\nu \rightarrow \infty} \text{g-lip}_X(E^\nu f - Ef) \leq \limsup_{\ell \rightarrow \infty} 2\mathbb{E}[L\mathbf{1}_{A_\ell^c}] = 0.$$

Since $\mathbb{E}[|f(\xi, 0)|] < \infty$, by SLLN, one has $E^\nu f(0) \rightarrow Ef(0)$, $\mathbb{P}$ -a.s. Therefore, we have

$$d_X(E^\nu f, Ef) = \max\{|E^\nu f(0) - Ef(0)|, \text{g-lip}_X(E^\nu f - Ef)\} \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

3.3.3 Proof of Theorem 3.9

Since $|I| < \infty$, after reindexing we may assume that $I = \mathbb{N}$ and that $\Xi_\ell = \emptyset$ for all $\ell > T$, for some $T \in \mathbb{N}$. Let $\{A_\ell\}_{\ell \in \mathbb{N}}$ and $\{B_\ell\}_{\ell \in \mathbb{N}}$ be as defined in the proof of Theorem 3.8. By $\cup_\ell \Xi_\ell = \Xi$, one has $P(A_T) = 1$. Let $\mathcal{H} = \{f(\cdot, 0)\} \cup \mathcal{G}_{A_T}$. Observe that L is an integrable envelope of $\mathcal{G}_{A_T}$. Let $L_2(\xi) = |f(\xi, 0)| + L(\xi)$, so that L_2 is $\mathcal{A}$ -measurable. By assumption, $\mathbb{E}[L_2^2] \leq 2\mathbb{E}[|f(\xi, 0)|^2] + 2\mathbb{E}[L^2] < \infty$; hence L_2 is a square-integrable envelope of $\mathcal{H}$. As in the proof of Theorem 3.8, the class $\mathcal{G}_{A_T}$ and also $\mathcal{H}$ are pointwise measurable. Moreover, for every $\delta > 0$, the classes

$$\mathcal{H}_\delta = \{h_1 - h_2 \mid h_1, h_2 \in \mathcal{H}, \int (h_1 - h_2)^2 dP < \delta^2\}, \quad \mathcal{H}_\infty^2 = \{(h_1 - h_2)^2 \mid h_1, h_2 \in \mathcal{H}\}$$

are pointwise measurable, by dominated convergence theorem and the square-integrable envelope L_2 . Since $A_T = \sqcup_{\ell \in [T]} B_\ell$, the collection of subgraphs in $\mathcal{H}$ is contained in

$$\mathcal{D} \cup \left\{ C_1 \cup \dots \cup C_T \cup (A_T^c \times (-\infty, 0)) \mid C_\ell \in \mathcal{C}_\ell, \ell \in [T] \right\},$$

where

$$\mathcal{D} = \left\{ \{(\xi, r) \mid r < f(\xi, 0)\} \right\}, \quad \mathcal{C}_\ell = \left\{ \{(\xi, r) \in B_\ell \times \mathbb{R} \mid r < g_{x,y}(\xi)\} \mid x, y \in D, x \neq y \right\}.$$

For each $\ell \in [T]$, the class $\mathcal{C}_\ell$ is a VC-class by the proof of Lemma 3.11, while $\mathcal{D}$ is a singleton class and hence is trivially a VC-class. By Lemma 2.2, the class of the above finite unions is a VC-class. Therefore, $\mathcal{H}$ is a VC-subgraph class. By the control of uniform entropy from [54, Theorem 2.6.7] and then invoking [54, Theorem 2.5.2], the class $\mathcal{H}$ is P -Donsker. Thus the empirical process

$$\sqrt{\nu}(\mathbb{E}^\nu h - \mathbb{E}h), \quad h \in \mathcal{H},$$

converges weakly in $\ell^\infty(\mathcal{H})$ to a centered tight Gaussian process $\mathbb{G}_P$. Since $P(A_T) = 1$, we have $\text{g-lip}_X(E^\nu f - Ef) = \sup_{g \in \mathcal{G}_\Xi} |\mathbb{E}^\nu[g \mathbf{1}_{A_T}] - \mathbb{E}[g \mathbf{1}_{A_T}]|$, $\mathbb{P}$ -a.s. Hence, one has

$$\sqrt{\nu} d_X(E^\nu f, Ef) = \sup_{h \in \mathcal{H}} |\sqrt{\nu}(\mathbb{E}^\nu h - \mathbb{E}h)|, \quad \mathbb{P}\text{-a.s.}$$

The mapping $z \mapsto \sup_{h \in \mathcal{H}} |z(h)|$ is the supremum norm on $\ell^\infty(\mathcal{H})$, hence is continuous. The continuous mapping theorem yields

$$\sqrt{\nu} d_X(E^\nu f, Ef) \rightarrow \sup_{h \in \mathcal{H}} |\mathbb{G}_P h| \quad \text{in distribution.}$$

The limit is finite almost surely because $\mathbb{G}_P$ is tight as an $\ell^\infty(\mathcal{H})$ -valued random element. Hence $\sqrt{\nu} d_X(E^\nu f, Ef) = O_{\mathbb{P}}(1)$.

3.3.4 Proof of Proposition 3.10

Let $X = [0, 1]$, $d = 1$, $\Xi = \mathbb{N}$, $\mathcal{A} = 2^{\mathbb{N}}$, and fix any $r^\nu \in (0, 1) \downarrow 0$. For any $k \in \mathbb{N}$, let $g_k : \mathbb{R} \rightarrow \mathbb{R}$ be the triangular-wave function:

$$g_k(x) = \int_0^x \mathbf{1}_{[0,1]}(t) (-1)^{\lfloor 2^k t \rfloor} dt.$$

It is easy to verify that $g_k(0) = 0$, $\text{g-lip}_X g_k = 1$, and g_k is semialgebraic. For any $K \in \mathbb{N}$ and any $(b_1, b_2, \dots) \in \mathbb{R}^{\mathbb{N}}$, there exists a point $x_K \in [0, 1]$ such that g_k is locally affine at x_K with $g'_k(x_K) = \text{sgn}(b_k)$ for each $k \leq K$ such that $b_k \neq 0$. Hence, we obtain

$$\sum_{k=1}^K |b_k| = \sum_{k=1}^K |b_k| \|g_k|_X\|_{\text{lip}} \geq \left\| \sum_{k=1}^K b_k g_k|_X \right\|_{\text{lip}} \geq \text{lip} \left(\sum_{k=1}^K b_k g_k|_X \right)(x_K) = \sum_{k=1}^K |b_k|.$$

Hence $\left\| \sum_{k=1}^K b_k g_k|_X \right\|_{\text{lip}} = \sum_{k=1}^K |b_k|$ for any $K \in \mathbb{N}$. When $\sum_{k=1}^\infty |b_k| < \infty$, for any $K_1, K_2 \in \mathbb{N}$, we have $\left\| \sum_{k=K_1}^{K_2} b_k g_k|_X \right\|_{\text{lip}} \leq \sum_{k=K_1}^{K_2} |b_k|$, so that the sequence $(\sum_{k=1}^K b_k g_k|_X)_K$ is Cauchy. Since $\text{Lip}(X)$ is complete, we obtain $\sum_{k=1}^\infty b_k g_k|_X \in \text{Lip}(X)$ with

$$\left\| \sum_{k=1}^\infty b_k g_k|_X \right\|_{\text{lip}} = \lim_{K \rightarrow \infty} \left\| \sum_{k=1}^K b_k g_k|_X \right\|_{\text{lip}} = \sum_{k=1}^\infty |b_k|.$$

Let δ_k be the Dirac measure on k , $p_k \in [0, 1]$ such that $\sum_{k=1}^{\infty} p_k = 1$, and, $P = \sum_{k=1}^{\infty} p_k \delta_k$. Let $\xi, \xi^1, \dots$ be iid $\mathbb{N}$ -valued discrete random variables on $(\Omega, \mathcal{F}, \mathbb{P})$ with the law P . For any $\nu \in \mathbb{N}$, define the empirical estimation $\mathbf{p}_k^\nu = \frac{1}{\nu} \sum_{i=1}^{\nu} \mathbf{1}_{\{\xi^i = k\}}$. Let $f : \mathbb{N} \times \mathbb{R} \rightarrow \mathbb{R}$ be $f(\xi, x) = g_\xi(x)$. Then, Assumptions 3.1, 3.2, and 3.7 hold. Note that $\sum_{k=1}^{\infty} |\mathbf{p}_k^\nu - p_k| \leq 2$, so that

$$d_X(E^\nu f, Ef) = \left\| \sum_{k=1}^{\infty} (\mathbf{p}_k^\nu - p_k) g_k|_X \right\|_{\text{lip}} = \sum_{k=1}^{\infty} |\mathbf{p}_k^\nu - p_k| \geq \sum_{k: \mathbf{p}_k^\nu = 0} |p_k|,$$

where the right-hand side is the *missing mass* as defined in [8, (1)]. Let $\mathbf{U}^\nu = \sum_{k: \mathbf{p}_k^\nu = 0} |p_k|$. Then, by [8, (2) and Proposition 4], there exists a specific law P on Ξ such that

$$\mathbb{E}[\mathbf{U}^\nu] > (\sqrt{r^\nu} + \nu^{-1/4})/2, \quad \mathbb{P}(|\mathbf{U}^\nu - \mathbb{E}[\mathbf{U}^\nu]| \geq \varepsilon) \leq 2 \exp(-\nu \varepsilon^2).$$

Let $\varepsilon = (\sqrt{r^\nu} + \nu^{-1/4})/4$ and we compute

$$\mathbb{P}\left(\mathbf{U}^\nu \leq \frac{\sqrt{r^\nu} + \nu^{-1/4}}{4}\right) \leq \mathbb{P}\left(|\mathbf{U}^\nu - \mathbb{E}[\mathbf{U}^\nu]| \geq \frac{\sqrt{r^\nu} + \nu^{-1/4}}{4}\right) \leq 2 \exp(-\sqrt{\nu}/16) \rightarrow 0,$$

as $\nu \rightarrow \infty$. Therefore, for any $M > 0$, one has $\sqrt{r^\nu} + \nu^{-1/4} > 4Mr^\nu$ eventually. Hence

$$\limsup_{\nu \rightarrow \infty} \mathbb{P}\left(d_X(E^\nu f, Ef) \leq Mr^\nu\right) \leq \limsup_{\nu \rightarrow \infty} \mathbb{P}\left(\mathbf{U}^\nu \leq \frac{\sqrt{r^\nu} + \nu^{-1/4}}{4}\right) = 0,$$

so that $d_X(E^\nu f, Ef) \neq O_{\mathbb{P}}(r^\nu)$.

4 Establishing Separability and Definability

In this section, we identify classes of functions satisfying Assumptions 3.2 and 3.7, respectively.

4.1 Separability

Recall that Assumption 3.2 requires the set $\{f(\xi, \cdot)|_X \mid \xi \in \Xi\}$ to be contained in a separable subspace of $(\text{Lip}(X), \|\cdot\|_{\text{lip}})$. The following countability condition is an immediate sufficient condition.

Proposition 4.1 (countability). *Under Assumption 3.1, if the collection $\{f(\xi, \cdot)|_X \mid \xi \in \Xi\}$ is countable, then Assumption 3.2 holds.*

Proof. Let $E = \{f(\xi, \cdot)|_X \mid \xi \in \Xi\}$. Since E is countable, the set of all finite rational linear combinations of elements of E is countable and dense in $\text{span } E$. Hence, $\text{span } E \subset \text{Lip}(X)$ is separable. $\square$

When ξ has a discrete distribution, we may restrict Ξ to its countable support. Then $\{f(\xi, \cdot)|_X \mid \xi \in \Xi\}$ is countable, and Assumption 3.2 follows from Proposition 4.1. Discrete distributions arise naturally in stochastic programming, for example, in scenario formulations [49, Section 2.4.1]. Countability also arises through *quantization*, which plays a fundamental role in applications such as signal processing [25].

Example 4.2 (quantization). *Let $J \subset \Xi$ be countable, and let $Q : \Xi \rightarrow J$ be called a quantizer. Standard examples include rounding, truncation, and other discretizations arising from finite-precision arithmetic. If the function $(\xi, x) \mapsto f(Q(\xi), x)$ satisfies Assumption 3.1, then it also satisfies Assumption 3.2.*

To further appreciate the countability condition, it is instructive to modify the negative result in Proposition 3.6 by replacing the uniform distribution on $[0, 1]$ with a discrete distribution supported on a countable set $\Xi' \subset [0, 1]$. By Proposition 4.1 and Theorem 3.3, this change alone suffices to eliminate the failure exhibited in Proposition 3.6: the function f remains unchanged, and only the law of ξ is modified.

Although perhaps evident, we note that countability of X cannot replace countability of Ξ , since the negative result in [53, Theorem 3] continues to hold when $X = \{(0, 0)\} \cup \{(0, k^{-1})\}_{k \in \mathbb{N}}$. While countability is convenient and easy to verify, separability is strictly more general. The following result provides another sufficient condition for Assumption 3.2 without requiring $\{f(\xi, \cdot)|_X \mid \xi \in \Xi\}$ to be countable.

Proposition 4.3 (continuous differentiability). *If X is compact with $0 \in X$ and $f(\xi, \cdot)$ is continuously differentiable on $\mathbb{R}^d$ for any $\xi \in \Xi$, then Assumption 3.2 holds.*

We defer the proof of Proposition 4.3 to Section 4.3.1.

4.2 Definability

We now turn to definability and examine Assumption 3.7 more closely. We illustrate its scope through three increasingly general classes of functions.

4.2.1 Jointly definable functions

We first consider functions jointly definable in an NIP structure. Let $\mathcal{M}$ be an NIP expansion of $\mathcal{R}$. A function $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ is jointly definable in $\mathcal{M}$ if $\Xi \subset \mathbb{R}^s$ for some $s \in \mathbb{N}$ and $\text{gph } f$ is definable in $\mathcal{M}$. As shown below, every jointly definable function satisfies Assumption 3.7.

Proposition 4.4 (joint definability). *Given $\Xi \subset \mathbb{R}^s$, if a function $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ is jointly definable in some NIP expansion $\mathcal{M}$ of $\mathcal{R}$, then Assumption 3.7 holds.*

Proof. Since f is definable in $\mathcal{M}$, the set $\text{gph } f$ is definable by some formula $\phi(\xi, x, r; z)$ with $\xi \in \mathbb{R}^s$, $x \in \mathbb{R}^d$, $r \in \mathbb{R}$, and parameter $z \in \mathbb{R}^k$. Let $I = \{1\}$, $\Xi_\ell = \Xi$, $\mathcal{M}_\ell = \mathcal{M}$, $\phi_\ell(x, r; (\xi, z)) = \phi(\xi, x, r; z)$, $k_\ell = s + k$, $z_{\ell, \xi} = (\xi, z) \in \mathbb{R}^{k_\ell}$ for $\ell \in I$. Then Assumption 3.7 holds. $\square$

Functions jointly definable in an expansion of $\mathcal{R}$ enjoy robust closure properties. In particular, joint definability is preserved under restriction to definable sets, finite Cartesian products, coordinate projections, composition with definable maps, pointwise sums and products, and pointwise maxima and minima; see [55, Chapter 1, §2]. These elementary closure properties do not require NIP. The NIP assumption is nevertheless mild and, by Proposition 2.5, is satisfied in particular by every o-minimal structure.

Corollary 4.5 (joint o-minimality). *Let $(\Xi, \mathcal{A})$ be a measurable space, and let $\xi, \xi^1, \xi^2, \dots$ be Ξ -valued iid random variables on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Suppose that $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $X \subset \mathbb{R}^d$ satisfy Assumption 3.1. If f is jointly definable in an o-minimal expansion of $\mathcal{R}$, then*

$$\lim_{\nu \rightarrow \infty} d_X(E^\nu f, Ef) = 0, \quad \mathbb{P}\text{-a.s.}$$

If, in addition, $\mathbb{E}[|f(\xi, 0)|^2] < \infty$ and $\mathbb{E}[L^2] < \infty$, then for any $\delta > 0$, there exists a constant $C_\delta < \infty$ such that for all sufficiently large $\nu \in \mathbb{N}$, with probability at least $1 - \delta$, one has

$$d_X(E^\nu f, Ef) \leq C_\delta \nu^{-1/2}.$$

Proof. The claim follows from Propositions 2.5 and 4.4, Theorems 3.8 and 3.9. $\square$

As shown in [21, Section 5.2], [9, Section 6.2], and [5], the training losses of many modern deep learning models are jointly definable in an o-minimal structure, notably the real exponential field $\mathcal{R}_{\text{exp}}$. Hence, Theorem 3.8 applies under mild integrability and Lipschitz assumptions. We emphasize, however, that joint definability of f does not imply definability of the expectation function Ef , which may require additional continuity and definability assumptions on the distribution P .

4.2.2 Functions with uniformly definable slices

Joint definability requires Ξ to be a subset of a Euclidean space, which may be inconvenient or unnecessary. In this subsection, we show that uniform definability, corresponding to Assumption 3.7 with $|I| = 1$, provides a natural generalization of joint definability. We begin with the following sufficient condition.

Proposition 4.6. *A function $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ satisfies Assumption 3.7 if there exist a function $Z : \Xi \rightarrow \mathbb{R}^k$ and a jointly definable function $g : \mathbb{R}^k \times \mathbb{R}^d \rightarrow \mathbb{R}$ on an NIP expansion $\mathcal{M}$ of $\mathcal{R}$ such that $f(\xi, x) = g(Z(\xi), x)$ for all $(\xi, x) \in \Xi \times \mathbb{R}^d$.*

Proof. Let $\Xi_\ell = \Xi$ and $\mathcal{M}_\ell = \mathcal{M}$ for $\ell \in I = \{1\}$. Since g is jointly NIP definable, there exists a formula $\phi(x, r, z; b)$ with $x \in \mathbb{R}^d, r \in \mathbb{R}, z \in \mathbb{R}^k$, and parameter $b \in \mathbb{R}^p$ such that $\text{gph } g = \{(z, x, r) \mid \mathcal{M} \models \phi(x, r, z; b)\}$. Let $\phi_\ell = \phi$. For any $\xi \in \Xi$, we have $\text{gph } f(\xi, \cdot) = \text{gph } g(Z(\xi), \cdot) = \{(x, r) \mid \mathcal{M} \models \phi(x, r, Z(\xi); b)\}$, hence $f(\xi, \cdot)$ is definable by the formula ϕ with parameter $(Z(\xi), b)$. $\square$

It is worth noting that the function $Z : \Xi \rightarrow \mathbb{R}^k$ may be arbitrary and need not be definable in any structure; consequently, Ξ need not be a Euclidean subset. Indeed, even when $\Xi = \mathbb{R}$, the class of functions satisfying the assumptions of Proposition 4.6 already contains functions that are not jointly definable in any NIP structure, as the following example shows.

Example 4.7 (uniformly definable but not jointly definable). *Consider $f(\xi, x) = \mathbf{1}_{\mathbb{Z}}(\xi)x^2$. Let $Z(\xi) = \mathbf{1}_{\mathbb{Z}}(\xi)$ and define $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $g(t, x) = tx^2$. Then $f(\xi, x) = g(Z(\xi), x)$. By Proposition 4.6, Assumption 3.7 holds, since g is semialgebraic. However, the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is not jointly definable in any NIP expansion of $\mathcal{R}$. Indeed, if f were definable in such an expansion, then $\mathbb{Z} = \{\xi \in \mathbb{R} \mid f(\xi, 1) = 1\}$ would also be definable, contradicting Proposition 2.6.*

As in the jointly definable case, an analogue of Corollary 4.5 holds when the family $\{f(\xi, \cdot) \mid \xi \in \Xi\}$ is uniformly definable. In particular, one obtains convergence of $d_X(E^\nu f, Ef)$ and, under the additional conditions $\mathbb{E}[|f(\xi, 0)|^2] < \infty$ and $\mathbb{E}[L^2] < \infty$, the rate $O_{\mathbb{P}}(\nu^{-1/2})$. We omit the formal statement for brevity.

4.2.3 Functions with piecewise uniformly definable slices

We now consider the full generality of Assumption 3.7. It extends uniform definability by covering Ξ with sets $\{\Xi_\ell\}_{\ell \in I}$ and requiring that, for each $\ell \in I$, the restrictions of the family $\{f(\xi, \cdot) \mid \xi \in \Xi_\ell\}$ to $\{q_k\}_{k \in \mathbb{N}}$ be uniformly definable in some NIP structure $\mathcal{M}_\ell$.

When $|I| < \infty$, Theorems 3.8 and 3.9 yield convergence and rate estimates analogous to those in Corollary 4.5. As the following example shows, piecewise uniform definability is strictly more general than uniform definability, even when $|I| = 2$.

Example 4.8 (piecewise uniformly definable but not uniformly). *Consider the NIP expansions $\mathcal{M}_1 = (\mathbb{R}, +, \cdot, 0, 1, <, 2^{\mathbb{Z}})$ and $\mathcal{M}_2 = (\mathbb{R}, +, \cdot, 0, 1, <, 3^{\mathbb{Z}})$; see [26, Theorem 6.5]. Let $\Xi = \{1, 2\}$, $\Xi_1 = \{1\}$, and $\Xi_2 = \{2\}$. Define $f(1, x) = \text{dist}(x, \{0\} \cup 2^{\mathbb{Z}})$ and $f(2, x) = \text{dist}(x, \{0\} \cup 3^{\mathbb{Z}})$. Then $f(\xi, \cdot)$ is uniformly definable in $\mathcal{M}_1$ when $\xi \in \Xi_1$ and uniformly definable in $\mathcal{M}_2$ when $\xi \in \Xi_2$. Hence, Assumption 3.7 holds. However, f is not uniformly definable in any single NIP expansion $\mathcal{M}$ of $\mathcal{R}$. Indeed, if both $f(1, \cdot)$ and $f(2, \cdot)$ were definable in $\mathcal{M}$, then their zero sets would be definable, and hence so would $2^{\mathbb{Z}}$ and $3^{\mathbb{Z}}$. Since $\log_2 3 \notin \mathbb{Q}$, [27, Theorem 1.3] implies that $\mathcal{M}$ defines $\mathbb{Z}$. By Proposition 2.6, $\mathcal{M}$ then has IP, contradicting NIP.*

When I is countably infinite, countable decompositions $\Xi = \cup_{\ell \in I} \Xi_\ell$ are closely related to nonuniform learnability in learning theory [45, Chapter 7] and are standard in empirical process theory [57, 58]. In this setting, Theorem 3.8 still yields $d_X(E^\nu f, Ef) \rightarrow 0$. However, unlike in the finite-piece case $|I| < \infty$, Proposition 3.10 shows that this convergence can be arbitrarily slow.

A natural question in Assumption 3.7 is how to choose the pieces $\{\Xi_\ell\}_{\ell \in I}$. This choice may be evident in a scenario formulation but is less apparent in general. In the sequel, we show that, for a broad class of functions, there is a universal construction that exploits the countability of the languages of the structures $\{\mathcal{M}_\ell\}_{\ell \in I}$. Our result requires the following assumption, based on the notion of a *totally Borel* structure; see [51, Definition 1.1.1].

Assumption 4.9 (slicewise definability). *Given a measurable space $(\Xi, \mathcal{A})$ and a function $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$, assume that there exists a family $\{\mathcal{M}_\ell\}_{\ell \in I}$ of NIP expansions of $\mathcal{R}$ indexed by $I \subset \mathbb{N}$, such that, for any $\ell \in I$,*

- (i) *the language of $\mathcal{M}_\ell$ is countable, and*
- (ii) *$\mathcal{M}_\ell$ is totally Borel, meaning that every set definable in $\mathcal{M}_\ell$ is Borel.*

Moreover, for any $\xi \in \Xi$, $f(\xi, \cdot)$ is definable in $\mathcal{M}_\ell$ for some $\ell \in I$.

In Assumption 4.9, neither an explicit cover $\{\Xi_\ell\}_{\ell \in I}$ nor the uniform definability required in Assumption 3.7 is assumed. Instead, for each $\xi \in \Xi$, the slice $f(\xi, \cdot)$ is definable in some totally Borel structure $\mathcal{M}_\ell$ in a countable language, with $\ell \in I$, without any explicit uniformity in ξ .

Theorem 4.10 (slicewise definability). *Let $(\Xi, \mathcal{A})$ be a measurable space, and let $\xi, \xi^1, \xi^2, \dots$ be Ξ -valued iid random variables on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Suppose that $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $X \subset \mathbb{R}^d$ satisfy Assumptions 3.1 and 4.9. Then*

$$\lim_{\nu \rightarrow \infty} d_X(E^\nu f, Ef) = 0, \quad \mathbb{P}\text{-a.s.}$$

The proof of Theorem 4.10 is deferred to Section 4.3.2 and mainly verifies Assumption 3.7. Informally, we enumerate all formulas arising from the structures $\{\mathcal{M}_\ell\}_{\ell \in I}$ and, after reindexing, define Ξ_ℓ as the set of ξ whose value sequence $\{f(\xi, q_j)\}_{j \in \mathbb{N}}$ can be represented by the ℓ th formula, where $\{q_j\}_{j \in \mathbb{N}}$ is a countable dense subset of X . This enumeration is possible because I is countable and each $\mathcal{M}_\ell$ has a countable language. The proof then concludes by showing that each Ξ_ℓ is universally measurable relative to $\mathcal{A}$, using the totally Borel hypothesis.

By [30, Proposition 1.1], every o-minimal structure is totally Borel. Hence any countable-language o-minimal structure is admissible for Assumption 4.9; important examples include the real ordered field $\mathcal{R}$ and the real exponential field $\mathcal{R}_{\text{exp}}$. The restricted analytic field $\mathcal{R}_{\text{an}}$ defines

globally subanalytic functions. Although its language contains a symbol for every restricted analytic function and is therefore uncountable, most functions arising in practice involve only countably many such functions across all slices. Retaining only these functions together with the ordered-field language yields a countable-language reduct of $\mathcal{R}_{\text{an}}$, which is o-minimal.

Corollary 4.11 (countable-language o-minimal slices). *Let $(\Xi, \mathcal{A})$ be a measurable space, and let $\xi, \xi^1, \xi^2, \dots$ be Ξ -valued iid random variables on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Let $\mathcal{M}$ be a countable-language o-minimal expansion of $\mathcal{R}$. Suppose that $f : \Xi \times \mathbb{R}^d \rightarrow \mathbb{R}$ and $X \subset \mathbb{R}^d$ satisfy Assumption 3.1 and, for any $\xi \in \Xi$, the function $f(\xi, \cdot)$ is definable in the structure $\mathcal{M}$. Then*

$$\lim_{\nu \rightarrow \infty} d_X(E^\nu f, Ef) = 0, \quad \mathbb{P}\text{-a.s.}$$

Proof. The claim follows from [30, Proposition 1.1], Proposition 2.5, and Theorem 4.10. $\square$

Remark 4.12. *Much of the nonsmooth optimization literature invokes o-minimality to rule out pathological behavior; see, e.g., [10, 21, 11]. In Corollary 4.11, and more generally in Theorem 4.10, however, the convergence of $d_X(E^\nu f, Ef)$ follows from a combinatorial property, namely NIP, which every o-minimal structure possesses. By contrast, the works cited above rely primarily on geometric properties, expressed through stratifiability in various senses, to establish convergence of subgradient-type methods. Such stratifiability is neither necessary nor sufficient for convergence in the Lipschitz pseudometric, even when the number of strata is uniformly bounded; see Section 4.3.3 for details. This distinction highlights the different roles of combinatorial and geometric regularity.*

Remark 4.13. *In Theorem 4.10, two countability assumptions are imposed: the index set I is countable, and each structure $\mathcal{M}_\ell$ has a countable language. Our proof relies essentially on both assumptions to enumerate all formulas arising from the family $\{\mathcal{M}_\ell\}_{\ell \in I}$. It is natural to ask whether either assumption can be removed. We conjecture that neither assumption can be dispensed with without additional hypotheses.*

4.3 Proofs

We collect the deferred proofs and details from Sections 4.1 and 4.2.

4.3.1 Proof of Proposition 4.3

By compactness of X , the set $E = \{h|_X \mid h \in C^1(\mathbb{R}^d)\}$ is a subspace of $(\text{Lip}(X), \|\cdot\|_{\text{lip}})$. Since $\{f(\xi, \cdot)|_X \mid \xi \in \Xi\} \subset E$, it suffices to show that E is separable. Let $Y = \text{con } X$, which is also compact. For $h, g \in C^1(\mathbb{R}^d)$, by the mean value theorem, $\text{g-lip}_X(h - g) \leq \sup_{y \in Y} \|\nabla h(y) - \nabla g(y)\|_*$. Hence $\|h|_X - g|_X\|_{\text{lip}} \leq \max\{|h(0) - g(0)|, \sup_{y \in Y} \|\nabla h(y) - \nabla g(y)\|_*\}$. Define the linear map $\Phi : C^1(\mathbb{R}^d) \rightarrow \mathbb{R} \times C(Y; \mathbb{R}^d)$ by $\Phi(h) = (h(0), \nabla h|_Y)$. Since Y is compact with Euclidean metric, $C(Y; \mathbb{R}^d)$ is separable [1, Lemma 3.99]. Therefore $\mathbb{R} \times C(Y; \mathbb{R}^d)$ with norm $(a, b) \mapsto \max\{|a|, \|b\|_\infty\}$ is separable, where $\|b\|_\infty = \sup_{y \in Y} \|b(y)\|_*$. The image $\Phi(C^1(\mathbb{R}^d))$ is a subset of the separable metric space $\mathbb{R} \times C(Y; \mathbb{R}^d)$ and is therefore separable [1, Corollary 3.5]. Let H be a countable subset of $C^1(\mathbb{R}^d)$ such that $\Phi(H)$ is dense in $\Phi(C^1(\mathbb{R}^d))$. For any $h \in C^1(\mathbb{R}^d)$, there exist $\{h_k\}_{k \in \mathbb{N}} \subset H$ such that $\Phi(h_k) \rightarrow \Phi(h)$ in the above norm, which, by the above inequality, implies $\|h_k|_X - h|_X\|_{\text{lip}} \rightarrow 0$. Hence $\{h|_X \mid h \in H\}$ is countable dense in E , and E is a separable subset of $(\text{Lip}(X), \|\cdot\|_{\text{lip}})$. Thus Assumption 3.2 holds.

4.3.2 Proof of Theorem 4.10

Since the structure $\mathcal{M}_\ell$ has countable language, the set of formulas in the language of $\mathcal{M}_\ell$ is countable; see [14, Proposition 1.3.4]. Hence, the set of all possible formulas in the structures $\{\mathcal{M}_\ell\}_{\ell \in I}$ is also countable. We can therefore enumerate all relevant formulas that have at least $d+1$ variables as $\{\phi_i(x, r; z_i)\}_{i \in \mathbb{N}}$, where $x \in \mathbb{R}^d, r \in \mathbb{R}, z_i \in \mathbb{R}^{k_i}, k_i \in \{0\} \cup \mathbb{N}$, and ϕ_i is defined in the language of $\mathcal{M}_{\ell(i)}$ with $\ell(i) \in \mathbb{N}$. Let $\{q_j\}_{j \in \mathbb{N}}$ be a countable dense subset of X . We then verify Assumption 3.7 and the conclusion follows from Theorem 3.8.

To this end, for each $i \in \mathbb{N}$, we focus on the formula ϕ_i and let

$$Z_i = \bigcap_{j=1}^{\infty} \{z \in \mathbb{R}^{k_i} \mid \mathcal{M}_{\ell(i)} \models (\exists! r) \phi_i(q_j, r; z)\},$$

where the quantifier $\exists!$ denotes unique existence and is definable in first-order logic. Hence, the set $\{z \in \mathbb{R}^{k_i} \mid \mathcal{M}_{\ell(i)} \models (\exists! r) \phi_i(q_j, r; z)\}$ is definable in $\mathcal{M}_{\ell(i)}$. Since $\mathcal{M}_{\ell(i)}$ is totally Borel, Z_i is Borel. For any $i, j \in \mathbb{N}$, define $G_{i,j} = \{(v, z) \in \mathbb{R}^{\mathbb{N}} \times Z_i \mid \mathcal{M}_{\ell(i)} \models \phi_i(q_j, v_j, z)\}$ and

$$H_i = \bigcap_{j=1}^{\infty} G_{i,j} = \left\{ (v, z) \in \mathbb{R}^{\mathbb{N}} \times Z_i \mid \mathcal{M}_{\ell(i)} \models \phi_i(q_j, v_j, z), \forall j \in \mathbb{N} \right\},$$

where the space $\mathbb{R}^{\mathbb{N}}$ is endowed with the product topology. Let $E_{i,j} = \{(v_j, z) \in \mathbb{R} \times \mathbb{R}^{k_i} \mid \mathcal{M}_{\ell(i)} \models \phi_i(q_j, v_j, z)\}$, which is definable in $\mathcal{M}_{\ell(i)}$, and therefore, is Borel. The set $G_{i,j}$ is the preimage of a Borel set $E_{i,j} \cap (\mathbb{R} \times Z_i)$ under the continuous coordinate projection $(v, z) \mapsto (v_j, z)$, so that $G_{i,j}$, and also H_i , are Borel on $\mathbb{R}^{\mathbb{N}} \times \mathbb{R}^{k_i}$. Since $\mathbb{R}^{\mathbb{N}}$ and $\mathbb{R}^{k_i}$ are Polish, using [17, Proposition 8.4.4], the following set is universally measurable relative to $\mathcal{B}(\mathbb{R}^{\mathbb{N}})$:

$$V_i = \{v \in \mathbb{R}^{\mathbb{N}} \mid \exists z \in Z_i, (v, z) \in H_i\},$$

which, simply put, is the collection of all value sequences on $\{q_j\}_{j \in \mathbb{N}}$ realized by the formula ϕ_i with parameters that define a unique value at every q_j . Consider the following mapping

$$\Psi : \Xi \longrightarrow \mathbb{R}^{\mathbb{N}}, \quad \xi \longmapsto (f(\xi, q_1), f(\xi, q_2), \dots),$$

which is $\mathcal{A}/\mathcal{B}(\mathbb{R}^{\mathbb{N}})$ -measurable since $f(\cdot, x)$ is $\mathcal{A}$ -measurable for all $x \in \mathbb{R}^d$; see [23, Proposition 2.4]. For each $i \in \mathbb{N}$, define

$$\Xi_i = \Psi^{-1}(V_i) \subset \Xi.$$

By using [17, Lemma 8.4.6], the set Ξ_i is universally measurable relative to $\mathcal{A}$.

By construction, for any $i \in \mathbb{N}$ and any $\xi \in \Xi_i$, there exists $z \in Z_i$ such that $\mathcal{M}_{\ell(i)} \models \phi_i(q_j, f(\xi, q_j); z)$ for all $j \in \mathbb{N}$. Meanwhile, by construction of Z_i , we have $\mathcal{M}_{\ell(i)} \models \phi_i(q_j, r; z)$ if and only if $r = f(\xi, q_j)$.

To show $\cup_i \Xi_i = \Xi$, suppose not. Then, there exists $\xi \in \Xi \setminus (\cup_i \Xi_i)$. By assumption, the function $f(\xi, \cdot)$ is definable in some structure in $\{\mathcal{M}_\ell\}_{\ell \in I}$. Hence, there exists $i \in \mathbb{N}, z \in \mathbb{R}^{k_i}$, and a formula ϕ_i such that $\text{gph } f(\xi, \cdot) = \phi_i(\mathbb{R}^{d+1}; z)$, so that $\xi \in \Xi_i$, a contradiction.

4.3.3 Missing details in Remark 4.12

Let $\mathcal{T}_S(x)$ denote the tangent cone to a set S at x . When S is a C^1 manifold, $\mathcal{T}_S(x)$ coincides with the tangent space to S at x ; see [42, Example 6.8]. We begin by recalling the definition of Whitney C^p -stratification used in [10, 21].

Definition 4.14. A Whitney C^p -stratification of $S \subset \mathbb{R}^d$ is a finite partition of S into connected, locally closed, embedded C^p -submanifolds S_k , called strata, satisfying the following conditions.

- (i) **(Frontier)** For $i \neq j$, if $S_i \cap \text{cl } S_j \neq \emptyset$, then $S_i \subset \text{cl } S_j$.
- (ii) **(Whitney-(a))** Fix $i \neq j$. If $S_i \ni x_k \rightarrow y \in S_j$ and $\mathcal{T}_{S_i}(x_k) \rightarrow T$, one has $\mathcal{T}_{S_j}(y) \subset T$.

Here, the convergence of tangent spaces is understood in the Grassmannian topology. We next show that Whitney C^p -stratifiability is neither necessary nor sufficient for convergence in the Lipschitz pseudometric. Since one may take $f(\xi, x) = g(x)$, where g is deterministic and essentially arbitrary, it is immediate that stratifiability is not necessary for the convergence of $d_X(E^\nu f, Ef)$. As the following analysis of the construction in Proposition 3.6 shows, it is not sufficient either, even when the number of strata is uniformly bounded.

We first recall the construction of $f : \Xi \times \mathbb{R}^2 \rightarrow \mathbb{R}$ in [53, Theorem 3] below:

$$f(\xi, x) = \max\{g(\xi, x), 0\} + 35\|x\|_2^2,$$

where $g(\xi, x) = x_1 + q(\xi, x_2)$ with $q : \Xi \times \mathbb{R} \rightarrow \mathbb{R}$. Moreover, for any $\xi \in \Xi$, the function $q(\xi, \cdot)$ is $C^{1,1}$ on $\mathbb{R}$ and C^2 on $\mathbb{R} \setminus \{0\}$. Without loss of generality, we remove the quadratic term and focus on $r(\xi, x) = \max\{g(\xi, x), 0\}$. Let $I_- = (-\infty, 0)$, $I_0 = \{0\}$, $I_+ = (0, \infty)$, and a piecewise affine $h : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $h(y_1, y_2) = \max\{y_1, 0\}$. Then, it is easy to verify that $\text{gph } h$ admits a Whitney C^∞ -stratification. In particular, $\text{gph } h$ is disjoint union of strata $\mathcal{P} = \{P_{\alpha, \beta}\}_{\alpha, \beta \in \{-, 0, +\}}$, where

$$P_{\alpha, \beta} = \{(y_1, y_2, h(y_1, y_2)) \mid y_1 \in I_\alpha, y_2 \in I_\beta\}.$$

For any $\xi \in \Xi$, define the following mapping $\Phi_\xi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ as

$$\Phi_\xi(x_1, x_2, t) = (x_1 + q(\xi, x_2), x_2, t),$$

which is a C^1 -diffeomorphism. Observe that $\Phi_\xi(\text{gph } r(\xi, \cdot)) = \text{gph } h$ and Φ_ξ is C^2 on $\mathbb{R} \times (\mathbb{R} \setminus \{0\}) \times \mathbb{R}$. For any $\xi \in \Xi$, consider the collection $\mathcal{Q}_\xi = \{Q_{\xi, \alpha, \beta}\}_{\alpha, \beta \in \{-, 0, +\}}$, where

$$Q_{\xi, \alpha, \beta} = \{(x_1, x_2, r(\xi, x_1, x_2)) \mid x_1 + q(\xi, x_2) \in I_\alpha, x_2 \in I_\beta\} = \Phi_\xi^{-1}(P_{\alpha, \beta}).$$

The collection $\mathcal{Q}_\xi$ forms a finite partition of $\text{gph } r(\xi, \cdot)$ into connected, locally closed submanifolds, and the frontier condition is inherited from the stratification of h because Φ_ξ is a homeomorphism. The collection $\mathcal{Q}_\xi$ consists of C^2 submanifolds and has at most nine elements for every $\xi \in \Xi$. Thus, $\mathcal{Q}_\xi$ is a finite C^2 -stratification.

It remains to check Whitney-(a). Fix two distinct indices (α, β) and (α', β') . The only nontrivial case is when $Q_{\xi, \alpha, \beta} \subset \text{cl } Q_{\xi, \alpha', \beta'}$, or equivalently, since Φ_ξ is a homeomorphism, $P_{\alpha, \beta} \subset \text{cl } P_{\alpha', \beta'}$. Let $V_{\alpha, \beta}$ and $V_{\alpha', \beta'}$ denote the constant tangent spaces of $P_{\alpha, \beta}$ and $P_{\alpha', \beta'}$, respectively. Since $\mathcal{P}$ is the Whitney stratification of $\text{gph } h$, we have $V_{\alpha, \beta} \subset V_{\alpha', \beta'}$. Take any sequence $z_k \in Q_{\xi, \alpha', \beta'}$ such that $z_k \rightarrow z \in Q_{\xi, \alpha, \beta}$, and suppose $\mathcal{T}_{Q_{\xi, \alpha', \beta'}}(z_k) \rightarrow T$. Write $z_k = \Phi_\xi^{-1}(a_k)$ and $z = \Phi_\xi^{-1}(b)$, where $a_k \in P_{\alpha', \beta'}$ and $b \in P_{\alpha, \beta}$. Since Φ_ξ is a homeomorphism, $a_k \rightarrow b$. Moreover, since Φ_ξ^{-1} is C^1 ,

$$\mathcal{T}_{Q_{\xi, \alpha', \beta'}}(z_k) = \nabla \Phi_\xi^{-1}(a_k) V_{\alpha', \beta'} \rightarrow \nabla \Phi_\xi^{-1}(b) V_{\alpha', \beta'}.$$

Hence $T = \nabla \Phi_\xi^{-1}(b) V_{\alpha', \beta'}$. On the other hand, $\mathcal{T}_{Q_{\xi, \alpha, \beta}}(z) = \nabla \Phi_\xi^{-1}(b) V_{\alpha, \beta}$. Because $\nabla \Phi_\xi^{-1}(b)$ is an invertible linear map and $V_{\alpha, \beta} \subset V_{\alpha', \beta'}$, we obtain

$$\mathcal{T}_{Q_{\xi, \alpha, \beta}}(z) = \nabla \Phi_\xi^{-1}(b) V_{\alpha, \beta} \subset \nabla \Phi_\xi^{-1}(b) V_{\alpha', \beta'} = T.$$

This verifies Whitney-(a).

5 Applications

In the preceding sections, we studied the convergence of $E^\nu f$ to Ef in the Lipschitz pseudometric. We now examine its consequences and properties, particularly the implications for other notions of convergence. For simplicity, we use the Euclidean norm $\|\cdot\|_2$ throughout this section.

5.1 Uniform convergence of function values

Uniform convergence of function values ensures consistency of global solutions in stochastic programming; see [49, Section 9.2.6]. We show below that convergence in the Lipschitz pseudometric controls the uniform distance on bounded sets.

Proposition 5.1. *Let $0 \in X \subset \mathbb{R}^d$ and $h, g : \mathbb{R}^d \rightarrow \mathbb{R}$. For any bounded set $U \subset X$, one has*

$$\sup_{x \in U} |h(x) - g(x)| \leq (1 + \sup_{z \in U} \|z\|_2) d_X(h, g).$$

Proof. For any $x \in U$, we have

$$|h(x) - g(x)| \leq |h(0) - g(0)| + \text{g-lip}_X(h - g) \|x\|_2 \leq (1 + \|x\|_2) d_X(h, g).$$

The claim follows by taking the supremum over U . $\square$

The boundedness requirement cannot be removed. Indeed, let $d = 1$, $X = \mathbb{R}$, $g^\nu(x) = \frac{1}{\nu}|x|$, and $g \equiv 0$. Then, one has $d_X(g^\nu, g) = \frac{1}{\nu} \rightarrow 0$, whereas $\sup_{x \in X} |g^\nu(x) - g(x)| = \infty$ for every $\nu \in \mathbb{N}$. Applying Proposition 5.1 with $h = E^\nu f$ and $g = Ef$ shows that a Lipschitzian SLLN implies a uniform SLLN for function values on every bounded subset of X .

5.2 Uniform convergence of subdifferentials

We next show that the Lipschitz pseudometric controls first-order approximations, including subdifferentials. For a locally Lipschitz function $h : \mathbb{R}^d \rightarrow \mathbb{R}$, several notions of subdifferential are available; see [42, Chapter 8]. In particular, the regular, limiting, and Clarke subdifferentials are defined by [42, Definition 8.3 and 8(32)]

$$\begin{aligned} \widehat{\partial}h(x) &= \{v \in \mathbb{R}^d \mid h(x_k) \geq h(x) + \langle v, x_k - x \rangle + o(\|x_k - x\|_2) \text{ as } x_k \rightarrow x\}, \\ \partial h(x) &= \{v \in \mathbb{R}^d \mid \exists x_k \rightarrow x, h(x_k) \rightarrow h(x), \exists v_k \rightarrow v, v_k \in \widehat{\partial}h(x_k)\}, \\ \overline{\partial}h(x) &= \text{con } \partial h(x). \end{aligned}$$

We show below that the Hausdorff distances between the corresponding limiting and Clarke subdifferentials are uniformly controlled by the Lipschitz pseudometric.

Proposition 5.2 (subdifferentials). *Let $h, g : \mathbb{R}^d \rightarrow \overline{\mathbb{R}}$ be locally Lipschitz on an open set $X \subset \mathbb{R}^d$ with $0 \in X$. Then, we have*

$$\sup_{x \in X} d(\overline{\partial}h(x), \overline{\partial}g(x)) \leq \sup_{x \in X} d(\partial h(x), \partial g(x)) \leq \sup_{x \in X} \text{lip}(h - g)(x) \leq d_X(h, g).$$

Proof. Fix $x \in X$. For the limiting case, by [42, Corollary 10.9] and the fact that h, g are locally Lipschitz on X , one has $\partial h(x) \subset \partial g(x) + \partial(h - g)(x)$. Moreover, by [42, Theorem 9.13], $\partial(h - g)(x) \subset \text{lip}(h - g)(x)\mathbb{B}$, hence $\text{exs}(\partial h(x); \partial g(x)) \leq \text{lip}(h - g)(x)$. The claim follows by interchanging h and g , taking the supremum over $x \in X$, and applying Lemma 2.1. For the Clarke case, using [29, Proposition 1.4], we obtain $d(\overline{\partial}h(x), \overline{\partial}g(x)) \leq d(\partial h(x), \partial g(x))$ for any $x \in X$. $\square$

Notably, the set X in Proposition 5.2 need not be bounded. Applying Proposition 5.2 to $E^\nu f$ and Ef and invoking the results of Section 3, we obtain the following uniform SLLN for subdifferentials over a possibly unbounded set.

Corollary 5.3. *Suppose that the assumptions of one of Theorems 3.3, 3.8 or 4.10 hold. Then, for every set $U \subset \text{int } X$, $\mathbb{P}$ -a.s.,*

$$\lim_{\nu \rightarrow \infty} \sup_{x \in U} d(\partial E^\nu f(x), \partial Ef(x)) = 0 \quad \text{and} \quad \lim_{\nu \rightarrow \infty} \sup_{x \in U} d(\bar{\partial} E^\nu f(x), \bar{\partial} Ef(x)) = 0.$$

Suppose instead that the assumptions of Theorem 3.9 hold. Then, for every $\delta > 0$, there exists $C_\delta < \infty$ such that, for all sufficiently large $\nu \in \mathbb{N}$, there is an event $\Omega_{\delta, \nu} \subset \Omega$ with $\mathbb{P}(\Omega_{\delta, \nu}) \geq 1 - \delta$ on which

$$\sup_{x \in U} d(\bar{\partial} E^\nu f(x), \bar{\partial} Ef(x)) \leq \sup_{x \in U} d(\partial E^\nu f(x), \partial Ef(x)) \leq C_\delta \nu^{-1/2}.$$

Since both the limiting and Clarke subdifferential mappings are outer semicontinuous, the desired consistency follows directly from Corollary 5.3. Uniform SLLNs for Clarke subdifferentials have been studied extensively, whereas substantially less is known for limiting subdifferentials. A uniform law for enlarged Clarke subdifferentials was established in [47] and shown in [40], under suitable conditions, to be equivalent to a graphical convergence result. Whether the enlargement could be removed under Assumption 3.1 was posed as an open question in [47, Remark 2]. This question was later answered negatively in [53], even for stochastic convex functions.

Uniform laws without enlargement have nevertheless been established for several special classes of functions; see [49, p. 449] and [52, 43]. In particular, the work [52] introduced a separability-type condition on subdifferential mappings that guarantees uniform convergence of Clarke subdifferentials. Our Theorem 3.3 may be viewed as similar to, but complementary to, [52] in the setting of Clarke subdifferentials. However, the results in [52] do not extend directly to limiting subdifferentials, whose values need not be convex. More recently, the work [43] established a uniform law for a class of convex-composite functions under a finite-VC-dimension assumption on the inner smooth mapping. Its scope, however, is restricted to subdifferentially regular functions and therefore does not cover the more general setting considered here.

Another related work [44] studies limit fields generated by gradients, Clarke Jacobians, and more general conservative mappings. Under joint definability of the parametrized family in a fixed o-minimal structure, the limit field admits a variational stratification. As noted in [44, Remark 4.6], slicewise definability alone is insufficient for their conclusion. In contrast, Theorem 4.10 requires only slicewise definability of f , provided that the relevant structures have countable languages. Recently, the work [2] develops an algorithmic approach to finding stationary points of stochastic convex problems without resorting to surrogate stationarity measures, such as small gradients of the Moreau envelope. In this sense, their approach avoids the need for a uniform SLLN for subdifferentials, but fundamentally relies on convexity.

A natural question is whether an analogue of Corollary 5.3 holds for regular subdifferentials. Such a result is delicate and may fail in general because the regular subdifferential can be empty. Nevertheless, arguments similar to those in Section 3.3 could yield uniform convergence for other first-order approximation objects, including generalized directional derivatives and metric slopes.

We conclude this subsection with the following result, which extends [43, Theorem 1] to more general Lipschitz functions and shows that the bound in terms of the local Lipschitz modulus in Proposition 5.2 is sharp for continuously differentiable functions.

Proposition 5.4. *Suppose that $h, g : \mathbb{R}^d \rightarrow \mathbb{R}$ are Lipschitz on an open $X \subset \mathbb{R}^d$. Let $D \subset X$ be the set of points where h and g are both differentiable. Then, D has full measure and*

$$\sup_{x \in X} d(\bar{\partial}h(x), \bar{\partial}g(x)) \leq \sup_{x \in X} \text{lip}(h - g)(x) \leq \sup_{x \in D} \|\nabla h(x) - \nabla g(x)\|_2.$$

If h and g are continuously differentiable on X , then both inequalities are equalities.

Proof. By Rademacher's theorem [42, Theorem 9.60], the set $D = \text{dom}(\nabla h) \cap \text{dom}(\nabla g) \cap X$ has full Lebesgue measure in X . Let $f = h - g$, hence f is Lipschitz on X and $\nabla f(x) = \nabla h(x) - \nabla g(x)$ for any $x \in D$. Using [42, Theorem 9.61], for any $y \in X$, one has $\sup_{v \in \bar{\partial}f(y)} \|v\|_2 \leq \sup_{x \in D} \|\nabla f(x)\|_2 = \sup_{x \in D} \|\nabla h(x) - \nabla g(x)\|_2$. By [42, Theorem 9.13], we obtain $\text{lip} f(y) = \sup_{v \in \bar{\partial}f(y)} \|v\|_2$, which completes the proof. $\square$

5.3 Finite-sample identification of solutions

Finite-sample identification in stochastic programming [46, 32, 48] is another application of our Lipschitzian SLLNs. Under mild assumptions, almost surely, every accumulation point of global solutions to the SAA problems in (2) is a global solution to the true problem in (1); see [3, Theorem 2.3]. A stronger finite-sample identification phenomenon is discussed in [49, Section 5.33]: under suitable polyhedral convexity and sharpness assumptions, and when the distribution has finite support, for all sufficiently large $\nu \in \mathbb{N}$,

$$\text{argmin}_{x \in X} E^\nu f(x) \subset \text{argmin}_{x \in X} Ef(x), \quad \mathbb{P}\text{-a.s.}$$

Thus, after finitely many samples, every solution of the SAA problem is an exact solution of the true problem. The sharpness condition is defined as follows. We say that the minimizer set of Ef over X is *sharp* if there exists $c > 0$ such that

$$Ef(y) \geq \inf_{x \in X} Ef(x) + c \text{dist}(y, \text{argmin}_{x \in X} Ef(x)), \quad \forall y \in X;$$

see [49, Equation 5.138] and [46, Assumption (A)]. This condition is closely related to the notion of weak sharp minima in [12].

An essential ingredient in the analyses of [46, 32, 48] is a uniform SLLN for convex subdifferentials under the stated assumptions; see, e.g., [46, Lemma 2.4]. Our Lipschitzian SLLNs yield the following analogous finite-sample identification result for a substantially broader class of problems, without requiring polyhedrality, finite support of the distribution, or convexity of the objective.

Proposition 5.5 (global solution). *Suppose that the assumptions of one of Theorems 3.3, 3.8 or 4.10 hold. If the minimizer set of Ef over X is nonempty and sharp, then, for all sufficiently large $\nu \in \mathbb{N}$,*

$$\text{argmin}_{x \in X} E^\nu f(x) \subset \text{argmin}_{x \in X} Ef(x), \quad \mathbb{P}\text{-a.s.}$$

Proof. Let $g^\nu = E^\nu f - Ef$ and $A = \text{argmin}_{x \in X} Ef(x)$. For any $y \in X$ and $x \in A$, one has

$$E^\nu f(y) - E^\nu f(x) = Ef(y) - Ef(x) + g^\nu(y) - g^\nu(x) \geq c \text{dist}(y, A) - \text{g-lip}_X(g^\nu) \|y - x\|_2.$$

By $\inf_{z \in X} E^\nu f(z) \leq E^\nu f(x)$ and then taking the supremum over $x \in A$ gives

$$E^\nu f(y) - \inf_{z \in X} E^\nu f(z) \geq (c - \text{g-lip}_X(g^\nu)) \text{dist}(y, A).$$

By Theorems 3.3, 3.8 or 4.10, $\text{g-lip}_X(g^\nu) \rightarrow 0$ $\mathbb{P}$ -a.s., so eventually $c > \text{g-lip}_X(g^\nu)$. Now let $y \in \text{argmin}_{x \in X} E^\nu f(x)$. Then the left-hand side above is zero, and therefore $\text{dist}(y, A) = 0$. Since Ef is continuous on X , the set A is closed relative to X . Thus $y \in A$. $\square$

Unlike the local consequences established in Section 5.2, the finite-sample identification result in Proposition 5.5 relies essentially on convergence of the global Lipschitz modulus $\text{g-lip}_X(E^\nu f - Ef)$. For nonconvex problems, however, obtaining global solutions to the SAA problem can be difficult, motivating an analogous result for stationary points. The uniform convergence of subdifferentials in Corollary 5.3 yields such an identification result under a sharpness-type condition.

Proposition 5.6 (stationary points). *Suppose that the assumptions of one of Theorems 3.3, 3.8 or 4.10 hold and that, for $S \subset U \subset \text{int } X$, one has*

$$\inf_{x \in U \setminus S} \text{dist}(0, \partial Ef(x)) \geq \eta \quad (6)$$

for some tolerance $\eta > 0$. Then, for any $\varepsilon^\nu \downarrow 0$ and all sufficiently large $\nu \in \mathbb{N}$, $\mathbb{P}$ -a.s., one has

$$S^\nu = \{x \in U \mid \text{dist}(0, \partial E^\nu f(x)) \leq \varepsilon^\nu\} \subset S.$$

Proof. By Corollary 5.3, $\mathbb{P}$ -a.s., $\rho^\nu = \sup_{x \in U} d(\partial E^\nu f(x), \partial Ef(x)) \rightarrow 0$. For every $y \in S^\nu$,

$$\text{dist}(0, \partial Ef(y)) \leq \text{dist}(0, \partial E^\nu f(y)) + \rho^\nu \leq \varepsilon^\nu + \rho^\nu.$$

For all sufficiently large ν , $\varepsilon^\nu + \rho^\nu < \eta$. Hence $y \notin U \setminus S$, and therefore $y \in S$. $\square$

5.4 Permanence properties

We record several operations under which convergence in the Lipschitz pseudometric is preserved. Linear operations, restrictions, and Lipschitz changes of variables require no additional regularity. Smooth outer compositions are also admissible on bounded sets. These permanence properties provide calculus rules that carry the above results to broader classes of problems.

Proposition 5.7 (permanence). *Let $0 \in X \subset \mathbb{R}^d$, and suppose that, for every $i \in [m]$, the functions $h_i^\nu, h_i \in \text{Lip}(X)$ satisfy $d_X(h_i^\nu, h_i) \rightarrow 0$, as $\nu \rightarrow \infty$. Then the following statements hold.*

- (a) *For any $a_1, \dots, a_m \in \mathbb{R}$, one has $d_X(\sum_{i=1}^m a_i h_i^\nu, \sum_{i=1}^m a_i h_i) \rightarrow 0$, as $\nu \rightarrow \infty$.*
- (b) *If $0 \in Z \subset X$, then $d_Z(h_i^\nu, h_i) \rightarrow 0$ as $\nu \rightarrow \infty$, for every $i \in [m]$.*
- (c) *Let $0 \in Y \subset \mathbb{R}^p$, and let $T : Y \rightarrow X$ be Lipschitz with $T(0) = 0$. Then $d_Y(h_i^\nu \circ T, h_i \circ T) \rightarrow 0$ for every $i \in [m]$.*
- (d) *Suppose, in addition, that X is bounded. Define $H^\nu = (h_1^\nu, \dots, h_m^\nu)$ and $H = (h_1, \dots, h_m)$. Then, for every C^1 function $G : \mathbb{R}^m \rightarrow \mathbb{R}$, one has $d_X(G \circ H^\nu, G \circ H) \rightarrow 0$ as $\nu \rightarrow \infty$.*

Proof. Part (a) follows from the triangle inequality and homogeneity of $\|\cdot\|_{\text{lip}}$. Part (b) is a special case of (c). For part (c), let $e^\nu = h_i^\nu - h_i$ and L_T be a Lipschitz constant of map T on the set Y . Since $T(0) = 0$, one has $|e^\nu(T(0))| = |e^\nu(0)|$ and $\text{g-lip}_Y(e^\nu \circ T) \leq \text{g-lip}_X(e^\nu)L_T$. Hence $d_Y(h_i^\nu \circ T, h_i \circ T) \leq \max\{1, L_T\}d_X(h_i^\nu, h_i) \rightarrow 0$. For part (d), since X is bounded, Proposition 5.1 gives $\delta_i^\nu = \sup_{x \in X} |h_i^\nu(x) - h_i(x)| \rightarrow 0$ for every $i \in [m]$, while $L_i^\nu = \text{g-lip}_X(h_i^\nu - h_i) \rightarrow 0$. Let $\delta^\nu = (\sum_{i=1}^m (\delta_i^\nu)^2)^{1/2}$, $L^\nu = (\sum_{i=1}^m (L_i^\nu)^2)^{1/2}$, and $L = (\sum_{i=1}^m \text{g-lip}_X(h_i)^2)^{1/2}$. Then $\delta^\nu \rightarrow 0$ and $L^\nu \rightarrow 0$. The ranges of H^ν and H are eventually contained in a common compact convex set K . Let $M = \sup_{z \in K} \|\nabla G(z)\|_2$, and let β be a modulus of continuity of ∇G on K . By the fundamental theorem of calculus, for all $x, y \in X$,

$$|G(H^\nu(x)) - G(H(x)) - G(H^\nu(y)) + G(H(y))| \leq (ML^\nu + \beta(\delta^\nu)L)\|x - y\|_2.$$

Moreover, $|G(H^\nu(0)) - G(H(0))| \leq M\delta^\nu$. Hence $d_X(G \circ H^\nu, G \circ H) \rightarrow 0$. $\square$

For illustration, let $h^\nu, h, g^\nu, g \in \text{Lip}(\mathbb{R}^d)$ satisfying $d_{\mathbb{R}^d}(h^\nu, h) \rightarrow 0$ and $d_{\mathbb{R}^d}(g^\nu, g) \rightarrow 0$. By Proposition 5.2 and the stability of the Hausdorff distance under Minkowski addition,

$$\sup_{x \in \mathbb{R}^d} d(\partial h^\nu(x) + \partial g^\nu(x), \partial h(x) + \partial g(x)) \rightarrow 0.$$

This conclusion is weaker than a direct comparison of the subdifferentials of the sums, since, in general, $\partial(h + g)(x) \subset \partial h(x) + \partial g(x)$. By Proposition 5.7, however, $d_{\mathbb{R}^d}(h^\nu + g^\nu, h + g) \rightarrow 0$, and hence Proposition 5.2 yields

$$\sup_{x \in \mathbb{R}^d} d(\partial(h^\nu + g^\nu)(x), \partial(h + g)(x)) \rightarrow 0.$$

Thus, the Lipschitz pseudometric provides a convenient surrogate for controlling Hausdorff distances between subdifferentials, owing to its favorable calculus properties.

So far, we have considered only the unconstrained case. Let $\mathcal{N}_X(x)$ be the limiting normal cone to a set X at x ; see [42, Definition 6.3]. We next incorporate the constraint set in (1), as illustrated by the following result.

Proposition 5.8 (constrained problem). *Let $Y \subset \mathbb{R}^d$, with $0 \in Y$, be an open superset of a nonempty closed set X . For $h^\nu, h : \mathbb{R}^d \rightarrow \mathbb{R}$, locally Lipschitz on Y , one has*

$$\begin{aligned} \sup_{x \in X} d(\partial h^\nu(x) + \mathcal{N}_X(x), \partial h(x) + \mathcal{N}_X(x)) &\leq d_Y(h^\nu, h), \\ \sup_{x \in X} d(\partial(h^\nu + \iota_X)(x), \partial(h + \iota_X)(x)) &\leq d_Y(h^\nu, h). \end{aligned}$$

Proof. Since the claim is trivial when $d_Y(h^\nu, h) = \infty$, suppose that it is finite. Then $\text{g-lip}_Y(h^\nu - h) \leq d_Y(h^\nu, h)$. By [42, Corollary 10.9] and [42, Theorem 9.13], for every $x \in X$, $\partial h^\nu(x) \subset \partial h(x) + \text{g-lip}_Y(h^\nu - h)\mathbb{B}$. Interchanging h^ν and h gives $\partial h(x) \subset \partial h^\nu(x) + \text{g-lip}_Y(h^\nu - h)\mathbb{B}$. Adding $\mathcal{N}_X(x)$ to both inclusions and using [42, Example 4.13, equation (4)(5)] yields the first inequality.

Now let $H^\nu = h^\nu + \iota_X$ and $H = h + \iota_X$. Since X is nonempty and closed and h^ν, h are continuous on Y , the functions H^ν and H are proper and lower semicontinuous on a neighborhood of any $x \in X$. Applying the same argument as in the first claim to $H^\nu = H + (h^\nu - h)$ and $H = H^\nu + (h - h^\nu)$ yields the second inequality. $\square$

Therefore, our previous consistency results extend straightforwardly to the constrained setting. We omit the formal statement for brevity.

References

- [1] C. D. Aliprantis and K. C. Border. *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer, 3rd edition, 2006.
- [2] F. Areces, J. Duchi, and M. Sommers. Finding a stationary point of a stochastic convex problem. *arXiv preprint arXiv:2607.06883*, 2026.
- [3] Z. Artstein and R. J-B Wets. Consistency of minimizers and the SLLN for stochastic programs. *Journal of Convex Analysis*, 2(1-2):1–17, 1995.
- [4] R. J. Aumann. Integrals of set-valued functions. *Journal of Mathematical Analysis and Applications*, 12(1):1–12, 1965.

- [5] G. Bareilles, A. Gehret, J. Aspman, J. Lepšová, and J. Mareček. Deep learning as the disciplined construction of tame objects. *arXiv preprint arXiv:2509.18025*, 2025.
- [6] G. Beer. *Bornologies and Lipschitz Analysis*. CRC Press, 2023.
- [7] G. Beer and M. J. Hoffman. The Lipschitz metric for real-valued continuous functions. *Journal of Mathematical Analysis and Applications*, 406(1):229–236, 2013.
- [8] D. Berend and A. Kontorovich. The missing mass problem. *Statistics & Probability Letters*, 82(6):1102–1110, 2012.
- [9] J. Bolte and E. Pauwels. Conservative set valued fields, automatic differentiation, stochastic gradient methods and deep learning. *Mathematical Programming*, 188(1):19–51, 2021.
- [10] J. Bolte, A. Daniilidis, A. Lewis, and M. Shiota. Clarke subgradients of stratifiable functions. *SIAM Journal on Optimization*, 18(2):556–572, 2007.
- [11] J. Bolte, T. Le, and E. Pauwels. Subgradient sampling for nonsmooth nonconvex minimization. *SIAM Journal on Optimization*, 33(4):2542–2569, 2023.
- [12] J. V. Burke and M. C. Ferris. Weak sharp minima in mathematical programming. *SIAM Journal on Control and Optimization*, 31(5):1340–1359, 1993.
- [13] J. V. Burke, X. Chen, and H. Sun. The subdifferential of measurable composite max integrands and smoothing approximation. *Mathematical Programming*, 181(2):229–264, 2020.
- [14] C. C. Chang and H. J. Keisler. *Model Theory*, volume 73. Elsevier, 3rd edition, 1990.
- [15] H. Chase and J. Freitag. Model theory and machine learning. *Bulletin of Symbolic Logic*, 25(3):319–332, 2019.
- [16] F. H. Clarke. *Optimization and Nonsmooth Analysis*. SIAM, 1990.
- [17] D. L. Cohn. *Measure Theory*. Birkhäuser Advanced Texts Basler Lehrbücher. Springer, 2nd edition, 2013.
- [18] M. Coste. *An Introduction to O-minimal Geometry*. RAAG Notes. Institut de Recherche Mathématiques de Rennes, 1999.
- [19] M. Coste. *An Introduction to Semialgebraic Geometry*. RAAG Notes. Institut de Recherche Mathématiques de Rennes, 2002.
- [20] D. Davis and D. Drusvyatskiy. Graphical convergence of subgradients in nonconvex optimization and learning. *Mathematics of Operations Research*, 47(1):209–231, 2022.
- [21] D. Davis, D. Drusvyatskiy, S. Kakade, and J. D. Lee. Stochastic subgradient method converges on tame functions. *Foundations of Computational Mathematics*, 20(1):119–154, 2020.
- [22] T. Devale, P. Devulapalli, and S. Hanneke. Uniform convergence beyond Glivenko-Cantelli. In Matus Telgarsky and Jonathan Ullman, editors, *International Conference on Algorithmic Learning Theory*, volume 313 of *Proceedings of Machine Learning Research*, pages 1–21. PMLR, 23–26 Feb 2026.

- [23] G. B. Folland. *Real Analysis: Modern Techniques and Their Applications*. John Wiley & Sons, 2nd edition, 1999.
- [24] D. J. Foster, A. Sekhari, and K. Sridharan. Uniform convergence of gradients for non-convex learning and optimization. *Advances in Neural Information Processing Systems*, 31:8759–8770, 2018.
- [25] R. M. Gray and D. L. Neuhoff. Quantization. *IEEE Transactions on Information Theory*, 44(6):2325–2383, 2002.
- [26] A. Günaydin and P. Hieronymi. Dependent pairs. *The Journal of Symbolic Logic*, 76(2):377–390, 2011.
- [27] P. Hieronymi. Defining the set of integers in expansions of the real field by a closed discrete set. *Proceedings of the American Mathematical Society*, 138(6):2163–2168, 2010.
- [28] A. D. Ioffe. An invitation to tame optimization. *SIAM Journal on Optimization*, 19(4):1894–1917, 2009.
- [29] A. Iusem and A. Seeger. Distances between closed convex cones: Old and new results. *Journal of Convex Analysis*, 17(3-4):1033–1055, 2010.
- [30] T. Kaiser. First order tameness of measures. *Annals of Pure and Applied Logic*, 163(12):1903–1927, 2012.
- [31] M. Karpinski and A. Macintyre. Polynomial bounds for VC dimension of sigmoidal and general Pfaffian neural networks. *Journal of Computer and System Sciences*, 54(1):169–176, 1997.
- [32] A. J. Kleywegt, A. Shapiro, and T. Homem-de-Mello. The sample average approximation method for stochastic discrete optimization. *SIAM Journal on Optimization*, 12(2):479–502, 2002.
- [33] L. S. Krapp, M. Vermeil, and L. Wirth. On tameness, measurability and the independence property. *arXiv preprint arXiv:2506.08733*, 2025.
- [34] M. C. Laskowski. Vapnik-Chervonenkis classes of definable sets. *Journal of the London Mathematical Society*, 2(2):377–384, 1992.
- [35] M. Ledoux and M. Talagrand. *Probability in Banach Spaces: Isoperimetry and Processes*, volume 23. Springer, 1991.
- [36] R. Livni and P. Simon. Honest compressions and their application to compression schemes. In *Conference on Learning Theory*, pages 77–92. PMLR, 2013.
- [37] A. Macintyre and E. D. Sontag. Finiteness results for sigmoidal “neural” networks. In *ACM Symposium on Theory of Computing*, pages 325–334, 1993.
- [38] S. Mei, Y. Bai, and A. Montanari. The landscape of empirical risk for nonconvex losses. *Annals of Statistics*, 46(6A):2747–2774, 2018.
- [39] H. D. Nguyen, J. Westerhout, and X. Guo. On rates of convergence for sample average approximations without smoothness. *arXiv preprint arXiv:2604.25153*, 2026.

- [40] V. I. Norkin and R. J-B Wets. On strong graphical law of large numbers for random semicontinuous mappings. *Vestnik Sankt-Peterburgskogo Universiteta*, Seriya 10(3):102–111, 2013.
- [41] R. T. Rockafellar. Favorable classes of Lipschitz-continuous functions in subgradient optimization. In E. Nurminski, editor, *Progress in Nondifferentiable Optimization*, IIASA Collaborative Proceedings Series, pages 125–144. International Institute for Applied Systems Analysis, Laxenburg, Austria, 1982.
- [42] R. T. Rockafellar and R. J-B Wets. *Variational Analysis*. Springer, 3rd printing-2009 edition, 1998.
- [43] F. Ruan. On the uniform convergence of subdifferentials in stochastic optimization and learning. *Mathematics of Operations Research*, to appear, 2025.
- [44] S. Schechtman. The gradient’s limit of a definable family of functions admits a variational stratification. *SIAM Journal on Optimization*, 36(2):1075–1099, 2026.
- [45] S. Shalev-Shwartz and S. Ben-David. *Understanding Machine Learning: From Theory to Algorithms*. Cambridge University Press, 2014.
- [46] A. Shapiro and T. Homem-de-Mello. On the rate of convergence of optimal solutions of Monte Carlo approximations of stochastic programs. *SIAM Journal on Optimization*, 11(1):70–86, 2000.
- [47] A. Shapiro and H. Xu. Uniform laws of large numbers for set-valued mappings and subdifferentials of random functions. *Journal of Mathematical Analysis and Applications*, 325(2): 1390–1399, 2007.
- [48] A. Shapiro, T. Homem-de Mello, and J. Kim. Conditioning of convex piecewise linear stochastic programs. *Mathematical Programming*, 94(1):1–19, 2002.
- [49] A. Shapiro, D. Dentcheva, and A. Ruszczyński. *Lectures on Stochastic Programming: Modeling and Theory*. SIAM, 3rd edition, 2021.
- [50] P. Simon. *A Guide to NIP Theories*. Lecture Notes in Logic. Cambridge University Press, 2015.
- [51] C. I. Steinhorn. Chapter XVI: Borel structures and measure and category logics. In *Model-Theoretic Logics*, volume 8, pages 579–597. Association for Symbolic Logic, 1985.
- [52] P. Terán. On a uniform law of large numbers for random sets and subdifferentials of random functions. *Statistics & Probability Letters*, 78(1):42–49, 2008.
- [53] L. Tian and J. O. Royset. Failure of uniform laws of large numbers for subdifferentials and beyond. *arXiv preprint arXiv:2511.16568*, 2025.
- [54] A. W. van der Vaart and J. A. Wellner. *Weak Convergence and Empirical Processes*. Springer, 2nd edition, 2023.
- [55] L. van den Dries. *Tame Topology and O-minimal Structures*, volume 248. Cambridge University Press, 1998.

- [56] L. van den Dries and C. Miller. Geometric categories and o-minimal structures. *Duke Mathematical Journal*, 84(2), 1996.
- [57] A. W. van der Vaart. New Donsker classes. *Annals of Probability*, 24(4):2128–2140, 1996.
- [58] A. W. van der Vaart and J. A. Wellner. Preservation theorems for Glivenko-Cantelli and uniform Glivenko-Cantelli classes. In *High Dimensional Probability II*, pages 115–133. Springer, 2000.
- [59] A. J. Wilkie. Model completeness results for expansions of the ordered field of real numbers by restricted Pfaffian functions and the exponential function. *Journal of the American Mathematical Society*, 9(4):1051–1094, 1996.
- [60] H. Xu. Uniform exponential convergence of sample average random functions under general sampling with applications in stochastic programming. *Journal of Mathematical Analysis and Applications*, 368(2):692–710, 2010.