

Leveraging tails for adaptation

SERGIOS AGAPIOU¹ ISMAËL CASTILLO² PAUL EGELS³

¹*Department of Mathematics and Statistics, University of Cyprus, Nicosia, Cyprus.*

E-mail: agapiou.sergios@ucy.ac.cy

²*Sorbonne Université, LPSM; 4, place Jussieu, 75005 Paris, France.*

E-mail: ismael.castillo@sorbonne-universite.fr

³*Sorbonne Université, LPSM; 4, place Jussieu, 75005 Paris, France.*

E-mail: paul.egels@sorbonne-universite.fr

We consider contraction of Bayesian posterior distributions in nonparametric settings where coefficients of a function over a basis or dictionary are given priors with p -exponential tails, including Laplace tails ($p = 1$) and heavier tails ($p < 1$). It is shown that contraction rates improve as p decreases and that full adaptation to smoothness, up to logarithmic factors, is obtained in an appropriate $p \rightarrow 0$ regime. As applications, we consider both series priors in white noise regression and shallow ReLU neural networks in random design regression. In particular, we show that overparametrised shallow ReLU networks can adapt to any regularity $0 \leq \beta \leq 2$. Through a simulation study, we show strong empirical agreement with the behavior predicted by our theory.

Keywords: Frequentist analysis of Bayesian procedures, adaptation, nonparametric regression, Bayesian neural networks, overparametrization.

1. Introduction

A central goal in nonparametric statistics is *adaptation*: the ability of an estimator to perform simultaneously and optimally across a wide variety of settings with little to no tuning. When inference is carried out over a class of functional spaces, it is desirable that the estimator automatically adapts to unknown features of these spaces, such as smoothness, geometry, sparsity or other finer structural properties. A large body of literature has focused on adaptation: Lepski’s method Lepski [1990, 1991], thresholding Donoho et al. [1995] and model selection Barron et al. [1999] are amongst the most well-known non-Bayesian approaches. Bayesian methods, on the other hand, have a natural ability to achieve adaptation, as we discuss in more detail below, by choosing prior distributions that are flexible enough to achieve this task (one possibility is for instance to draw certain prior parameters at random in a hierarchical Bayes fashion).

Recently, motivated by the remarkable empirical success of deep learning methods, there has been a growing interest in understanding how neural networks can automatically learn structural parameters, such as smoothness of functions or ‘effective’ dimensions, for instance in regression settings exhibiting a compositional structure as in Schmidt-Hieber [2020], Kohler and Langer [2021] or for data lying on geometric structures (e.g. Nakada and Imaizumi [2020]). While the above works prove that optimal convergence rates can indeed be achieved by *appropriately* choosing the network’s architecture, this choice still typically depends on parameters unknown to the statistician. In principle a solution to this *adaptation* problem could be to compare empirically risks for different architectures (by a method such as model selection or cross-validation); this would be particularly demanding computationally, and in practice overparameterized networks are used. These have a width or a depth larger than what traditional bias–variance trade-offs would suggest as optimal ones, but still empirically perform remarkably well. It is therefore particularly desirable to understand statistical properties of overparam-

terized networks, in particular, whether they can achieve adaptation to smoothness or to other structural parameters.

In this work, we take a step in this direction by showing, in a simple regression setting, that overparameterized prior distributions, with similar tails to those arising as outputs of typical neural networks with random Gaussian weights, have natural adaptation properties. Our results are relevant both for adaptation properties of classical projection estimators in regression as well as for neural network-type estimates—here for simplicity we consider shallow networks—. We now briefly review adaptation methods from a Bayesian perspective.

In the Bayesian setting, Gaussian processes (henceforth GPs) are among the most celebrated and widely used priors in machine learning and statistical applications (see e.g. [Rasmussen and Williams \[2006\]](#), [Liu et al. \[2020\]](#)). Thanks to pioneering work by [van der Vaart and van Zanten \[2008\]](#) building on the general contraction rate theory of [Ghosal et al. \[2000\]](#), contraction rates of posterior distributions for GPs are now fairly well understood from a theoretical perspective. While *per se* Gaussian processes are not adaptive to smoothness (see [Castillo \[2008\]](#)), a large body of literature has shown that GPs can be made adaptive to (homogeneous) smoothness [van der Vaart and van Zanten \[2009\]](#), [Szabó et al. \[2013\]](#), [Castillo et al. \[2014\]](#), [Knapik et al. \[2016\]](#), [Rousseau and Szabo \[2017\]](#) as well as to geometry in manifold settings [Berenfeld et al. \[2024\]](#), [Yang and Dunson \[2016\]](#), [Tang et al. \[2026\]](#), provided they are properly rescaled (depending on the observed data) or their parameters are suitably estimated or drawn from a hyperprior.

However, several drawbacks are associated with Gaussian process approaches. First, achieving adaptation requires an additional computational layer in order to adjust scale or regularity parameters, which can render GP regression computationally demanding, especially in high-dimensional and data-rich regimes, see for example [Agapiou et al. \[2014\]](#). A number of works aim to mitigate this complexity by developing scalable approximations of GPs, for instance through sparse variational inference [Nieman and Szabó \[2025\]](#) or local approximations (see e.g. [Szabó and Zhu \[2026\]](#) for further references). Second, and more fundamentally, recent negative results show that even properly rescaled GPs may fail to adapt when finer notions of adaptivity are required, such as adaptation to compositional structures [Giordano et al. \[2022\]](#) (see also [Abraham and Deo \[2023\]](#) in the context of inverse problems) or inhomogeneous smoothness [Agapiou and Wang \[2024\]](#). In order to obtain stronger forms of adaptation, a number of works advocate and study deep Bayesian methods, such as Bayesian deep GPs [Damianou and Lawrence \[2013\]](#), [Finocchio and Schmidt-Hieber \[2023\]](#), [Castillo and Randrianarisoa \[2025\]](#) and Bayesian deep (Gaussian) neural networks [Kong and Kim \[2025\]](#). These approaches exhibit stronger adaptive properties compared to standard GPs, notably with respect to compositional structures. Nevertheless, with the currently available theory, they still require hyperparameter (or hyperprior) estimation. Contributions on posterior rates for Bayesian deep networks include [Polson and Ročková \[2018\]](#) for spike-and-slab priors on weights (see also [Chérif-Abdellatif \[2020\]](#), [Bai et al. \[2020\]](#), [Ohn and Lin \[2024\]](#) for variational Bayes counterparts) and [Lee and Lee \[2022\]](#), [Kong et al. \[2023\]](#), [Kong and Kim \[2025\]](#); a review on Bayesian deep neural networks can be found in [Arbel et al. \[2026\]](#).

It turns out that replacing the Gaussian distribution with heavier-tailed distributions can lead to improved contraction rates. One example is the Laplace–Besov prior, which replaces Gaussian coefficients by double–exponential ones and has recently been studied in a variety of nonparametric settings, including regression, density estimation, and inverse problems [Agapiou and Wang \[2024\]](#), [Giordano \[2023\]](#), [Dolera et al. \[2024\]](#). These works build on a more general theory for p -exponential priors, corresponding to densities proportional to $\exp(-|x|^p/p)$ and whose tails interpolate between Laplace ($p = 1$) and Gaussian ($p = 2$), which was initiated in [Agapiou et al. \[2021\]](#) and later complemented by [Agapiou and Savva \[2024\]](#). Interestingly, although the contraction rates obtained in [Agapiou et al. \[2021\]](#) for such p -exponential priors ($1 \leq p < 2$) are not minimax optimal, for $1 \leq p < 2$ they are still polynomially faster than the corresponding Gaussian rates ($p = 2$). Moreover, when scaling and/or

regularity hyperparameters are drawn at random, these works establish adaptation properties of p -exponential priors ($1 \leq p \leq 2$) with respect to inhomogeneous smoothness in an L_2 -sense (see also [Agapiou et al. \[2026\]](#) for further discussion on adaptation to inhomogeneous smoothness). However, such p -exponential priors still require some form of hyperparameter estimation in order to achieve adaptation, and proof techniques crucially rely on log-concavity of the prior density, which only holds for $p \geq 1$.

Very recently, [Agapiou and Castillo \[2024\]](#) initiated the theoretical study of even heavier-tailed priors, for instance with polynomially decaying tails such as the Student- t distribution. One main advantage of such priors is that they only require a fixed universal deterministic (in particular, data-independent) rescaling, so that no hyperprior or hyperparameter estimation is needed to achieve adaptation. In [Agapiou et al. \[2026\]](#), these heavier-tailed priors were shown to possess strong adaptive properties with respect to non-homogeneously smooth Besov functions, this time in any L_r -norm ($r \geq 1$). When applied to overparameterized deep Bayesian neural networks, [Castillo and Egels \[2025\]](#) showed that they yield posteriors that adapt to compositional structures and manifold geometry, again without requiring hyperparameter estimation.

The primary goal of the present paper is to understand adaptation properties of p -exponential type priors over the full range of tail indices $0 < p \leq 2$, and in particular in the (formally defined below) limiting regime $p \rightarrow 0$. There are two main motivations for investigating the “small p regime” for p -exponential priors. The first is that the existing results on contraction rates for p -exponential priors indicate that the rate may (further) improve as p decreases below 1, suggesting that adaptation may be obtained for such priors by formally letting $p \rightarrow 0$ *without* the need to sample hyperparameters. The second motivation comes from the study of deep learning methods, in particular Bayesian neural networks with Gaussian priors on network weights. Indeed, several recent works show that, conditionally on the input, the output of deep neural networks with i.i.d. Gaussian weights exhibits heavy-tailed behavior, [Vladimirova et al. \[2019\]](#), [Zavatone-Veth and Pehlevan \[2021\]](#), [Noci et al. \[2021\]](#). More precisely, such outputs are of (generalized) sub-Weibull type with index $L/2$ (see e.g. [Vladimirova et al. \[2020\]](#)), which is closely related to the p -exponential distribution with $p = 2/L$, where L denotes the depth of the network. These results suggest that, for deep neural networks with i.i.d. Gaussian weights, increasing depth naturally induces marginally p -exponential outputs, in the regime where p approaches 0 inversely proportionally to the network depth.

Our study, on the one hand, complements the work of [Agapiou et al. \[2021\]](#), and builds a bridge between the very heavy-tailed (polynomial) case of [Agapiou and Castillo \[2024\]](#), where posteriors exhibit powerful adaptive properties, and the lighter heavy-tailed priors such as the Laplace-Besov case, which are more commonly used in practice. On the other hand, this will enable us to analyse *overparameterized* shallow ReLU neural networks with p -exponential weights, and show that the latter satisfy a remarkable adaptation to smoothness property.

We now summarize our main contributions:

1. In a white-noise regression setting, we study series priors with independent p -exponentially distributed coefficients ($p > 0$) and derive an upper bound on the contraction rate of ρ -posteriors ($0 < \rho < 1$) toward the unknown regression function. This bound considerably broadens previously known results of [Agapiou et al. \[2021\]](#) for $1 \leq p \leq 2$. Notably, the obtained rate becomes near-optimal as the tail parameter p approaches 0, that is, as the prior becomes heavier-tailed. Also, previous works for $p \in [1, 2]$ left open the question whether the obtained rates could be improved; here we provide a matching lower bound, valid also for standard posteriors ($\rho = 1$).
2. In random design nonparametric regression, we study a class of *overparameterized* ReLU shallow neural network priors, where the number of neurons is taken explicitly much larger than the oracle number that achieves the optimal minimax rate. An independent p -exponential prior

- ($0 < p \leq 1$) is placed on the weights of the hidden layer, and an upper bound on the contraction rate of the corresponding ρ -tempered posterior is obtained. Although different in nature from the series prior setting, these neural network rates share a similar type of improvement as p approaches 0, with a rate that is near-minimax for small p ;
3. In both of the above settings, we formally show that taking the limit $p \rightarrow 0$ as the sample size n goes to infinity, leads to fully adaptive posteriors over smoothness classes. These results can be linked to earlier findings of [Agapiou and Castillo \[2024\]](#) for series priors and [Castillo and Egels \[2025\]](#) for neural networks. In particular, in the latter case, our results provide further insight into the adaptivity of deep Gaussian Bayesian neural networks (here small p may be interpreted as mimicking the effect of increasing depth in a network; see also the discussion in Section 5).
 4. We provide numerical experiments confirming that in both frameworks adaptation automatically occurs for the resulting estimators as the tail-index p approaches 0. In particular, ReLU shallow networks with overparametrised width of order n and weights with p -exponential distributions and small deterministic scalings indeed achieve adaptation to smoothness in practical experiments, closely matching the predicted behaviour from our theory.

Frequentist analysis of posterior distributions. We consider a family of probability distributions $(P_f^{(n)})$ parametrized by $f \in \mathcal{F}$, where $\mathcal{F}$ is a (possibly infinite-dimensional) parameter space and $n \geq 1$ is an integer representing the informativeness of the observed data. Given data $X^{(n)} \sim P_{f_0}^{(n)}$ (sometimes simply denoted X) generated from a ‘true’ $f_0 \in \mathcal{F}$, we estimate f_0 from $X^{(n)}$ using a Bayesian procedure. Starting from a prior distribution Π on $\mathcal{F}$ and a parameter $\rho \in (0, 1]$, the ρ -tempered (or fractional) posterior distribution $\Pi_\rho[\cdot | X]$ by, for any measurable set B ([Ghosal and van der Vaart \[2017\]](#), [Castillo \[2024\]](#)),

$$\Pi_\rho[B | X] = \frac{\int_B \exp(\rho \ell_n(f, X)) d\Pi(f)}{\int \exp(\rho \ell_n(f, X)) d\Pi(f)}, \quad (1)$$

where $\ell_n(f, X) := \log p_f^{(n)}(X)$ denotes the log-likelihood. For $\rho = 1$, the fractional posterior distribution coincides with the usual posterior $\Pi[\cdot | X]$ given by Bayes’ formula. In this paper, for technical simplicity we focus on ρ -tempered posteriors with fixed $\rho < 1$, although we expect most results to carry over to the case $\rho = 1$ (see also Section 5 for more details).

Contraction rates of ρ -posteriors. For any positive loss function d on $\mathcal{F} \times \mathcal{F}$, we say that the ρ -posterior contracts around f_0 at the rate $\varepsilon_n \rightarrow 0$ in d -loss if

$$E_{f_0} \Pi_\rho(d(f, f_0) \leq M \varepsilon_n | X) \rightarrow 1, \quad \text{as } n \rightarrow \infty, \quad (2)$$

with $M > 0$ a sufficiently large constant and E_{f_0} the expectation under $P_{f_0}^{(n)}$.

p -exponential distributions. The prior distributions Π we consider below are based on the p -exponential distribution on $\mathbb{R}$: for $p > 0$, this distribution has density given by, for $t \in \mathbb{R}$,

$$h_p(t) \propto \exp \left\{ -\frac{|t|^p}{p} \right\}. \quad (3)$$

Such distribution generalizes the Gaussian ($p = 2$) and Laplace ($p = 1$) distributions, with possibly heavier tails as the parameter p gets closer to 0. More generally, for any $p > 0$, we say that a density

function h has p -tails (or p -exponential tails) if

$$h \text{ is symmetric about } 0 \quad (h(-t) = h(t) \text{ for all } t), \quad (4)$$

if for some constants $c_0, c_1 > 0$, one has

$$h(t) \geq c_0 e^{-c_1 t^p} \quad t \geq 0, \quad (5)$$

and, denoting by $\bar{H}(x) := \int_x^{+\infty} h(t) dt$ the survival function associated with h , if for $q \in (0, p]$ and positive constants d_0, d_1, M_0 ,

$$\bar{H}(x) \leq d_0 e^{-d_1 x^q} \quad x \geq M_0. \quad (6)$$

For example, the p -exponential density h_p defined in (3) satisfies conditions (4)–(5)–(6) for any $q < p$ (see e.g. Lemma 9 for the case $p < 1$). Other commonly used prior distributions such as Weibull and Generalized-Weibull (see e.g. Vladimirova et al. [2020]) distributions also have p -tails in the above sense.

Outline. In Section 2, we state our results for p -exponential series priors in white noise regression, both for given p and in a regime $p \rightarrow 0$, together with a corresponding matching lower bound. Section 3 focuses on results for overparameterized shallow ReLU neural network posteriors. A simulation study illustrating both settings can be found in Section 4, while the discussion in Section 5 puts our results in perspective. Part of the proofs of the main results can be found in Section 6. The Supplementary material contains the remaining proofs, a number of additional results as well as some technical Lemmata.

2. Series priors

In this section, for technical simplicity we focus on the prototypical nonparametric Gaussian white noise model: for $f \in L^2[0, 1]$ set of squared-integrable functions on $[0, 1]$, one observes

$$dZ^{(n)}(t) = f(t)dt + dW(t)/\sqrt{n}, \quad t \in [0, 1],$$

where W is standard Brownian motion and $n \geq 1$. For $(\varphi_k)_{k \geq 1}$ an orthonormal basis of $L^2([0, 1])$, for the canonical inner product, one denotes $f_k := \langle f | \varphi_k \rangle$ its basis coefficients. The white noise model above induces, once projected into the basis (φ_k) , observations in the so-called Gaussian sequence model (see the book by Giné and Nickl [2015])

$$X_k := f_k + \frac{1}{\sqrt{n}} \xi_k, \quad k \geq 1, \quad (7)$$

where ξ_k are i.i.d. $\mathcal{N}(0, 1)$ random variables. The observation sequence from the model (7) will be denoted $X := X^{(n)} \sim P_f^{(n)}$ and associated with the log-likelihood

$$\ell_n(f, X) := -\frac{n}{2} \sum_{k \geq 1} (X_k - f_k)^2. \quad (8)$$

Definition of the prior. We define a prior $\Pi := \Pi(p, \alpha)$ on $f \in L^2[0, 1]$, identified as the sequence of its (square summable) coefficients $(f_k)_{k \geq 1}$ by setting

$$f_k = \sigma_k \zeta_k, \quad (9)$$

where ζ_k are i.i.d. random variables with a density h satisfying the p -tails conditions (4)–(5)–(6) for some $p > 0$. A possible choice of scaling sequence $(\sigma_k)_{k \geq 1}$ is, for $\alpha > 0$,

$$\sigma_k = k^{-1/2-\alpha}. \quad (10)$$

Regularity assumption on f_0 , targeted rate. Equipped with the previously defined prior distribution $\Pi = \Pi(p, \alpha)$ and from the likelihood formula (8), one defines (for any $\rho < 1$) the ρ -posterior $\Pi_\rho[\cdot | X]$ using (1). We study these ρ -posteriors under the assumption $X^{(n)} \sim P_{f_0}^{(n)}$, where f_0 belongs to the hyperrectangle, for some $\beta, L > 0$,

$$\mathcal{F}^\beta(L) := \left\{ f = (f_k) : \max_{k \geq 1} \left(|f_k| k^{\beta+1/2} \right) \leq L \right\}. \quad (11)$$

The contraction rate of $\Pi_\rho[\cdot | X]$ around f_0 will involve an interplay between the prior parameters (p, α) and the true function smoothness parameter β . We define

$$\gamma = \gamma(p, \alpha, \beta) := \beta + \frac{p}{2}(\alpha - \beta), \quad (12)$$

and the associated rate

$$\varepsilon_n = \varepsilon_n(p, \alpha, \beta) := \begin{cases} n^{-\frac{\beta}{2\gamma+1}} = n^{-\frac{\beta}{2\beta+p(\alpha-\beta)+1}}, & \alpha > \beta, \\ n^{-\frac{\alpha}{2\alpha+1}}, & \alpha \leq \beta. \end{cases} \quad (13)$$

2.1. Contraction rate for p -tails series priors.

For a prior (9)–(10) with p -tails distributions, the rate (13) turns out to be the L^2 -contraction rate of the associated ρ -posterior. For $g \in L^2[0, 1]$, let $\|g\|_2^2 = \int_0^1 g(u)^2 du$.

Theorem 1 (Upper bound). *Let $p > 0$ and $\alpha, \beta > 0$. Suppose $f_0 \in \mathcal{F}^\beta(L)$ for some $L > 0$, assume $X^{(n)} \sim P_{f_0}^{(n)}$ from the model (7). Then, for any $\rho \in (0, 1)$, starting from the prior $\Pi = \Pi(p, \alpha)$ defined in (9)–(10), as $n \rightarrow \infty$, we have*

$$E_{f_0} \Pi_\rho [\|f - f_0\|_2 \leq M \varepsilon_n | X] \rightarrow 1,$$

where ε_n is given in (13) and $M > 0$ is a large enough constant.

The proof can be found in Section 6. The rate obtained in Theorem 1 depends on the choice of the prior smoothness parameter α compared to the true smoothness β . In particular, there is an elbow in the rate at $\alpha = \beta$. In the undersmoothing case $\alpha < \beta$, the p -tails posterior contracts at the (slower than minimax) rate $n^{-\alpha/(2\alpha+1)}$; in the matching case $\alpha = \beta$, the minimax rate is attained. In the oversmoothing case $\alpha > \beta$, the rate is $n^{-\beta/(2\beta+p(\alpha-\beta)+1)}$ and improves as p gets smaller. In particular, this rate (available for all $p > 0$) matches the one obtained over the range $1 \leq p \leq 2$ in Agapiou et al. [2021]. When $p = 2$, this rate corresponds to the known (sharp) rate of contraction of Gaussian processes obtained in van der Vaart and van Zanten [2008] and Castillo [2008].

Remark 1. Although Theorem 1 assumes f_0 belongs in the hyperrectangle $\mathcal{F}^\beta(L)$ the same result holds true over (the richer) Hilbert–Sobolev Balls: this claim is formally proved for the oversmoothing prior ($\alpha > \beta$) with $p < 1$ in Theorem 6 of the Appendix.

Theorem 2 (Lower bound). Let $p \in (0, 1]$, $L > 0$ and $\alpha > \beta > 0$. Assume the prior on f is defined as in (9)–(10) for the specific choice of density $h = h_p$ as in (3). Then there exists a function $f_0 \in \mathcal{F}^\beta(L)$ such that, if $X^{(n)} \sim P_{f_0}^{(n)}$ from the model (7), then for any $\rho \in (0, 1]$, as $n \rightarrow \infty$,

$$E_{f_0} \Pi_\rho[\|f - f_0\|_2 < m \cdot \varepsilon_n \mid X] \rightarrow 0,$$

where ε_n is given in (13) and $m > 0$ is a small enough constant.

The proof of this result can be found in Appendix B.2. Theorem 2 shows that in the heavier than Laplace case ($p < 1$), the contraction rate ε_n obtained in Theorem 1 is tight, in the sense that there exists some function $f_0 \in \mathcal{F}^\beta(L)$ towards which the p -tails posterior (note that Theorem 2 allows also for the standard posterior $\rho = 1$) cannot contract at a faster rate than ε_n . We believe such lower bound can be obtained also for lighter tails $1 < p < 2$ with similar proof techniques, but here have focused on the (harder) case $p \leq 1$. In the next section, we further explore the rate improvement noted above in a regime $p \rightarrow 0$.

2.2. Adaptation with varying tails.

We showed that whenever p decreases the contraction rate of the p -exponential process posterior improves. In this section, in order to obtain minimax contraction rate and smoothness adaptation, we design priors for which p naturally decreases towards 0.

Let $(\sigma_k)_{k \geq 1}$ be a positive sequence, $(p_k)_{k \geq 1}$ be a sequence such that $0 < p_k \leq 1$ for all $k \geq 1$. Consider a prior Π on coefficients $(f_k)_{k \geq 1}$ with

$$f_k = \sigma_k \zeta_k,$$

where ζ_k are independent p_k -exponential random variables with respective densities h_{p_k} defined in (3). For some $\beta > 0$, consider the following quantities:

$$r_k := \frac{k^{-\beta-1/2}}{\sigma_k} \quad \text{and} \quad z_k := \frac{r_k^{p_k}}{p_k}. \quad (14)$$

The next Theorem provides a result under generic conditions; practical choices of p_k are considered in Corollary 1. For any positive number $s > 0$, we denote

$$N_s := \lfloor n^{\frac{1}{2s+1}} \rfloor. \quad (15)$$

Theorem 3. Let $f_0 \in \mathcal{F}^\beta(L)$ for some $\beta > 0$ and $L \geq 1$. Assume $X^{(n)} \sim P_{f_0}^{(n)}$ from the model (7). Consider the prior Π defined above this statement and assume that $(\sigma_k)_{k \geq 1}$ and $(p_k)_{k \geq 1}$ are chosen such that, for n large enough,

$$\forall k > N_\beta, \quad r_k^{p_k} \geq 8. \quad (16)$$

Furthermore, assume there exists a constant $\eta > 1$, such that

$$\sum_{k \geq 1} \exp\{-z_k/4\} < +\infty \quad \text{and} \quad \sum_{k \leq N_\beta} z_k \lesssim N_\beta \log^\eta n. \quad (17)$$

Then, for any $\rho \in (0, 1)$, there exist a constant $\eta' > 2\eta$, such that, as $n \rightarrow \infty$, we have

$$E_{f_0} \Pi_\rho \left[\|f - f_0\|_2 \leq n^{-\frac{\beta}{2\beta+1}} \log^{\eta'} n \mid X \right] \rightarrow 1.$$

The proof of Theorem 3 can be found in Section B.3. Theorem 3 provides prior conditions on the scaling σ_k and tail parameters p_k for which the posterior contracts at minimax rate $n^{-\beta/(2\beta+1)}$ up to logarithmic factors. Compatibility condition (16) ensures that the prior is oversmoothing in a way (roughly $\sigma_k \lesssim k^{-\beta-1/2}$) and that p_k does not go too fast to 0; an examination of Condition (17) suggests that a natural choice for $1/p_k$ is a logarithmic-type growth in terms of k . This is confirmed in the next Corollary 1, whose proof can be found in Section B.4, which provides two natural examples for which these conditions are satisfied.

Corollary 1. *The conditions of Theorem 3 are met for the following choices of (p_k) , (σ_k)*

1. *Polynomial decay of the scaling. For $\alpha > \beta > 0$, $p_1 = p_2 = 1$,*

$$\sigma_k = k^{-1/2-\alpha} \quad \text{and} \quad \forall k \geq 3, \quad p_k = (\log k)^{-1} \log \log k. \quad (18)$$

2. *Faster than polynomial decay. For $\gamma > 0$, $p_1 = p_2 = 1$ and $c > 2.1 > \log 8$,*

$$\sigma_k = \exp(-\log^{1+\gamma} k) \quad \text{and} \quad \forall k \geq 3, \quad p_k = c(\log k)^{-(1+\gamma)}. \quad (19)$$

The first natural choice matches the oversmoothing polynomial choice of sequence σ_k in Theorem 1. Compared to Theorem 1 (fixed p) for which the rate was polynomially slower than minimax, now choosing p_k going to 0 leads to contraction at quasi-minimax rate (up to log factors). With the choice (18) one obtains one-sided adaptation on the range $\alpha > \beta$. In order to obtain full minimax adaptation, we use the second choice of sequence $\sigma_k = \exp(-\log^{1+\gamma} k)$ which ensures oversmoothing thanks to its faster than polynomial decay. This automatic oversmoothing choice matches the ones introduced in Agapiou and Castillo [2024], Agapiou et al. [2026] and similarly shows full adaptation without being restricted by hyperparameters (the choice of γ is free and need not depend on β).

3. Overparameterized shallow neural network priors

In this section we focus on random-design nonparametric Gaussian regression. Given an integer $n \geq 1$, we observe n i.i.d. pairs of random variables $(X_i, Y_i) \in [0, 1] \times \mathbb{R}$, with

$$Y_i = f(X_i) + \xi_i, \quad (20)$$

where (X_i) are i.i.d. from a distribution P_X on $[0, 1]$ and independently ξ_i are i.i.d. $\mathcal{N}(0, 1)$ variables. For simplicity, we work in dimension 1, but all our results extend to an input space $[0, 1]^d$, $d \geq 2$, in a straightforward way. The sample $((X_i, Y_i))_{1 \leq i \leq n}$ has distribution denoted P_f^n , where P_f is the law of

a single pair (X_i, Y_i) ; the corresponding log-likelihood is

$$\ell_n(f, (X, Y)) := -\frac{1}{2} \sum_{i=1}^n (Y_i - f(X_i))^2.$$

Given a prior Π on $f : [0, 1] \rightarrow \mathbb{R}$ to be defined below, the ρ -posterior $\Pi_\rho[\cdot | X, Y]$ is formed using (1). We study it assuming $((X_i, Y_i))_{1 \leq i \leq n} =: (X, Y)$ is distributed from $P_{f_0}^n$, where the true f_0 belongs to a Hölder ball: denoting by $D^j f$ the j -th derivative of f , define

$$\mathcal{H}^\beta(L) := \left\{ f : [0, 1] \rightarrow \mathbb{R} : \max_{0 \leq i \leq \underline{\beta}} \|D^i f\|_\infty + \sup_{x \neq y} \frac{|D^{\underline{\beta}} f(x) - D^{\underline{\beta}} f(y)|}{|x - y|^{\beta - \underline{\beta}}} \leq L \right\}, \quad (21)$$

for $L \geq 0$, $\beta \in (0, 2]$ and where $\underline{\beta} \in \{0, 1\}$ is the largest integer strictly smaller than β . We aim to characterize the contraction rate of $\Pi_\rho[\cdot | X, Y]$ around f_0 using the population loss

$$\|f - f_0\|_{2, P_X}^2 := \int (f - f_0)^2 dP_X. \quad (22)$$

3.1. Definition of the prior

Shallow neural networks. Functions in $\mathcal{H}^\beta(L)$ can naturally be approximated by shallow neural networks (henceforth SNN). The realization of a shallow ReLU neural network with $M \geq 1$ neurons is a function

$$f_M : x \in [0, 1] \mapsto \sum_{k=0}^{M-1} w_k (v_k \times x + a_k)_+ + b, \quad (23)$$

where $b \in \mathbb{R}$ and $w_k, v_k, a_k \in \mathbb{R}$ for all $k \in \{0, \dots, M-1\}$. The ReLU activation function $x \mapsto (x)_+ := \max(0, x)$ is piecewise linear, so that the function f_M is piecewise affine.

To construct an approximation of a smooth function $f \in \mathcal{H}^\beta(L)$ with an SNN, one can consider a uniform subdivision of $[0, 1]$ in M intervals $I_k := [k/M, (k+1)/M)$, and approximate f by the piecewise affine function f_M^* changing slope and interpolating f precisely at the boundary points of I_k . Lemma 10 (see Section D.3) formalizes this intuition and controls the approximation error. For any $s \in \mathbb{R}$, let us denote $N_s := 2^{m_s}$, where m_s is the closest integer solution to $2^{m_s} = n^{1/(2s+1)}$, for $n \in \mathbb{N}$.

Ideal ‘oracle’ approximator. Suppose first, to fix ideas, that the smoothness parameter β of $f_0 \in \mathcal{H}^\beta(L)$ is known. In this case one could set $M = N_\beta$ and

$$f_{N_\beta}^*(x) = f_0(0) + \sum_{k=0}^{N_\beta-1} w_{0;k} \left(x - \frac{k}{N_\beta} \right)_+. \quad (24)$$

Lemma 10 implies $\|f_0 - f_{N_\beta}^*\|_\infty \leq 2LN_\beta^{-\beta}$, for $\beta \in (0, 2]$, which is the optimal approximation error for β -Hölder functions.

Priors on overparameterized shallow neural networks. We now define an *overparameterized* prior on shallow networks (that is, with possibly much larger than oracle number of neurons). Let us choose

a number of neurons $N_\alpha \geq N_\beta$ (equivalently $\alpha \leq \beta$; typically one can think of the choice $N_\alpha = n$, or $N_\alpha = n^{1-\delta}$ for some small $\delta > 0$) and set

$$f = \sum_{k=0}^{N_\alpha-1} w_k (x - a_k)_+ + b, \quad w_k \stackrel{iid}{=} \sigma_n \zeta_k, \quad a_k = k/N_\alpha, \quad b \sim \pi_b, \quad (25)$$

for some deterministic $\sigma_n > 0$ to be chosen, ζ_k independent and identically distributed random variables with density h satisfying conditions (4)–(6), and where the prior π_b on the bias is any distribution with symmetric, continuous and strictly positive density on $\mathbb{R}$, independent of the prior on the weights (w_k) (note also that, for simplicity, we have taken the shifts a_k to be deterministic).

3.2. Contraction results for overparameterized SNN priors

For any $\rho \in (0, 1)$ and P, Q probability measures, the ρ -Rényi divergence is defined as

$$D_\rho(P, Q) := \frac{1}{\rho - 1} \log \int \left(\frac{dP}{dQ} \right)^\rho dQ. \quad (26)$$

The next result examines contraction rates of ρ -posterior distributions under SNN priors.

Theorem 4. *Let $\beta \in (0, 2]$ and $f_0 \in \mathcal{H}^\beta(L)$. Consider data $(X_i, Y_i)_{1 \leq i \leq n}$ generated from $P_{f_0}^n$ in model (20). Let $0 \leq \alpha \leq \beta$ and Π be the overparameterized SNN prior defined in (25), with $0 < p \leq 1$, $q > 0$ and some σ_n to be specified below. For any $\rho \in (0, 1)$ and D_ρ as in (26), there exists a large enough constant $M > 0$, such that, as $n \rightarrow \infty$,*

$$E_{f_0} \Pi_\rho \left[\left\{ f : \frac{1}{n} D_\rho(P_f^n, P_{f_0}^n) \geq M \varepsilon_n^2 \right\} \mid X, Y \right] \rightarrow 0,$$

where, letting $\varepsilon_n^* = n^{-\frac{\beta}{1+2\beta}}$ and $\varepsilon_n^+ = n^{-2/5}$ (equal to ε_n^* for $\beta = 2$), ε_n is given by

$$i) \text{ (Oracle } \sigma_n) \text{ for } \sigma_n = N_\alpha^{-\frac{2}{2+p}} N_\beta^{\frac{2}{2+p}-\beta} \log^{-\frac{2}{q(2+p)}}(n),$$

$$\varepsilon_n = \varepsilon_n^* (N_\alpha N_\beta)^{\frac{p}{2+p}} \log^{-\frac{p}{q(2+p)}}(n);$$

$$ii) \text{ (Non-oracle } \sigma_n) \text{ for } \sigma_n = \varepsilon_n^+ / N_\alpha$$

$$\varepsilon_n = \varepsilon_n^* n^p \left(\frac{1-\beta}{2+4\beta} + \frac{1}{5} + \frac{1}{2+4\alpha} \right).$$

Corollary 2. *In the setting of Theorem 4, assume further $\|f_0\|_\infty \leq F$, for some $F > 0$. Define $T_F : f \mapsto -F \vee (f \wedge F)$ and consider the “clipped” posterior $\Pi_\rho^F[\cdot \mid X, Y] = \Pi_\rho[\cdot \mid X, Y] \circ T_F^{-1}$. As $n \rightarrow \infty$,*

$$E_{f_0} \Pi_\rho^F \left[\{ f : \|f - f_0\|_{2, P_X} \geq M' \varepsilon_n \} \mid X, Y \right] \rightarrow 0,$$

where $M' > 0$ is a large enough constant (depending on F) and ε_n is defined as in Theorem 4.

The proofs of Theorem 4 and Corollary 2 can be found in Section B.5. These results give conditions on the deterministic prior scaling σ_n to obtain posterior contraction rates in $L_2(P_X)$ -loss (or in Rényi divergence if clipping is not applied) with overparameterized SNN p -tailed priors. The first choice of σ_n is said to be oracle as it depends on the unknown ‘true’ smoothness β , while the second choice does not. Although both lead to polynomially slower than minimax rates (here the minimax rate is ε_n^*), similarly to the series priors in Section 2, this extra polynomial factor decreases as p goes to 0.

Remark 2. *The case of lighter tails is also interesting (since it includes the case of Gaussian priors on the weights and bias) and is examined in Theorem 9 in the supplement. As expected, the rate becomes worse as p increases.*

To obtain a minimax adaptive contraction rate, we now devise an overparameterized SNN prior with suitably chosen deterministic prior scalings σ_n and decaying tail parameter p_n .

Theorem 5. *Let $\beta \in (0, 2]$, and $f_0 \in \mathcal{H}^\beta(L)$. Consider data $(X_i, Y_i)_{1 \leq i \leq n}$ generated from $P_{f_0}^n$ in model (20). Let Π be the overparameterized SNN prior defined as in (25), for $\alpha = 0$ (hence the width is $N_\alpha \asymp n$), $\sigma_n = n^{-t}$ with $t > 2.5$, and ζ_k independent and identically distributed according to p_n -exponential distributions (defined in (3)) with $p_n = 2/\log n$, $n \geq 8$. For any $\rho \in (0, 1)$, there exists a large enough constant $M > 0$, such that, as $n \rightarrow \infty$,*

$$E_{f_0} \Pi_\rho \left[\left\{ f : \frac{1}{n} D_\rho(P_f^n, P_{f_0}^n) \geq M \varepsilon_n^2 \right\} \mid X, Y \right] \rightarrow 0,$$

for any $\varepsilon_n \geq \varepsilon_n^* \sqrt{\log n}$, where $\varepsilon_n^* = n^{-\frac{\beta}{1+2\beta}}$ and D_ρ is Rényi divergence. Assuming further $\|f_0\|_\infty \leq F$ and considering the clipped posterior $\Pi_\rho^F[\cdot \mid X, Y]$ as in Corollary 2, for a large enough constant $M' > 0$, as $n \rightarrow \infty$,

$$E_{f_0} \Pi_\rho^F \left[\left\{ f : \|f - f_0\|_{2, P_X} \geq M' \varepsilon_n \right\} \mid X, Y \right] \rightarrow 0.$$

The proof can be found in Section B.6. Theorem 5 shows that, for an overparameterized (with $\alpha = 0$) SNN prior with p_n -exponential weights rescaled by a sufficiently small polynomial factor σ_n and sufficiently heavy tails $p_n = 2/\log n$, we obtain a minimax fully *adaptive* contraction rate (up to a $\sqrt{\log n}$ factor). This choice of tail parameter $p_n = 2/\log n$ on an SNN p -exponential prior can be (informally) thought of as having a deep neural network prior with i.i.d. Gaussian weights and depth $L = \log n$ (see Section 5 for more discussion).

While the last result is theoretically appealing since it establishes adaptation over the full range $\beta \in (0, 2]$ up to logarithmic factors, neural network priors of width n can become computationally demanding for large sample sizes. To alleviate this, one can instead use networks of slightly smaller width N_α , $0 \leq \alpha < 2$ (and small α e.g. $\alpha = .5$), still yielding adaptation over the range $\beta \in [\alpha, 2]$. Also, the condition $t > 2.5$ on the scalings σ_n , can be relaxed, facilitating posterior sampling. See Remarks 4 and 5 in the supplement for details.

Moreover, the proof of Theorem 5 carries over, with minor modifications, to the choice

$$\sigma_n = \exp(-a(\log n)^{1+\gamma}), \quad p_n = \frac{b}{(\log n)^{1+\gamma}}, \quad (27)$$

for $\gamma > 0$ and $a, b > 0$ such that $ab > \log 8$, corresponding to the second regime of Corollary 1. See Remark 3 in the supplement for details.

4. Simulations

In this section we provide a numerical simulation study. Even though results with series priors in Section 2 were formulated in the white noise model, we compare the performance of the various priors considered in the following random design regression setting:

$$y_i = f(x_i) + \xi_i, \quad i = 1, \dots, n, \quad (28)$$

where $f : [0, 1] \rightarrow \mathbb{R}$ unknown, the $x_i \in [0, 1]$ are i.i.d. with uniform distribution on $[0, 1]$ and ξ_i are independent $N(0, \sigma^2)$ for $\sigma = 1/4$ (the latter value has no special meaning and is chosen for easier visualization in the pictures below).

We define a true function through the series expansion

$$f_0(x) = \sum_{k=1}^{\infty} f_{0,k} \varphi_k(x),$$

for $\varphi_k(x) = \sqrt{2} \cos((k - 1/2)\pi x)$ and $f_{0,k} = (3/2)k^{-3/2} \sin k$. This function has regularity $\beta = 1$ in the sense of (11). We generate noisy observations $Y = \{y_1, \dots, y_n\}$ according to the above random design regression model, for $n = 400$ and $n = 4000$. We consider the following priors:

- series priors as in (9), (10) defined over the same basis as the truth, $\{\varphi_k\}$, with p -exponential tails for $p = 2, 1, 1/2, 1/4$ and $\alpha = 2$;
- series priors defined over $\{\varphi_k\}$, with p_k -exponential tails varying with frequency k (and n), either as (18) for $\alpha = 2$ or as (19) for $\gamma = 1/2$;
- Cauchy HT(α) for $\alpha = 2$ and OT for $\gamma = 1/2$ series priors defined over $\{\varphi_k\}$. These are series priors as in (9) but with Cauchy-distributed ζ_k , with $\sigma_k = k^{-1/2-\alpha}$ or $\sigma_k = \exp(-\log^{1+\gamma} k)$, respectively. We use them as a benchmark as they have been shown to be partially adaptive for $\alpha \geq \beta$ and fully adaptive for any $\gamma > 0$, respectively, Agapiou and Castillo [2024], Agapiou et al. [2026];
- shallow neural network priors defined as in (25) with $\alpha = 1/2$, for $p = 2, 1, 1/2, 1/4$ and oracle choice of σ_n , see Theorem 4(i);
- shallow neural network priors defined as in (25) with $\alpha = 1/2$, for $p = 2, 1, 1/2, 1/4$ and $\sigma_n = \varepsilon_n^+ / N_\alpha$ as in Theorem 4(ii);
- shallow neural network priors defined as in (25) with $\alpha = 1/2$ and p varying with n , $p_n = 1/\log n$, $\sigma_n = n^{-7/5-0.01}$, and with $\alpha = 0$ (width- n) for p_n, σ_n as in (27), with $\gamma = 1/2$, $a = 1/4$ and $b = 4 \log 8 + 0.01$.

Note that compared to the shallow neural network priors, the series priors are somewhat *favorable* by the fact that the basis $\{\varphi_j\}$ on which they are defined coincides with the one on which the true function is defined, which is not the case for the ReLU prior. In particular, both the true function and draws from the studied series priors are restricted to take the value zero at the right edge ($x = 1$). Similarly, shallow neural networks with the *oracle* choice of σ_n use knowledge of the regularity β of the truth f_0 , which is unrealistic in practice. For the series priors, we chose $\alpha = 2$ so that we are in the oversmoothing prior regime $\alpha > \beta$, in which the differences in performance depending on p arise. For shallow neural network priors, we chose $\alpha = 1/2$ so that we are in the overparameterized regime studied in Section 3. Note that the first varying- p shallow network prior used (corresponding to $\alpha = 1/2, p_n = 1/\log n, \sigma_n = n^{-7.5-0.01}$) is not strictly covered by the results of Section 3, however our theory suggests that it should perform well, at least asymptotically for large n (see Remarks 4 and 5 in the supplement). For the second such prior, a and b were chosen so that p_n did not become too

small for the sample sizes n considered, while still satisfying the conditions under which our theory holds. Smaller values of p_n necessitated a smaller step size in the posterior sampling algorithm used and hence more iterations (see the next paragraph for details). As this prior already incurred a higher computational cost due to its width n , compared with the width $\sqrt{n}$ of the other neural network priors considered, the resulting increase in the number of iterations would have made posterior sampling computationally expensive. Furthermore, in Section 3, for convenience we assumed p -exponential priors on the weights as in (3), while in our implementation we use p -exponential tailed distributions in the broader sense of (4)–(6). To summarize what we expect from our theory (and which is fully empirically confirmed by the results below), the two Cauchy priors and the varying p -exponential tail series and neural network priors are expected to be optimal, while for the rest of the priors, smaller p is better.

Regarding the implementation, we discretized the unit interval using 200 uniformly spaced points, while we truncated the spectral series (for the truth or the series priors) up to $k = 200$. For sampling the posterior we employed the (whitened) preconditioned Crank-Nicholson (wpCN) algorithm, [Chen et al. \[2018\]](#), [Cotter et al. \[2013\]](#), which is a derivative-free Metropolis-Hastings algorithm robust with respect to dimension (truncation level). The vanilla version of pCN is suitable for Gaussian priors, while the whitened version is suitable for non-Gaussian priors admitting a transformation $f = T(\xi)$, where ξ is a sequence of i.i.d. standard normal variables and f is a random draw from the non-Gaussian prior of interest, see Algorithm 2 in [Chen et al. \[2018\]](#). For example, to get series priors with p -exponential tails for $p > 0$, we employed the transformation $T(\xi) = |\xi|^{2/p-1}\xi$ which transforms $\xi \sim N(0, 1)$ to a random variable with p -exponential tails, while for the shallow neural network priors this transformation is composed with the map taking the sequence of weights and biases to function realizations as in (25). For all considered priors, we initialized the Markov chains using draws from the prior. In all runs, we tuned the proposal step size (i.e., the scaling of the proposals) to achieve an acceptance rate of approximately 30%. For all priors, the wpCN algorithm is less efficient compared to the Gaussian series prior case, in the sense that smaller step sizes are required to maintain this target acceptance rate. To keep the total distance explored by the sampler roughly constant across experiments, we scaled the total number of iterations inversely proportional to the step size. In all runs, we retained 20,000 samples. Depending on the prior used, a thinning factor proportional to the scaling applied to the total number of iterations was used.

To compare the performance of the considered priors quantitatively, we averaged errors over 100 realizations of the data, for each of $n = 400$ and $n = 4000$. Figure 1 shows one such realization for each choice of n . In particular, we employed two types of errors. The first one is the L_2 -error of the posterior means, hence after averaging we estimate the error $E_{f_0} \|\hat{f} - f_0\|_2$, for $\hat{f}$ the posterior mean. The second error estimates

$$E_{f_0} E_{\Pi[\cdot|Y]} \|f - f_0\|_2,$$

where the inner expectation is estimated by taking the average of the L_2 -errors of the (thinned) Markov chain samples after burn-in, and the outer by averaging over the 100 data realizations. The latter error captures the contraction of the whole posterior around f_0 .

The computed average errors are presented in Table 1. We also computed standard deviations which were of lower order compared to the averages, hence for ease of readability we did not include them. The errors appear to be consistent with our theory. Overall, the best performers are the priors with heavier tails (either small p or Cauchy) and the (diminishing) varying tail priors. Shallow neural network priors with the more practically relevant choice of $\sigma_n = \varepsilon_n^+ / N_\alpha$, for small p have similar performance to the corresponding shallow neural network priors with oracle σ_n .

Figures 2 and 3 show posterior means and 95% credible intervals for the various priors, and for $n = 400$ and $n = 4000$, respectively, for the data realizations shown in Figure 1. The conclusions are aligned to the ones in the previous paragraph.




Figure 1. Observed random design regression data, with $n = 400$ (left) and $n = 4000$ (right). True function f_0 in black solid, red points noisy observations according to model (28).

Prior	$n = 400$		$n = 4000$	
	Err A	Err B	Err A	Err B
Series $p = 2, \alpha = 2$	0.1194	0.1226	0.0742	0.0749
Series $p = 1, \alpha = 2$	0.0895	0.0962	0.0477	0.0495
Series $p = 1/2, \alpha = 2$	0.0814	0.0919	0.0400	0.0431
Series $p = 1/4, \alpha = 2$	0.0915	0.1050	0.0401	0.0444
Series varying $p, \alpha = 2$	0.0851	0.0984	0.0383	0.0427
Series varying $p, \gamma = 1/2$	0.0950	0.1184	0.0398	0.0461
Cauchy HT(α), $\alpha = 2$	0.0807	0.0920	0.0394	0.0431
Cauchy OT, $\gamma = 1/2$	0.0707	0.0906	0.0349	0.0397
SNN $p = 2, \alpha = 1/2$, oracle σ_n	0.2901	0.2908	0.2113	0.2114
SNN $p = 1, \alpha = 1/2$, oracle σ_n	0.1539	0.1559	0.1080	0.1082
SNN $p = 1/2, \alpha = 1/2$, oracle σ_n	0.0793	0.0837	0.0497	0.0505
SNN $p = 1/4, \alpha = 1/2$, oracle σ_n	0.0664	0.0718	0.0410	0.0423
SNN $p = 2, \alpha = 1/2, \sigma_n = \varepsilon_n^+/N_\alpha$	0.8569	0.8570	0.9018	0.9018
SNN $p = 1, \alpha = 1/2, \sigma_n = \varepsilon_n^+/N_\alpha$	0.3467	0.3472	0.3473	0.3474
SNN $p = 1/2, \alpha = 1/2, \sigma_n = \varepsilon_n^+/N_\alpha$	0.0999	0.1031	0.0797	0.0801
SNN $p = 1/4, \alpha = 1/2, \sigma_n = \varepsilon_n^+/N_\alpha$	0.0675	0.0727	0.0498	0.0509
SNN varying $p, \alpha = 1/2$	0.0729	0.0776	0.0481	0.0492
SNN varying $p, \alpha = 0, \gamma = 1/2$	0.0681	0.0750	0.0265	0.0290

Table 1. L_2 average errors of posterior mean (Error A) and L_2 contraction-type errors (Error B) for the considered priors, with number of observations $n = 400$ and $n = 4000$.

5. Discussion

This work shows that equipping the coefficients of a function (on a basis, or on a ReLU dictionary) with p -exponential priors leads to near (small p) or full (regime $p \rightarrow 0$) adaptation to smoothness. The presented results are obtained for simplicity in white noise (for series priors) and random design regression (for neural network priors). However, they hold much more generally for ρ -posteriors in other statistical models, see Section C.1 of the supplement.

We now comment on the overparameterized p -exponential shallow neural network prior. The prior architecture we have chosen originates from the piecewise affine approximation result Lemma 10. To recover f_0 , the prior randomly draws weights in the hidden layer and the weights of the input layer are fixed on a certain grid. This specific architecture allows us to define a “simplest possible”




Figure 2. Random design regression: true function (black), posterior mean (blue), 95% credible regions (grey), for $n = 400$. Top row: p -exponential series priors with $\alpha = 2$ and $p = 2, 1, 1/2, 1/4$ left to right. Second row: series priors with varying p -tails as in (18) with $\alpha = 2$ and as in (19) with $\gamma = 1/2$, Cauchy HT(α) with $\alpha = 2$ and Cauchy OT with $\gamma = 1/2$, left to right. Third row: shallow neural network priors as in (25) with $\alpha = 1/2$, oracle choice of σ_n and $p = 2, 1, 1/2, 1/4$ left to right. Fourth row: shallow neural network priors as in (25) with $\alpha = 1/2$, $\sigma_n = \varepsilon_n^+ / N_\alpha$ and $p = 2, 1, 1/2, 1/4$ left to right. Bottom row: shallow network priors as in (25) with $\alpha = 1/2$, $\sigma = n^{-7/5-0.01}$, $p_n = 1/\log n$ and with $\alpha = 0$, $\sigma_n = \exp(-(\log n)^{3/2}/4)$, $p_n = (4 \log 8 + 0.01)/(\log n)^{3/2}$.

overparameterized SNN, having much more active neurons in the hidden layer compared to the oracle network in Lemma 10. This choice also allows us to easily compare performances of different choices of p -exponential prior distributions on the weights. In particular, we show (e.g. in Theorem 4) that on this architecture heavier tailed priors perform better (than Gaussian for example). We now informally link this p -exponential shallow prior to deeper Gaussian priors. The recent works Vladimirova et al. [2019], Zavatone-Veth and Pehlevan [2021], Noci et al. [2021] have shown that in a deep NN with i.i.d. Gaussian weights, the distribution of the output of a layer at depth $L \geq 1$ is a random vector whose




Figure 3. Same setting as Figure 2 but here with $n = 4000$

coordinates (conditionally on the input $\mathbf{x}$ of the NN) are *marginally* $2/L$ -exponential, in the sense of equation (5) with $q = 2/L$ (see for instance Theorem 3.1 in [Vladimirova et al. \[2019\]](#)). The p -SNN prior in (25) can then be viewed as a ‘summary’ of an *i.i.d.* deep Gaussian prior with depth $L = 2/p$. Indeed, we take for prior on the single hidden layer a distribution whose marginals are the same as the output of an L -layered Gaussian DNN and removing dependencies. With this analogy, the improved performances of heavier priors (e.g. in Theorem 4) can be related to improved performances of DNN priors as the depth increases. Particularly looking at Theorem 5, we show that taking $p = 2/\log n$ provides a smoothness-adaptive procedure. Such result can be related to those of [Castillo and Egels \[2025\]](#) who showed that overparameterized heavy-tailed (with heavier polynomial tails rather than p -exponential as in the present work) DNN priors of depth $L \gtrsim \log n = 2/p$ are fully adaptive. Our results provide a first link between the theoretical results for overparameterized heavy-tailed priors and the more practically employed overparameterized Gaussian DNN prior (we are not aware of any

existing posterior contraction results for such Gaussian priors). The case of purely Gaussian weights for deeper networks will be addressed in future work.

Finally, for simplicity, we restricted to results to fractional posteriors with parameter $\rho < 1$. The main technical reason is that one does not need to build sieve sets and control their entropy, as is usually the case for classical posteriors ($\rho = 1$) when using the generic posterior contraction theory [Ghosal et al. \[2000\]](#). Still, we believe all our results go through also for $\rho = 1$: this can be formally proved in the white noise model (see Section C.4 in the Supplement for an example of such a result), but for more general models such as random design regression or classification, it is currently an open question; we refer to the discussion in [Agapiou and Castillo \[2024\]](#) for more details on this point.

6. Proof of Theorem 1

Throughout this section, the notation $a_n \lesssim b_n$ denotes inequalities up to a positive constant.

Proof. First, thanks to Lemmas 4 and 5, in the Gaussian white noise model (7), to obtain ρ -posterior contraction in L_2 -distance, it is enough to show, for some constant $C, D > 0$,

$$\Pi [\|f - f_0\|_2 \leq D\varepsilon_n] \geq \exp \left\{ -Cn\varepsilon_n^2 \right\}.$$

Let $(\delta_k)_{k \geq 1}$ be a $(n$ -dependent) sequence of positive numbers, to be chosen below, that verifies, for a large enough constant $D \geq 1$,

$$\sum_{k=1}^{\infty} \delta_k^2 \leq (D\varepsilon_n)^2. \quad (29)$$

By combining Parseval's identity and (29), the event $\{f = (f_k) : (f_k - f_{0,k})^2 \leq \delta_k^2 \text{ for all } k \geq 1\}$ is included in $\{f : \|f - f_0\|_2 \leq D\varepsilon_n\}$. Using that the variables $f_k = \sigma_k \zeta_k$ are independent, one deduces that it is enough to verify the inequality

$$\prod_{k \geq 1} \Pi [|f_k - f_{0,k}| \leq \delta_k] \geq \exp \left\{ -Cn\varepsilon_n^2 \right\}. \quad (30)$$

We now bound from below the individual probabilities in the last product, that is

$$\Pi [|f_k - f_{0,k}| \leq \delta_k] = P [|\sigma_k \zeta_k - f_{0,k}| \leq \delta_k] \geq c_0 \int_{(f_{0,k} - \delta_k)/\sigma_k}^{(f_{0,k} + \delta_k)/\sigma_k} e^{-c_1 |x|^p} dx, \quad (31)$$

for all $k \geq 1$. Since the last integrand is symmetric, we can assume without loss of generality that $f_{0,k} \geq 0$ in the bounds to follow.

We distinguish the two cases $\alpha > \beta$ and $\alpha \leq \beta$. First suppose $\alpha > \beta$. Recall the definitions of γ and N_γ in (12)–(15) and let us choose δ_k as follows

$$\delta_k := \begin{cases} 1/\sqrt{n}, & 1 \leq k \leq N_\gamma, \\ 2Lk^{-1/2-\beta}, & k > N_\gamma \end{cases}. \quad (32)$$

Let us check that this choice satisfies the constraint (29). By definition, for some $C_\beta > 0$,

$$\sum_{k=1}^{\infty} \delta_k^2 \leq \frac{N_\gamma}{n} + 4L^2 C_\beta N_\gamma^{-2\beta}.$$

Recall (13) the definition of ε_n , since $\alpha > \beta$, one has $\gamma > \beta$, so that $N_\gamma \leq N_\beta$ and $\varepsilon_n^2 \geq n^{-2\beta/(2\beta+1)} \geq N_\beta/n \geq N_\gamma/n$. Also, $N_\gamma^{-2\beta} \leq 2\varepsilon_n^2$ for large enough n (using $\lfloor x \rfloor^{-2\beta} \leq 2x^{-2\beta}$ for large x), so that the last display is bounded from above by $(1 + 8L^2C_\beta)\varepsilon_n^2$.

To bound (31) from below, let us first consider the case of indices $1 \leq k \leq N_\gamma$. In that case we bound from below the integrand in (31) by its smallest value, attained at $x = (f_{0,k} + \delta_k)/\sigma_k$ (since $f_{0,k} \geq 0$), so that, using $(a+b)^p \leq \kappa_p(a^p + b^p)$ ($\kappa_p = 1$ for $p \in (0, 1]$),

$$\begin{aligned} \Pi[|f_k - f_{0,k}| \leq \delta_k] &\geq \frac{2c_0\delta_k}{\sigma_k} \exp\left\{-\frac{c_1}{\sigma_k^p}(f_{0,k} + \delta_k)^p\right\} \\ &\geq \frac{2c_0\delta_k}{\sigma_k} \exp\left\{-\frac{c_1}{\sigma_k^p}\kappa_p(f_{0,k}^p + n^{-p/2})\right\} \geq \frac{2c_0\delta_k}{\sigma_k} \exp\left\{-\frac{c_2}{\sigma_k^p}2\kappa_p(Lk^{-1/2-\beta})^p\right\}, \end{aligned}$$

using the definition of δ_k , the regularity condition on f_0 and that $Lk^{-1/2-\beta} \gtrsim 1/\sqrt{n}$ for all $1 \leq k \leq N_\gamma$ for large enough n (since $N_\gamma \leq N_\beta$ for $\alpha > \beta$). Deduce, for such k 's and n 's,

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \frac{2c_0}{\sqrt{n}\sigma_k} \exp\left\{-2\kappa_p c_2 (Lk^{\alpha-\beta})^p\right\}.$$

In order to bound the product of these probabilities from below in (30), we distinguish two cases. If $p \leq 2$, one notes that $\gamma \leq \alpha$ by definition, so that $\sigma_k^{-1} \geq k^{1/2+\gamma}$. Lemma 6 (applied with α therein replaced by γ) then implies

$$\prod_{k=1}^{N_\gamma} \frac{2c_0}{\sqrt{n}\sigma_k} \geq e^{-(1/2+\gamma-\log(2c_0))N_\gamma} \geq e^{-C_0N_\gamma},$$

for some $C_0 > 0$. Then using the bound $\sum_{k=1}^N k^a \lesssim N^{a+1}$ for any $a > 0$ and integer N ,

$$\prod_{k=1}^{N_\gamma} \Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \exp\left\{-C_0N_\gamma - C_2N_\gamma^{p(\alpha-\beta)+1}\right\} \geq \exp\left\{-C_3N_\gamma^{p(\alpha-\beta)+1}\right\} \geq \exp\left\{-C_3n\varepsilon_n^2\right\},$$

noticing that $N_\gamma^{p(\alpha-\beta)+1} \leq n\varepsilon_n^2$ follows from the definitions of N_γ and ε_n . In the case that $p \geq 2$, one instead applies Lemma 7. Noting that the lower bound in that Lemma is itself bounded from below by $\exp(-C_4N_\gamma^{p(\alpha-\beta)+1})$, one obtains

$$\prod_{k=1}^{N_\gamma} \frac{2c_0}{\sqrt{n}\sigma_k} \geq \exp\left\{-\log(2c_0)N_\gamma - C_4N_\gamma^{p(\alpha-\beta)+1}\right\} \geq e^{-C_5N_\gamma^{p(\alpha-\beta)+1}},$$

so that the product in the last but one display can be bounded from below by $e^{-C_6N_\gamma^{p(\alpha-\beta)+1}}$ and same argument as for the case $p \leq 2$ can be used.

To bound (31) from below for indices $k > N_\gamma$, the choice of δ_k in (32) ensures that

$$[f_{0,k} - \delta_k, f_{0,k} + \delta_k] \supset [-Lk^{-1/2-\beta}, Lk^{-1/2-\beta}]$$

holds, using (32) and the regularity condition on f_0 . Further bounding the probability in (31) from below gives, in this case

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \Pi[|f_k| \leq Lk^{-1/2-\beta}] \geq \Pi[|\zeta_k| \leq Lk^{\alpha-\beta}].$$

Since ζ_k 's have density h and survival function $\bar{H}$, one has $\Pi[|\zeta_k| \leq Lk^{\alpha-\beta}] = 1 - 2\bar{H}(Lk^{\alpha-\beta})$. Gathering these bounds and combining with Condition (6) gives

$$\begin{aligned} \prod_{k > N_\gamma} \Pi[|f_k - f_{0,k}| \leq \delta_k] &\geq \prod_{k > N_\gamma} (1 - 2\bar{H}(Lk^{\alpha-\beta})) \\ &\geq \exp \left\{ \sum_{k > N_\gamma} \log(1 - 2d_0 e^{-d_1(Lk^{\alpha-\beta})^q}) \right\} \geq \exp \left\{ -4d_0 \sum_{k > N_\gamma} e^{-d_1(Lk^{\alpha-\beta})^q} \right\}, \end{aligned}$$

where we have used the inequality $\log(1 - 2x) \geq -4x$, valid for $x \in [0, 1/4]$. Since the series $\sum_k e^{-ck^\delta}$ converges for any given constants $c, \delta > 0$, one deduces that the last display converges to 1 as $n \rightarrow \infty$ and in particular is bounded from below by $1/2$ for n large enough. Gathering the previous bounds gives

$$\prod_{k \geq 1} \Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \exp \left\{ -Cn\varepsilon_n^2 \right\},$$

so that (30) is satisfied for large enough C , concluding the proof in the case $\alpha > \beta$. $\square$

Supplementary Material. The appendix contains the remaining proofs of the main results and some additional results referenced in the main text.

References

- K. Abraham and N. Deo. Deep Gaussian process priors for Bayesian inference in nonlinear inverse problems. *arXiv preprint arXiv:2312.14294*, 2023.
- S. Agapiou and I. Castillo. Heavy-tailed Bayesian nonparametric adaptation. *Ann. Statist.*, 52(4): 1433–1459, 2024. ISSN 0090-5364,2168-8966.
- S. Agapiou and A. Savva. Adaptive inference over Besov spaces in the white noise model using p-exponential priors. *Bernoulli*, 30(3):2275 – 2300, 2024.
- S. Agapiou and S. Wang. Laplace priors and spatial inhomogeneity in Bayesian inverse problems. *Bernoulli*, 30(2):878 – 910, 2024.
- S. Agapiou, J. M. Bardsley, O. Papaspiliopoulos, and A. M. Stuart. Analysis of the Gibbs sampler for hierarchical inverse problems. *SIAM/ASA J. Uncertain. Quantif.*, 2(1):511–544, 2014.
- S. Agapiou, M. Dashti, and T. Helin. Rates of contraction of posterior distributions based on p-exponential priors. *Bernoulli*, 27(3):1616 – 1642, 2021.
- S. Agapiou, I. Castillo, and P. Egels. Heavy-tailed and horseshoe priors for regression and sparse Besov rates. *Bernoulli*, 2026. To appear, arXiv:2505.15543.
- J. Arbel, K. Pitas, M. Vladimirova, and V. Fortuin. A Primer on Bayesian Neural Networks: Review and Debates. *Statistical Science*, 41(2):316 – 353, 2026.
- J. Bai, Q. Song, and G. Cheng. Efficient variational inference for sparse deep learning with theoretical guarantee. In *Advances in Neural Information Processing Systems*, 2020.

- A. Barron, L. Birgé, and P. Massart. Risk bounds for model selection via penalization. *Probab. Theory Related Fields*, 113(3):301–413, 1999. ISSN 0178-8051.
- C. Berenfeld, P. Rosa, and J. Rousseau. Estimating a density near an unknown manifold: A Bayesian nonparametric approach. *The Annals of Statistics*, 52(5):2081 – 2111, 2024.
- I. Castillo. Lower bounds for posterior rates with Gaussian process priors. *Electron. J. Stat.*, 2:1281–1299, 2008.
- I. Castillo. *Bayesian nonparametric statistics*, volume 2358 of *Lecture Notes in Mathematics*. Springer, Cham, 2024. Saint-Flour Probability Summer School LI—2023.
- I. Castillo and P. Egels. Posterior and variational inference for deep neural networks with heavy-tailed weights. *Journal of Machine Learning Research*, 26(122):1–58, 2025.
- I. Castillo and T. Randrianarisoa. Deep horseshoe Gaussian processes. *The Annals of Statistics*, 53(5): 1886 – 1912, 2025.
- I. Castillo, G. Kerkycharian, and D. Picard. Thomas Bayes’ walk on manifolds. *Probab. Theory Related Fields*, 158(3-4):665–710, 2014. ISSN 0178-8051,1432-2064.
- V. Chen, M. M. Dunlop, O. Papaspiliopoulos, and A. M. Stuart. Dimension-robust MCMC in Bayesian inverse problems. 2018. arXiv preprint 1803.03344.
- B.-E. Chérif-Abdellatif. Convergence rates of variational inference in sparse deep learning. In *Proceedings of the 37th International Conference on Machine Learning*, 2020.
- S. Cotter, G. Roberts, A. Stuart, and D. White. MCMC methods for functions: Modifying old algorithms to make them faster. *Statistical Science*, 28(3):424–446, 2013.
- A. Damianou and N. D. Lawrence. Deep Gaussian processes. In *Artificial intelligence and statistics*, pages 207–215. PMLR, 2013.
- E. Dolera, S. Favaro, and M. Giordano. On strong posterior contraction rates for Besov-Laplace priors in the white noise model. *arXiv preprint arXiv:2411.06981*, 2024.
- D. L. Donoho, I. M. Johnstone, G. Kerkycharian, and D. Picard. Wavelet shrinkage: asymptopia? *J. Roy. Statist. Soc. Ser. B*, 57(2):301–369, 1995. ISSN 0035-9246.
- G. Finocchio and J. Schmidt-Hieber. Posterior contraction for deep Gaussian process priors. *Journal of Machine Learning Research*, 24(66):1–49, 2023.
- S. Ghosal and A. van der Vaart. *Fundamentals of nonparametric Bayesian inference*. Cambridge University Press, Cambridge, 2017.
- S. Ghosal, J. K. Ghosh, and A. W. van der Vaart. Convergence rates of posterior distributions. *Ann. Statist.*, 28(2):500–531, 2000.
- E. Giné and R. Nickl. *Mathematical foundations of infinite-dimensional statistical models*, volume 40. Cambridge University Press, 2015.
- M. Giordano. Besov-Laplace priors in density estimation: optimal posterior contraction rates and adaptation. *Electronic Journal of Statistics*, 17(2):2210 – 2249, 2023.
- M. Giordano, K. Ray, and J. Schmidt-Hieber. On the inability of Gaussian process regression to optimally learn compositional functions. In *Advances in Neural Information Processing Systems*, volume 35, 2022.
- B. T. Knapik, B. T. Szabó, A. W. van der Vaart, and J. H. van Zanten. Bayes procedures for adaptive inference in inverse problems for the white noise model. *Probab. Theory Related Fields*, 164(3-4): 771–813, 2016.
- M. Kohler and S. Langer. On the rate of convergence of fully connected deep neural network regression estimates. *The Annals of Statistics*, 49(4):2231 – 2249, 2021.
- I. Kong and Y. Kim. Posterior concentrations of fully-connected Bayesian neural networks with general priors on the weights. *Journal of Machine Learning Research*, 26(94), 2025.
- I. Kong, D. Yang, J. Lee, I. Ohn, G. Baek, and Y. Kim. Masked Bayesian neural networks: Theoretical guarantee and its posterior inference. In *International Conference on Machine Learning*. PMLR,

- 2023.
- K. Lee and J. Lee. Asymptotic properties for Bayesian neural network in Besov space. In *Advances in Neural Information Processing Systems*, volume 35, pages 5641–5653, 2022.
- O. V. Lepskii. A problem of adaptive estimation in Gaussian white noise. *Teor. Veroyatnost. i Primenen.*, 35(3):459–470, 1990.
- O. V. Lepskii. Asymptotically minimax adaptive estimation. I. Upper bounds. Optimally adaptive estimates. *Teor. Veroyatnost. i Primenen.*, 36(4):645–659, 1991.
- A. L’Huillier, L. Travis, I. Castillo, and K. Ray. Semiparametric inference using fractional posteriors. *Journal of Machine Learning Research*, 24(389):1–61, 2023.
- H. Liu, Y.-S. Ong, X. Shen, and J. Cai. When Gaussian process meets big data: A review of scalable GPs. *IEEE transactions on neural networks and learning systems*, 31(11):4405–4423, 2020.
- R. Nakada and M. Imaizumi. Adaptive approximation and generalization of deep neural network with intrinsic dimensionality. *JMLR*, 21(174):1–38, 2020.
- D. Nieman and B. Szabó. Adaptive Sparse Variational Approximations for Gaussian Process Regression. *Bayesian Analysis*, pages 1 – 20, 2025.
- L. Noci, G. Bachmann, K. Roth, S. Nowozin, and T. Hofmann. Precise characterization of the prior predictive distribution of deep ReLU networks. *Advances in Neural Information Processing Systems*, 34:20851–20862, 2021.
- I. Ohn and L. Lin. Adaptive variational Bayes: Optimality, computation and applications. *The Annals of Statistics*, 52(1):335–363, 2024.
- N. G. Polson and V. Ročková. Posterior concentration for sparse deep learning. *Advances in Neural Information Processing Systems*, 31, 2018.
- C. E. Rasmussen and C. K. I. Williams. *Gaussian processes for machine learning*. Adaptive Computation and Machine Learning. MIT Press, Cambridge, MA, 2006.
- J. Rousseau and B. Szabo. Asymptotic behaviour of the empirical Bayes posteriors associated to maximum marginal likelihood estimator. *The Annals of Statistics*, 45(2), 2017.
- J. Schmidt-Hieber. Nonparametric regression using deep neural networks with ReLU activation function. *The Annals of Statistics*, 48(4):1875 – 1897, 2020.
- B. Szabó and Y. Zhu. Vecchia Gaussian processes: Probabilistic properties, minimax rates and methodological developments, 2026. arXiv preprint 2410.10649.
- B. T. Szabó, A. W. van der Vaart, and J. H. van Zanten. Empirical Bayes scaling of Gaussian priors in the white noise model. *Electronic Journal of Statistics*, 7, 2013.
- T. Tang, N. Wu, X. Cheng, and D. Dunson. Adaptive Bayesian regression on data with low intrinsic dimensionality. *The Annals of Statistics*, 54(2):1080 – 1099, 2026.
- A. W. van der Vaart and J. H. van Zanten. Rates of contraction of posterior distributions based on Gaussian process priors. *Ann. Statist.*, 36(3):1435–1463, 2008. ISSN 0090-5364.
- A. W. van der Vaart and J. H. van Zanten. Adaptive Bayesian estimation using a Gaussian random field with inverse gamma bandwidth. *Ann. Statist.*, 37(5B):2655–2675, 2009.
- M. Vladimirova, J. Verbeek, P. Mesejo, and J. Arbel. Understanding priors in Bayesian neural networks at the unit level. In *Proceedings of the 36th International Conference on Machine Learning*, volume 97, pages 6458–6467. PMLR, 2019.
- M. Vladimirova, S. Girard, H. Nguyen, and J. Arbel. Sub-Weibull distributions: Generalizing sub-Gaussian and sub-exponential properties to heavier tailed distributions. *Stat*, 9(1):e318, 2020.
- Y. Yang and D. B. Dunson. Bayesian manifold regression. *The Annals of Statistics*, 44(2):876 – 905, 2016.
- J. Zavatone-Veth and C. Pehlevan. Exact marginal prior distributions of finite Bayesian neural networks. In *Advances in Neural Information Processing Systems*, volume 34, pages 3364–3375, 2021.

Supplementary material

This supplement is structured along three Sections. Section B contains the remaining proofs of the results stated in the main paper, along with, at the end of the Section, a few remarks on allowing slightly different parameter choices. Section C starts with a few comments on how to extend the results to other statistical models. It then also contains the statement and proof of a few additional results: the case of Sobolev truths; the proof of the lower bound of Theorem 2 in the special (and easier) case $p = 1$; an example of statement of an upper-bound rates for the classical posterior (case $\rho = 1$, in contrast to the case $\rho < 1$ considered in the main paper); an extension of Theorem 4 to the case of light tails $p > 1$. Finally, Section D gathers a few technical lemmas used along the proofs.

Appendix B: Remaining proofs

B.1. End of the proof of Theorem 1

Let us now focus on the case $\alpha \leq \beta$. The constants C_i , $i = 1, \dots, 6$ might change from the previous case. Again, we split along indices k , this time separating $1 \leq k \leq N_\alpha$ and $k > N_\alpha$. Noticing that $N_\alpha/n \leq \varepsilon_n^2 = n^{2\alpha/(2\alpha+1)}$, we use the inclusion

$$\bigcap_{k=1}^{N_\alpha} \left\{ f : (f_k - f_{0,k})^2 \leq \frac{D^2}{2n} \right\} \cap \left\{ f : \sum_{k>N_\alpha} (f_k - f_{0,k})^2 \leq \frac{(D\varepsilon_n)^2}{2} \right\} \subset \left\{ f : \|f - f_0\|_2^2 \leq (D\varepsilon_n)^2 \right\}.$$

First consider the case of indices $1 \leq k \leq N_\alpha$. One reproduces the same argument as for the case $\alpha > \beta$ above: setting $\delta_k := D/\sqrt{2n}$,

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \frac{2c_0\delta_k}{\sigma_k} \exp \left\{ -\frac{c_1}{\sigma_k^p} \kappa_p(f_{0,k}^p + n^{-p/2}) \right\} \geq \frac{2c_0\delta_k}{\sigma_k} \exp \left\{ -c_1 \kappa_p(L^p + \sigma_k^{-p} n^{-p/2}) \right\},$$

where we use that $|f_{0,k}| \leq Lk^{-1/2-\beta} \leq Lk^{-1/2-\alpha} = L\sigma_k$ for $\alpha \leq \beta$. Using further that $\sigma_k^{-1} \leq \sqrt{n}$ for $k \leq N_\alpha$, one obtains, with $\sigma_k = k^{-1/2-\alpha}$ by definition,

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \geq C_2 D \frac{k^{1/2+\alpha}}{\sqrt{n}} \exp\{-C_3\}.$$

Taking the product over $1 \leq k \leq N_\alpha$ and using Lemma 6 leads to

$$\prod_{k=1}^{N_\alpha} \Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \exp\{-(1/2 + \alpha)N_\alpha - (C_4 + \log D)N_\alpha\} \geq \exp\{-C_5 n \varepsilon_n^2\},$$

where $C_5 > 0$ is a large enough constant and we used $N_\alpha \lesssim n \varepsilon_n^2$ in the last inequality.

Second let us deal with indices $k > N_\alpha$. One has $\sum_{k>N_\alpha} f_{0,k}^2 \leq L^2 N_\alpha^{-2\alpha} \lesssim \varepsilon_n^2$. So for large enough D , it holds

$$\Pi \left[\sum_{k>N_\alpha} (f_k - f_{0,k})^2 \leq \frac{(D\varepsilon_n)^2}{2} \right] \geq \Pi \left[\sum_{k>N_\alpha} f_k^2 \leq \frac{(D\varepsilon_n)^2}{4} \right] \geq \Pi \left[\sum_{k>N_\alpha} (f_k^2 - \sigma_k^2 E[\zeta_k^2]) \leq \frac{(D\varepsilon_n)^2}{8} \right],$$

where for the last inequality we have used that $E[\zeta_k^2] = E[\zeta_1^2]$ is a fixed constant and $\sum_{k>N_\alpha} \sigma_k^2 \lesssim \varepsilon_n^2$ by definition of σ_k and N_α and once again taking D large enough. Looking now at the complement, Markov's inequality gives

$$\begin{aligned} \Pi \left[\sum_{k>N_\alpha} \left(f_k^2 - \sigma_k^2 E[\zeta_k^2] \right) > (D\varepsilon_n)^2/8 \right] &= P \left[\sum_{k>N_\alpha} \sigma_k^2 \left(\zeta_k^2 - E[\zeta_k^2] \right) > (D\varepsilon_n)^2/8 \right] \\ &\leq \frac{64}{(D\varepsilon_n)^4} \text{Var} \left[\sum_{k>N_\alpha} \sigma_k^2 \zeta_k^2 \right] \leq \frac{64}{(D\varepsilon_n)^4} \text{Var} [\zeta_1^2] \sum_{k>N_\alpha} \sigma_k^4 \leq \frac{C_7}{(D\varepsilon_n)^4} N_\alpha^{-1-4\alpha}. \end{aligned}$$

Since $N_\alpha^{-1-4\alpha} \lesssim N_\alpha^{-1} \varepsilon_n^4$, one obtains that the prior mass in the last display goes to 0 as $n \rightarrow \infty$. In particular, the last but one display is bounded from below by $1/2$ for large enough n . Putting together the above bounds in both regimes of k 's leads to

$$\Pi [\|f - f_0\|_2 \leq D\varepsilon_n] \geq \frac{1}{2} \exp \left\{ -C_5 n \varepsilon_n^2 \right\} \geq \exp \left\{ -C n \varepsilon_n^2 \right\},$$

which concludes the proof of the theorem.

B.2. Proof of Theorem 2

Proof. To simplify the notation, we give the proof first for the standard posterior $\rho = 1$. We also focus on the (harder) case $p < 1$: the proof for $p = 1$ is similar, though easier (the maximiser μ_k of the function h_k below is completely explicit in that case): for completeness we give it explicitly below, see Section C.3.

Let us choose f_0 as the function in $\mathcal{F}^\beta(L)$ defined through its basis coefficients by $f_{0,k} = Lk^{-1/2-\beta}$ for $\beta > 0, L > 0$. Let us recall the definitions of γ in (12) and $N_\gamma := \lfloor n^{1/(2\gamma+1)} \rfloor$, as well as the target rate $\varepsilon_n = \varepsilon_n(p, \alpha, \beta) = N_\gamma^{-\beta}$. Denoting $\|g\|_{N_\gamma}^2 := \sum_{k=1}^{N_\gamma} g_k^2$ for any square-integrable function g , it is enough to prove, for small $m > 0$ to be chosen and $n_\gamma = dN_\gamma$ for some small enough constant d to be chosen below, that, as $n \rightarrow \infty$,

$$E_{f_0} \Pi[\|f - f_0\|_{n_\gamma} \geq m\varepsilon_n \mid X] \rightarrow 1.$$

For a sequence $\mu = (\mu_k)$ to be defined below (that will correspond to the mode of the posterior over the corresponding coordinates), the triangle inequality gives $\|f_0 - \mu\|_{n_\gamma} \leq \|f_0 - f\|_{n_\gamma} + \|f - \mu\|_{n_\gamma}$. This implies

$$\Pi[\|f - f_0\|_{n_\gamma} \geq m\varepsilon_n \mid X] \geq \Pi[\|f - \mu\|_{n_\gamma} \leq m\varepsilon_n \mid X] \cdot \mathbb{1}\{\|f_0 - \mu\|_{n_\gamma} \geq 2m\varepsilon_n\}. \quad (33)$$

It now suffices to show that each term of the product of the right hand side of the last display goes to 1 in probability under $P_{f_0}^{(n)}$.

We start by showing $\Pi[\|f - \mu\|_{n_\gamma} \leq m\varepsilon_n \mid X]$ goes to 1 in probability under $P_{f_0}^{(n)}$. Consider the event

$$\mathcal{B}_n := \left\{ |\xi_k| \leq \sqrt{2 \log n}, \quad \text{for all } k = 1, \dots, n_\gamma \right\}.$$

Since $P_{f_0}^{(n)}(\mathcal{B}_n) \rightarrow 1$ (Lemma 8), it is sufficient to work on the event $\mathcal{B}_n$ defined above and show $E_{f_0} \Pi[\|f - \mu\|_{n_\gamma} > m\varepsilon_n \mid X] \mathbf{1}_{\mathcal{B}_n} \rightarrow 0$. Below are the steps to obtain such result.

Expression of the posterior. Since $f_k = \sigma_k \zeta_k$, where ζ_k is p -exp, one can express the posterior on the coefficient f_k as

$$d\Pi_k(\theta \mid X^n) = \frac{e^{-\frac{n}{2}(X_k - \theta)^2 - \frac{|\theta|^p}{p\sigma_k^p}} d\theta}{\int e^{-\frac{n}{2}(X_k - \theta)^2 - \frac{|\theta|^p}{p\sigma_k^p}} d\theta}.$$

Let use denote $\tilde{g}_k^+(\theta) := e^{-\frac{n}{2}(X_k - \theta)^2 - \frac{|\theta|^p}{p\sigma_k^p}} \mathbf{1}_{\theta \geq 0}$ and $\tilde{g}_k^-(\theta) := e^{-\frac{n}{2}(X_k - \theta)^2 - \frac{|\theta|^p}{p\sigma_k^p}} \mathbf{1}_{\theta < 0}$, as well as

$$G_k^+ := \int \tilde{g}_k^+(\theta) d\theta, \quad G_k^- := \int \tilde{g}_k^-(\theta) d\theta \quad \text{and} \quad w_k^+ := \frac{G_k^+}{G_k^+ + G_k^-}.$$

Re-normalizing $g_k^+ := \tilde{g}_k^+ / G_k^+$ and $g_k^- := \tilde{g}_k^- / G_k^-$, one obtains the decomposition

$$d\Pi_k(\theta \mid X^n) = w_k^+ g_k^+(\theta) d\theta + (1 - w_k^+) g_k^-(\theta) d\theta.$$

By Markov's inequality, for any $\mu = (\mu_k)$ to be chosen below,

$$\begin{aligned} \Pi[\|f - \mu\|_{n_\gamma} > m\varepsilon_n \mid X] \mathbf{1}_{\mathcal{B}_n} &\leq \frac{1}{(m\varepsilon_n)^2} \int \|f - \mu\|_{n_\gamma}^2 d\Pi(f \mid X) \mathbf{1}_{\mathcal{B}_n} \\ &\leq \frac{\mathbf{1}_{\mathcal{B}_n}}{(m\varepsilon_n)^2} \sum_{k=1}^{n_\gamma} \left[w_k^+ \int (f_k - \mu_k)^2 g_k^+(f_k) df_k + (1 - w_k^+) \int (f_k - \mu_k)^2 g_k^-(f_k) df_k \right]. \end{aligned} \quad (34)$$

Control on the event $\mathcal{B}_n$. In order to control the various terms that will appear below, we set a constant $M = M(p)$, such that

$$M(p) \geq 10^{2-p} \vee \left(1 + \frac{2^p}{2p(1-p)} \right)^{2-p} \vee \frac{8}{p} \left(\frac{3}{2} \right)^p. \quad (35)$$

Applying Lemma 8 shows that we can take $d > 0$ small enough in $n_\gamma = dN_\gamma$, such that, for all $k \leq n_\gamma$ and n large enough, on $\mathcal{B}_n$ and under $P_{f_0}^{(n)}$, we have

$$nX_k^2 \geq M \left(\frac{X_k}{\sigma_k} \right)^p. \quad (36)$$

Also note that, thanks to Lemma 8, for all $k \leq n_\gamma$, on $\mathcal{B}_n$, we have

$$X_k \geq f_{0,k}/2 > 0.$$

Laplace-type bound on the positive part, choice of μ_k . Since $X_k > 0$, provided we choose $\mu_k > 0$ on $\mathcal{B}_n$, we will see below that most of the posterior mass will come from the ‘positive part’. Since $w_k^+ \leq 1$, in order to control the term involving g_k^+ in (34), it suffices to bound from above, on the event

$\mathcal{B}_n$,

$$\sum_{k \leq n_\gamma} \int (\theta - \mu_k)^2 g_k^+(\theta) d\theta.$$

Let us define




Figure 4. Function h_k with $p = 0.1$, $X_k = \sigma_k = 1$ and $n = 80 > 10^{2-p}$

$$h_k(\theta) := -\frac{n}{2}(X_k - \theta)^2 - \frac{\theta^p}{p\sigma_k^p}, \quad k = 1, \dots, n_\gamma \quad (37)$$

for $\theta > 0$. Simple algebraic computation show that $h_k''' < 0$ and

$$h_k'(\theta) = n(X_k - \theta) - \frac{\theta^{p-1}}{\sigma_k^p}, \quad (38)$$

$$h_k''(\theta) = -n + (1-p)\frac{\theta^{p-2}}{\sigma_k^p}. \quad (39)$$

Hence, as $\theta > 0$ grows, h_k'' strictly decreases from $+\infty$ to $-n$ and vanishes at a single inflection point θ_2^* , such that $h_k''(\theta_2^*) = 0$, where

$$\theta_2^* := \left(\frac{n\sigma_k^p}{1-p} \right)^{\frac{1}{p-2}}. \quad (40)$$

Denote $\psi(u) := 4u^{1/(1+u)} + 4^{-u}u^{-u/(1+u)}$, for $u \in (0, 1)$. Using $u \log \frac{1}{u} \leq 1 + u$, available for all $u \in (0, 1)$ we obtain $\psi(u) \leq 4 + e < 7$. We can then compute

$$\begin{aligned} h'_k(\theta_2^*) &\geq h'_k(4\theta_2^*) = n \left(X_k - 4 \left(\frac{n\sigma_k^p}{1-p} \right)^{\frac{1}{p-2}} - \frac{4^{p-1}}{n\sigma_k^p} \left(\frac{n\sigma_k^p}{1-p} \right)^{\frac{p-1}{p-2}} \right) \\ &= n \left(X_k - (n\sigma_k^p)^{\frac{1}{p-2}} \psi(1-p) \right) \\ &\geq n(X_k - 7(n\sigma_k^p)^{\frac{1}{p-2}}). \end{aligned}$$

Since $M > 7^{2-p}$ in (35) and thanks to our choice of $d > 0$, by (36) we have $nX_k^2 > 7^{2-p}(X_k/\sigma_k)^p$ and thus $h'_k(\theta_2^*) \geq h'_k(4\theta_2^*) > 0$.

Thus, h'_k starts from $-\infty$ at 0, ends at $-\infty$ at $+\infty$ and takes a positive value at θ_2^* , therefore h'_k vanishes at exactly two critical points θ_m^* and θ_M^* , such that $\theta_m^* < \theta_2^* < \theta_M^*$ (these critical points obviously depend on k ; we do not make this explicit to avoid overloading the notation). In particular, at θ_M^* there is a local maximum of h_k (because $\theta_M^* > \theta_2^*$ implies $h''_k(\theta_M^*) < h''_k(\theta_2^*) = 0$).

First notice that $\lim_{\theta \rightarrow \infty} h_k(\theta) = -\infty$. Second, recalling (40) the definition of θ_2^* , we have $h_k(2\theta_2^*) > h_k(0)$, this shows that the global maximum of h_k is attained at θ_M^* and not at the boundary. Indeed

$$\begin{aligned} h_k(2\theta_2^*) - h_k(0) &= 2nX_k\theta_2^* - 2n(\theta_2^*)^2 - \frac{2^p}{p\sigma_k^p}(\theta_2^*)^p \\ &= 2n\theta_2^* \left(X_k - \left(1 + \frac{2^p}{2p(1-p)}\right)\theta_2^* \right) > 0, \end{aligned}$$

where the last inequality comes from choice of M large enough in (35).

We finally set, for any $k \leq n_\gamma$, μ_k as the global maximizer of h_k ,

$$\mu_k := \theta_M^*.$$

We apply now a Laplace's method-type argument, to upper bound, on the event $\mathcal{B}_n$,

$$\int (\theta - \mu_k)^2 g_k^+(\theta) d\theta := \frac{\int_0^\infty (\theta - \theta_M^*)^2 e^{h_k(\theta)} d\theta}{\int_0^\infty e^{h_k(\theta)} d\theta}.$$

Taylor's formula at $\theta_M^* = \mu_k$, provides $\xi_\theta \in [\theta \wedge \theta_M^*, \theta \vee \theta_M^*]$, such that

$$h_k(\theta) = h_k(\theta_M^*) + \frac{1}{2}(\theta - \theta_M^*)^2 h''_k(\xi_\theta). \quad (41)$$

Recall (39) the expression of h''_k . Plugging the simple bound $h''_k \geq -n$ in (41), allows us to bound the denominator

$$\int_0^\infty e^{h_k(\theta)} d\theta \geq e^{h_k(\theta_M^*)} \int_0^\infty e^{-\frac{n}{2}(\theta - \theta_M^*)^2} d\theta \geq \sqrt{\frac{\pi}{2n}} e^{h_k(\theta_M^*)}.$$

For the numerator, recall (40) the definition of θ_2^* and recall that we showed $h'_k(4\theta_2^*) > 0$, thus $2\theta_2^* \leq 4\theta_2^* \leq \theta_M^*$. We cut the integral at $2\theta_2^*$ and obtain

$$\int_0^\infty (\theta - \theta_M^*)^2 e^{h_k(\theta)} d\theta = \underbrace{\int_0^{2\theta_2^*} (\theta - \theta_M^*)^2 e^{h_k(\theta)} d\theta}_{I_1} + \underbrace{\int_{2\theta_2^*}^\infty (\theta - \theta_M^*)^2 e^{h_k(\theta)} d\theta}_{I_2}.$$

Bounding I_2 first, for $\theta \in [2\theta_2^*, \infty)$, we have $\xi_\theta \geq 2\theta_2^*$, hence $h''_k(\xi_\theta) \leq h''_k(2\theta_2^*) = -n(1 - 2^{p-2}) \leq -n/2$ and thus

$$I_2 \leq e^{h_k(\theta_M^*)} \int_{2\theta_2^*}^\infty (\theta - \theta_M^*)^2 e^{-\frac{n}{4}(\theta - \theta_M^*)^2} d\theta \leq e^{h_k(\theta_M^*)} \frac{4\sqrt{\pi}}{n\sqrt{n}}.$$

Now for I_1 , we know h_k decreases from 0 to θ_m^* and then increases up until $\theta_M^* \geq 2\theta_2^*$, thus

$$I_1 \leq (e^{h_k(0)} \vee e^{h_k(2\theta_2^*)}) \int_0^{2\theta_2^*} (\theta - \theta_M^*)^2 d\theta \leq 4\theta_2^*[(\theta_2^*)^2 + (\theta_M^*)^2](e^{h_k(0)} \vee e^{h_k(2\theta_2^*)}).$$

Considering the maximum on the right hand side, we showed in the study of h_k above, that, on the event $\mathcal{B}_n$, we have $h_k(2\theta_2^*) > h_k(0)$. Along with $\theta_M^* \leq X_k$ (from the critical equation $h'_k(\theta_M^*) = 0$ in (38)), this provides,

$$I_1 \leq 4\theta_2^*[(\theta_2^*)^2 + X_k^2]e^{h_k(2\theta_2^*)}.$$

Along with the denominator bound, we obtain

$$\frac{\int_0^\infty (\theta - \theta_M^*)^2 e^{h_k(\theta)} d\theta}{\int_0^\infty e^{h_k(\theta)} d\theta} \lesssim \sqrt{n}\theta_2^*[(\theta_2^*)^2 + X_k^2]e^{h_k(2\theta_2^*) - h_k(\theta_M^*)} + \frac{1}{n}.$$

We showed above that $4\theta_2^* \leq \theta_M^*$, so that $2\theta_2^* - \theta_M^* \leq -\theta_M^*/2$. Combined with an application of the mean-value theorem, and noticing that h'_k is decreasing in $[2\theta_2^*, \theta_M^*]$ (since it vanishes at θ_m^* and θ_M^* , while $h'_k(\theta_2^*) > 0$, where $\theta_m^* < 2\theta_2^* < \theta_M^*$), we have

$$h_k(2\theta_2^*) - h_k(\theta_M^*) \leq h'_k(2\theta_2^*)[2\theta_2^* - \theta_M^*] \leq -\frac{1}{2}\theta_M^* h'_k(2\theta_2^*).$$

Doing similar computations as for the bound of $h'_k(4\theta_2^*)$ above and using $nX_k^2 \geq 10^{2-p}(X_k/\sigma_k)^p$ (again $M \geq 10^{2-p}$ in (35)), we have (from $2 + e \leq 5$)

$$h'_k(2\theta_2^*) \geq n(X_k - 5(n\sigma_k^p)^{\frac{1}{p-2}}) \geq nX_k/2.$$

These inequalities combined provide $e^{h_k(2\theta_2^*) - h_k(\theta_M^*)} \leq e^{-\frac{1}{4}nX_k\theta_M^*}$ and thus

$$\frac{\int_0^\infty (\theta - \theta_M^*)^2 e^{h_k(\theta)} d\theta}{\int_0^\infty e^{h_k(\theta)} d\theta} \lesssim \sqrt{n}\theta_2^*[(\theta_2^*)^2 + X_k^2]e^{-\frac{1}{4}nX_k\theta_M^*} + \frac{1}{n}.$$

Recall $\theta_2^* \leq \theta_M^*$. Using the definition of θ_2^* in (40), the fact that $h'_k(\theta_M^*) = 0$ where h'_k is given in (38), as well as $(1-p)^{-\frac{1-p}{2-p}} \leq 2$, we obtain

$$\theta_M^* = X_k - \frac{(\theta_M^*)^{p-1}}{n\sigma_k^p} \geq X_k - (1-p)^{-\frac{1-p}{2-p}} (n\sigma_k^p)^{\frac{p-1}{p-2}-1} \geq X_k - 2(n\sigma_k^p)^{\frac{1}{p-2}}. \quad (42)$$

Since $M \geq 4^{2-p}$ in (35), we have $nX_k^2 \geq 4^{2-p}(X_k/\sigma_k)^p$ and the last display is further lower bounded from below by $X_k/2$, so that

$$\frac{\int_0^\infty (\theta - \theta_M^*)^2 e^{h_k(\theta)} d\theta}{\int_0^\infty e^{h_k(\theta)} d\theta} \lesssim \sqrt{n} \theta_2^* [(\theta_2^*)^2 + X_k^2] e^{-\frac{1}{8}nX_k^2} + \frac{1}{n}.$$

Summing up over $k \leq n_\gamma$ gives two terms on the right hand side. The first term is smaller than any polynomial power of n . Indeed, thanks to Lemma 8, on the event $\mathcal{B}_n$, we have $f_{0,k}/2 \leq X_k \leq 3f_{0,k}/2$. There exist constants c_1, c_2 , such that, recalling $\alpha > \beta$,

$$\sqrt{n} \sum_{k \leq n_\gamma} \theta_2^* [(\theta_2^*)^2 + X_k^2] e^{-\frac{1}{8}nX_k^2} \lesssim \sqrt{n} e^{-c_2 n n_\gamma^{-2\beta-1}} \sum_{k \leq n_\gamma} (n\sigma_k^p)^{\frac{1}{p-2}} [(n\sigma_k^p)^{\frac{2}{p-2}} + f_{0,k}^2] \lesssim n^{c_1} e^{-c_2 n^{\frac{p(\alpha-\beta)}{1+2\beta+p(\alpha-\beta)}}}.$$

The last displayed bound is $o(n_\gamma^{-2\beta})$ as $n \rightarrow \infty$. For the third term, simply notice that $n_\gamma/n = o(n_\gamma^{-2\beta})$, so that it holds

$$E_{f_0} \sum_{k \leq n_\gamma} w_k^+ \int (\theta - \mu_k)^2 g_k^+(\theta) d\theta \mathbf{1}_{\mathcal{B}_n} = o(n_\gamma^{-2\beta}). \quad (43)$$

Control on the weight of the negative part. We are only left to control

$$(1 - w_k^+) \int (f_k - \mu_k)^2 g_k^-(f_k) df_k,$$

on the event $\mathcal{B}_n$. We start by bounding the weight $1 - w_k^+ = (1 + G_k^+/G_k^-)^{-1} \leq G_k^-/G_k^+$. For $\theta < 0$ and $X_k \geq 0$, we have $(X_k - \theta)^2 \geq X_k^2$, so that

$$G_k^- = \int_{-\infty}^0 e^{-\frac{n}{2}(X_k - \theta)^2 - \frac{|\theta|^p}{p\sigma_k^p}} d\theta \leq e^{-\frac{nX_k^2}{2}} \int_0^\infty e^{-\frac{\theta^p}{p\sigma_k^p}} d\theta.$$

The change of variable $\theta = \sigma_k(pu)^{1/p}$ yields

$$G_k^- \leq p^{\frac{1}{p}-1} \Gamma(1/p) \sigma_k e^{-\frac{nX_k^2}{2}}.$$

For G_k^+ , restricting the integral on $[X_k/2, 3X_k/2]$, so that $(X_k - \theta)^2 \leq X_k^2/4$, provides

$$G_k^+ \geq \int_{X_k/2}^{3X_k/2} e^{-\frac{n}{2}(X_k - \theta)^2 - \frac{\theta^p}{p\sigma_k^p}} d\theta \geq e^{-\frac{nX_k^2}{8}} \int_{X_k/2}^{3X_k/2} e^{-\frac{\theta^p}{p\sigma_k^p}} d\theta \geq X_k e^{-\frac{nX_k^2}{8}} e^{-\frac{1}{p}(\frac{3}{2})^p \left(\frac{X_k}{\sigma_k}\right)^p}.$$

Combining, we obtain

$$1 - w_k^+ \lesssim \frac{\sigma_k}{X_k} \exp \left\{ -\frac{3}{8} n X_k^2 + \frac{1}{p} \left(\frac{3}{2} \right)^p \left(\frac{X_k}{\sigma_k} \right)^p \right\},$$

where the constant depends only on p . We now employ Lemma 8, on the event $\mathcal{B}_n$ and thanks to our choice of M in (35), we have for all $k \leq n_\gamma$ and large enough n ,

$$\frac{L}{2} \sigma_k \leq \frac{f_{0,k}}{2} \leq X_k \quad \text{and} \quad n X_k^2 \geq \frac{8}{p} \left(\frac{3}{2} \right)^p \frac{X_k^p}{\sigma_k^p}.$$

This leads to the bound, on the event $\mathcal{B}_n$ and under $P_{f_0}^{(n)}$,

$$1 - w_k^+ \lesssim \exp \left\{ -\frac{1}{16} n f_{0,k}^2 \right\}. \quad (44)$$

We are left to control

$$\int (\theta - \mu_k)^2 g_k^-(\theta) d\theta = \frac{\int_0^\infty (\theta + \theta_M^*)^2 e^{-\frac{n}{2}(X_k + \theta)^2 - \frac{\theta^p}{p\sigma_k^p}} d\theta}{\int_0^\infty e^{-\frac{n}{2}(X_k + \theta)^2 - \frac{\theta^p}{p\sigma_k^p}} d\theta}.$$

First use $(\theta + \theta_M^*)^2 \leq 2\theta^2 + 2(\theta_M^*)^2$ and upper bound similarly as before

$$\int_0^\infty \theta^2 e^{-\frac{n}{2}(X_k + \theta)^2 - \frac{\theta^p}{p\sigma_k^p}} d\theta \leq e^{-\frac{nX_k^2}{2}} \int_0^\infty \theta^2 e^{-\frac{\theta^p}{p\sigma_k^p}} d\theta \lesssim e^{-\frac{nX_k^2}{2}} p^{\frac{3}{p}-1} \Gamma(3/p) \sigma_k^3.$$

For the denominator, restricting the integral up to $c\sigma_k$, where $c > 0$ is a small enough constant to be chosen below, we obtain

$$\int_0^\infty e^{-\frac{n}{2}(X_k + \theta)^2 - \frac{\theta^p}{p\sigma_k^p}} d\theta \geq e^{-\frac{c^p}{p}} \int_0^{c\sigma_k} e^{-\frac{n}{2}(X_k + \theta)^2} d\theta \geq c\sigma_k e^{-\frac{c^p}{p}} e^{-\frac{n}{2}(X_k + c\sigma_k)^2}.$$

Finally, combining the two previous bounds yields

$$\int (\theta - \mu_k)^2 g_k^-(\theta) d\theta \lesssim (\theta_M^*)^2 + \sigma_k^2 e^{-\frac{n}{2}X_k^2 + \frac{n}{2}(X_k + c\sigma_k)^2}.$$

By Lemma 8, on the event $\mathcal{B}_n$, we have $L\sigma_k \leq 2X_k$ and $2X_k \leq 3f_{0,k}$, thus

$$\int (\theta - \mu_k)^2 g_k^-(\theta) d\theta \lesssim (\theta_M^*)^2 + \sigma_k^2 e^{2\frac{c}{L}(1+\frac{c}{L})nX_k^2} \lesssim (\theta_M^*)^2 + \sigma_k^2 e^{\frac{9}{2}\frac{c}{L}(1+\frac{c}{L})nf_{0,k}^2}.$$

Using the bound (44) above, we finally have

$$\sum_{k \leq n_\gamma} (1 - w_k^+) \int (\theta - \mu_k)^2 g_k^-(\theta) d\theta \lesssim \sum_{k \leq n_\gamma} ((\theta_M^*)^2 + \sigma_k^2 e^{\frac{9}{2}\frac{c}{L}(1+\frac{c}{L})nf_{0,k}^2}) e^{-\frac{1}{16}nf_{0,k}^2}.$$

Using $\theta_M^* = X_k - (\theta_M^*)^{p-1}/n\sigma_k^p \leq X_k \leq f_{0,k}/2$ and $L\sigma_k \leq 2X_k$, this bound becomes

$$\sum_{k \leq n_\gamma} (1 - w_k^+) \int (\theta - \mu_k)^2 g_k^-(\theta) d\theta \lesssim \sum_{k \leq n_\gamma} f_{0,k}^2 (1 + e^{\frac{9}{2}\frac{c}{L}(1+\frac{c}{L})nf_{0,k}^2}) e^{-\frac{1}{16}nf_{0,k}^2}.$$

We choose $c > 0$ so that $\frac{9}{2} \frac{c}{L} (1 + \frac{c}{L}) = \frac{1}{32}$ and we obtain, on $\mathcal{B}_n$, recalling $\alpha > \beta$ (hence $\gamma > \beta$)

$$\sum_{k \leq n_\gamma} (1 - w_k^+) \int (\theta - \mu_k)^2 g_k^-(\theta) d\theta \lesssim \sum_{k \leq n_\gamma} f_{0,k}^2 e^{-\frac{1}{32} n f_{0,k}^2} = o(n^{-1}) = o(\varepsilon_n^2). \quad (45)$$

Conclusion of the Proof. We gather the bounds (43) and (45) into equation (34) and finally obtain

$$E_{f_0} \Pi[\|f - \mu\|_{n_\gamma} > m\varepsilon_n \mid X] \mathbf{1}_{\mathcal{B}_n} = o(1).$$

We are only left to show that the indicator $\mathbf{1}\{\|f_0 - \mu\|_{n_\gamma} \geq 2m\varepsilon_n\}$ in (33) goes to 1 under $P_{f_0}^{(n)}$. Taking the expectation and using $\|\mu - f_0\|_{n_\gamma} \geq \|\mu - X\|_{n_\gamma} - \|X - f_0\|_{n_\gamma}$ by the triangle inequality,

$$P_{f_0}^{(n)}[\|\mu - f_0\|_{n_\gamma} \geq 2m\varepsilon_n] \geq P_{f_0}^{(n)}[\|X - f_0\|_{n_\gamma} \leq m\varepsilon_n, \|\mu - X\|_{n_\gamma} \geq 3m\varepsilon_n].$$

Recall the definition of the event $\mathcal{B}_n$, which satisfies $P_{f_0}^{(n)}(\mathcal{B}_n) \rightarrow 1$ (Lemma 8),

$$\mathcal{B}_n := \left\{ |\xi_k| \leq \sqrt{2 \log n}, \quad \text{for all } k = 1, \dots, n_\gamma \right\}.$$

On this event and under $P_{f_0}^{(n)}$, using $\alpha > \beta$, we have for n large enough

$$\|X - f_0\|_{n_\gamma}^2 = \sum_{k \leq n_\gamma} \frac{1}{n} |\xi_k|^2 \leq \frac{n_\gamma}{n} \times 2 \log n = o(\varepsilon_n^2).$$

Second, recall that $\mu_k = \theta_M^*$, defined by $n(X_k - \theta_M^*) = \sigma_k^{-p} (\theta_M^*)^{p-1}$, satisfying $\theta_M^* \leq X_k$ so that, since $p < 1$,

$$\|\mu - X\|_{n_\gamma}^2 = \sum_{k \leq n_\gamma} \frac{\sigma_k^{-2p}}{n^2} |\theta_M^*|^{2(p-1)} \geq \sum_{k \leq n_\gamma} \frac{\sigma_k^{-2p}}{n^2} |X_k|^{2(p-1)}.$$

On the event $\mathcal{B}_n$, thanks to Lemma 8, we have $|X_k| \asymp |f_{0,k}|$ and thus

$$\|\mu - X\|_{n_\gamma}^2 \gtrsim \sum_{k \leq n_\gamma} \frac{\sigma_k^{-2p}}{n^2} |f_{0,k}|^{2(p-1)} \gtrsim \frac{n_\gamma^{2(1+\beta+p(\alpha-\beta))}}{n^2} \gtrsim \varepsilon_n^2.$$

Combining the two previous facts shows that for a small enough constant m , for n large enough,

$$P_{f_0}^{(n)}[\|X - f_0\|_{n_\gamma} \leq m\varepsilon_n, \|\mu - X\|_{n_\gamma} \geq 3m\varepsilon_n] \geq P_{f_0}^{(n)}(\mathcal{B}_n),$$

where $P_{f_0}(\mathcal{B}_n) \rightarrow 1$. We have thus shown that the indicator in (33) goes to 1 in E_{f_0} -expectation, hence also in $P_{f_0}^{(n)}$ -probability, which concludes the proof. $\square$

B.3. Proof of Theorem 3

Proof. Recall $N_\beta = \lfloor n^{1/(2\beta+1)} \rfloor$ is the usual frequency cutoff. Using independence of the f_k drawn from the prior, it is sufficient to show

$$\prod_{k \geq 1} \Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \exp(-n\varepsilon_n^2),$$

where

$$\delta_k := \begin{cases} 1/\sqrt{n}, & 1 \leq k \leq N_\beta, \\ 2Lk^{-1/2-\beta}, & k > N_\beta \end{cases}. \quad (46)$$

Indeed, recalling $\varepsilon_n = n^{-\beta/(2\beta+1)} \log^{\eta'} n = N_\beta^{-\beta} \log^{\eta'} n = \sqrt{N_\beta/n} \log^{\eta'} n$, we have, for n large enough,

$$\sum_k \delta_k^2 = \sum_{k \leq N_\beta} \frac{1}{n} + 4L^2 \sum_{k \geq N_\beta} k^{-1-2\beta} \leq \frac{N_\beta}{n} + 4L^2 N_\beta^{-2\beta} \leq \varepsilon_n^2.$$

The following inclusions hold,

$$\{f : \forall k \geq 1, |f_k - f_{0,k}| \leq \delta_k\} \subset \left\{f : \sum_{k \geq 1} |f_k - f_{0,k}|^2 \leq \sum_{k \geq 1} \delta_k^2\right\} \subset \{f : \|f - f_0\|_2 \leq \varepsilon_n\},$$

establishing the claimed sufficiency of the condition above. Let us start with the indices $k \leq N_\beta$. We bound from below

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] = P\left[|\sigma_k \zeta_k - f_{0,k}| \leq \frac{1}{\sqrt{n}}\right] = \int_{(f_{0,k}-1/\sqrt{n})/\sigma_k}^{(f_{0,k}+1/\sqrt{n})/\sigma_k} h_{p_k}(x) dx,$$

for all $k \geq 1$. Since h_{p_k} the density of ζ_k is symmetric (see (3)), we can assume without loss of generality that $f_{0,k} \geq 0$ in the bounds to follow. We can then bound the previous integral using the smallest value of h_{p_k} , which leaves us with

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \frac{2}{\sigma_k \sqrt{n}} h_{p_k}\left(\frac{f_{0,k} + 1/\sqrt{n}}{\sigma_k}\right).$$

Using the lower bound on h_p given in Lemma 9, we obtain

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \gtrsim \frac{U_k}{\sqrt{n}} \exp\left\{-\frac{1}{p_k} \left(\frac{f_{0,k} + 1/\sqrt{n}}{\sigma_k}\right)^{p_k}\right\}, \quad (47)$$

where

$$U_k := \sqrt{p_k} \exp\left(\frac{1}{p_k} - \frac{p_k}{12}\right) \sigma_k^{-1}.$$

Using first the smoothness condition on f_0 , we have $f_{0,k} \leq Lk^{-1/2-\beta}$ and then using $k \leq N_\beta$ we obtain $k^{-1/2-\beta} \geq n^{-\frac{1/2+\beta}{2\beta+1}} = 1/\sqrt{n}$. Since $p_k \in (0, 1]$, we obtain

$$(f_{0,k} + 1/\sqrt{n})^{p_k} \leq (L+1)^{p_k} k^{-p_k(1/2+\beta)} \leq (L+1) k^{-p_k(1/2+\beta)}.$$

Using $\sigma_k \leq 1$ and $p_k \leq 1$, we have $\sqrt{p_k} e^{\frac{1}{p_k} - \frac{p_k}{12}} \geq 1 \geq \sigma_k$ and thus $U_k \geq 1$. Finally plugging these two inequalities in (47), we obtain,

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \gtrsim \frac{1}{\sqrt{n}} \exp \left\{ -\frac{L+1}{p_k} \left(\frac{k^{-\beta-1/2}}{\sigma_k} \right)^{p_k} \right\} = \frac{1}{\sqrt{n}} \exp\{-(L+1)z_k\}.$$

From this bound, it follows

$$\prod_{k \leq N_\beta} \Pi[|f_k - f_{0,k}| \leq \delta_k] \gtrsim \exp \left\{ -\sum_{k \leq N_\beta} (\log \sqrt{n} + (L+1)z_k) \right\}.$$

Employing the summability condition (17) on z_k , we obtain $C > 0$ and $\eta > 1$, such that

$$\prod_{k \leq N_\beta} \Pi[|f_k - f_{0,k}| \leq \delta_k] \geq e^{-CN_\beta \log^\eta n} \geq e^{-n\varepsilon_n^2}.$$

For the other part of the product, indexed by $k > N_\beta$, we have $\delta_k = 2Lk^{-1/2-\beta}$, such that

$$[f_{0,k} - \delta_k, f_{0,k} + \delta_k] \supset [-Lk^{-1/2-\beta}, Lk^{-1/2-\beta}].$$

Recalling $r_k = \sigma_k^{-1} k^{-1/2-\beta}$, we can then bound the probability of interest as

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \Pi[|f_k| \leq Lk^{-1/2-\beta}] = P[|\zeta_k| \leq L\sigma_k^{-1} k^{-1/2-\beta}] = P[|\zeta_k| \leq Lr_k].$$

Recalling that ζ_k has symmetric density h_{p_k} and survival function $\bar{H}_{p_k}$, we get

$$\Pi[|f_k - f_{0,k}| \leq \delta_k] \geq 1 - 2\bar{H}_{p_k}(Lr_k).$$

Taking now the product over $k > N_\beta$, using $L \geq 1$ and the monotonicity of $\bar{H}_{p_k}$, we obtain

$$\prod_{k > N_\beta} \Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \exp \left\{ \sum_{k > N_\beta} \log(1 - 2\bar{H}_{p_k}(r_k)) \right\}.$$

Thanks to the condition (16), we have $r_k \geq 1$. Using Lemma 9, we get

$$\bar{H}_{p_k}(r_k) \lesssim \frac{e^{\frac{1}{p_k}}}{\sqrt{p_k}} r_k e^{-r_k^{p_k}/p_k} \lesssim \exp \left\{ \frac{1}{2} \log \frac{1}{p_k} + \frac{1}{p_k} + \log r_k - \frac{r_k^{p_k}}{p_k} \right\}.$$

Using first $\frac{1}{2} \log \frac{1}{p_k} \leq \frac{1}{p_k}$ and then $\log r_k = \frac{1}{p_k} \log r_k^{p_k} \leq \frac{1}{2p_k} r_k^{p_k}$, we get

$$\bar{H}_{p_k}(r_k) \lesssim \exp \left\{ \frac{2}{p_k} - \frac{r_k^{p_k}}{2p_k} \right\} \lesssim \exp \left\{ -\frac{r_k^{p_k}}{4p_k} \right\} = \exp \left\{ -\frac{z_k}{4} \right\},$$

where the assumption $r_k^{p_k} \geq 8$ was used in the last inequality. As the general term of a convergent series (thanks to condition (17)) the previous bound goes to zero. Thus, for n large enough and $k > N_\beta$, we

have $\overline{H}_{p_k}(r_k) \leq 1/4$. We then use the inequality $\log(1 - 2x) \geq -4x$, available for $x \leq 1/4$, to obtain

$$\prod_{k > N_\beta} \Pi[|f_k - f_{0,k}| \leq \delta_k] \geq \exp \left\{ -4 \sum_{k > N_\beta} \overline{H}_{p_k}(r_k) \right\} \geq \exp \left\{ -C \sum_{k > N_\beta} \exp \left\{ -\frac{z_k}{4} \right\} \right\}.$$

This last sum goes to 0 when $n \rightarrow \infty$, as the remainder of a converging series, such that

$$\prod_{k > N_\beta} \Pi[|f_k - f_{0,k}| \leq \delta_k] \geq e^{-n\varepsilon_n^2},$$

for any $n\varepsilon_n^2 \rightarrow \infty$. □

B.4. Proof of Corollary 1

Proof. Let us look at the first case where $\sigma_k = k^{-1/2-\alpha}$ and $p_k = \log \log k / \log k$ for $k \geq 3$. We check the compatibility condition (16), we compute, for all $k > N_\beta$,

$$r_k^{p_k} = \left(\frac{k^{-1/2-\beta}}{\sigma_k} \right)^{p_k} = \left(k^{\alpha-\beta} \right)^{\frac{\log \log k}{\log k}} = (\log k)^{\alpha-\beta}.$$

Since $\alpha > \beta$, we have $(\log k)^{\alpha-\beta} \rightarrow \infty$, so that $r_k^{p_k} \geq 8$ for $k > N_\beta$ and n large enough. Finally to check the summability conditions (17), we compute

$$z_k = \frac{r_k^{p_k}}{p_k} = \frac{(\log k)^{1+\alpha-\beta}}{\log \log k},$$

with the appropriate adaptations whenever $k = 1, 2$. Such that, taking $\eta = 1 + \alpha - \beta > 1$, we obtain

$$\sum_{k \leq N_\beta} z_k \leq C + \frac{(\log N_\beta)^{\alpha-\beta}}{\log \log 3} \sum_{k=3}^{N_\beta} \log k \lesssim N_\beta (\log N_\beta)^{1+\alpha-\beta} = N_\beta \log^\eta n.$$

We also have that

$$\exp\{-z_k/4\} = \exp \left\{ -\frac{1}{4} \frac{(\log k)^{1+\alpha-\beta}}{\log \log k} \right\}$$

is the general term of a summable series since $\alpha > \beta$.

Now looking at the second case where $\sigma_k = \exp\{-\log^{1+\gamma} k\}$ and $p_k = c / \log^{1+\gamma} k$. We check the compatibility condition (16), where

$$r_k^{p_k} = \left(\frac{k^{-1/2-\beta}}{\sigma_k} \right)^{p_k} = \left(k^{-1/2-\beta} e^{\log^{1+\gamma} k} \right)^{\frac{c}{\log^{1+\gamma} k}} = e^{-(1/2+\beta) \frac{c}{\log^\gamma k} + c}$$

which, provided $c > 2.1 > \log 8$, satisfies $r_k^{p_k} \geq 8$, for $k > N_\beta$ and n large enough. Finally, we compute

$$z_k = \frac{r_k^{p_k}}{p_k} = e^{-(1/2+\beta) \frac{c}{\log^\gamma k} + c} \log^{1+\gamma} k.$$

Therefore, taking $\eta = 1 + \gamma$, we have, for large enough n ,

$$\sum_{k=3}^{N_\beta} z_k \lesssim N_\beta e^{c(1 - \frac{1/2+\beta}{\log^\gamma N_\beta})} (\log N_\beta)^{1+\gamma} \lesssim N_\beta \log^\eta n.$$

To check the rest of Condition (17), notice that, for large enough k , we have $\tilde{c} > 0$, so that

$$\exp\{-z_k/4\} = \exp\left\{-\frac{1}{4c} e^{c(1 - \frac{1/2+\beta}{\log^\gamma k})} \log^{1+\gamma} k\right\} \leq \exp\left\{-\tilde{c} \log^{1+\gamma} k\right\},$$

which is the general term of a converging series. $\square$

B.5. Proof of Theorem 4 and Corollary 2

Proof. Corollary 2 is directly obtained by applying (the second part of) Lemma 5 to lower bound $D_\rho(P_f^n, P_{f_0}^n)/n$ in Theorem 4. We now prove Theorem 4. In the random design regression setting of (20), Lemma 4 and 5 show that it is sufficient to get, for some constant $C > 0$,

$$\Pi[\|f - f_0\|_\infty \leq \varepsilon_n] \geq e^{-Cn\varepsilon_n^2}. \quad (48)$$

Recall $f_{N_\beta}^*$ is the approximating network in (24) obtained by Lemma 10 and satisfies $\|f - f_{N_\beta}^*\|_\infty \leq 2LN_\beta^{-\beta}$. For the considered $\varepsilon_n \gtrsim N_\beta^{-\beta} = \varepsilon_n^*$, the triangle inequality implies, for some constant $d > 0$ small enough, that $\Pi[\|f - f_0\|_\infty \leq \varepsilon_n] \geq \Pi[\|f - f_{N_\beta}^*\|_\infty \leq d\varepsilon_n]$.

We now relabel the approximating network $f_{N_\beta}^*$, so that it has larger width N_α (corresponding to the prior network). To this end, we define the following partition of the index set $\{0, \dots, N_\alpha - 1\}$ into

$$S_n = \left\{k = l \frac{N_\alpha}{N_\beta}, l = 0, 1, \dots, N_\beta - 1\right\} \quad \text{and} \quad T_n = \{0, \dots, N_\alpha - 1\} \setminus S_n \quad (49)$$

We then let

$$f^*(x) = b^* + \sum_{k=0}^{N_\alpha-1} w_k^* (x - a_k^*)_+,$$

where $b^* = f_0(0)$,

$$w_k^* = \begin{cases} 0, & \text{if } k \in T_n, \\ w_{0;l}, \text{ for } l = \frac{kN_\beta}{N_\alpha}, & \text{if } k \in S_n. \end{cases}$$

and

$$a_k^* = \begin{cases} 0, & \text{if } k \in T_n, \\ l/N_\beta, \text{ for } l = \frac{kN_\beta}{N_\alpha}, & \text{if } k \in S_n. \end{cases}$$

It is straightforward to check that $f^* = f_{N_\beta}^*$ (notice the different ‘star notation’), so that this is indeed a relabeling of the approximating shallow network, where only the N_β weights indexed by $k \in S_n$ are nonzero; for these indices, note that $a_k^* = k/N_\alpha$. According to Lemma 10 and after the relabeling,

the nonzero weights still satisfy $|w_0^*| \leq LN_\beta^{(1-\beta)_+}$ and $|w_k^*| \leq 2LN_\beta^{1-\beta}$ for $k = 1, \dots, N_\alpha - 1$ and $|b^*| \leq L$. With this relabeling, we now have

$$\Pi[\|f - f_0\|_\infty \leq \varepsilon_n] \geq \Pi[\|f - f^*\|_\infty \leq d\varepsilon_n].$$

Recalling $a_k = k/N_\alpha$ in the definition of the prior (25), notice that whenever $w_k^* \neq 0$, it holds $a_k^* = a_k$, so that $\sum_{k=1}^{N_\alpha-1} w_k^*((x - a_k)_+ - (x - a_k^*)_+) = 0$ and thus

$$f - f^* = \sum_{k=0}^{N_\alpha-1} w_k(x - a_k)_+ - \sum_{k=1}^{N_\alpha-1} w_k^*(x - a_k^*)_+ + (b - b^*) = \sum_{k=0}^{N_\alpha-1} (w_k - w_k^*)(x - a_k)_+ + (b - b^*).$$

From the triangle inequality, follows

$$\Pi[\|f - f_0\|_\infty \leq \varepsilon_n] \geq \Pi\left(\left\|\sum_{k=0}^{N_\alpha-1} (w_k - w_k^*)(x - a_k)_+\right\|_\infty \leq d\varepsilon_n/2, |b - b^*| \leq d\varepsilon_n/2\right).$$

Since $\sup_{x \in [0,1]} (x - a_k)_+ \leq 1$, the probability displayed above is further lower bounded by

$$\Pi\left(\sum_{k=0}^{N_\alpha-1} |w_k - w_k^*| \leq d\varepsilon_n/2, |b - b^*| \leq d\varepsilon_n/2\right).$$

Using independence, this probability can be split in the following product (recall $w_k^* = 0$ for $k \in T_n$), where we set $c = d/2$

$$\underbrace{\prod_{k \in S_n} \Pi(|w_k - w_k^*| \leq c\varepsilon_n/N_\alpha)}_I \times \underbrace{\prod_{k \in T_n} \Pi(|w_k| \leq c\varepsilon_n/N_\alpha)}_{II} \times \underbrace{\Pi(|b - b^*| \leq c\varepsilon_n)}_{III}. \quad (50)$$

We first study term III, which (denoting the density of π_b also by π_b), by symmetry, positivity and continuity, is lower bounded by $2rc\varepsilon_n$ for $r = \min_{x \in [-2L, 2L]} \pi_b(x) > 0$. The latter lower bound vanishes much slower than $\exp(-n\varepsilon_n^2)$, as long as $\varepsilon_n \gtrsim n^{-s}$ for some $s < 1/2$, as is the case for the considered $\varepsilon_n \gtrsim \varepsilon_n^*$.

We then study term II. Assuming $\varepsilon_n/(\sigma_n N_\alpha) \rightarrow \infty$, the symmetry and tail assumptions (4) and (6) on h , allow us to lower bound as follows

$$\Pi(|w_k| \leq c\varepsilon_n/N_\alpha) = 1 - 2\overline{H}\left(\frac{c\varepsilon_n}{\sigma_n N_\alpha}\right) \geq 1 - 2d_0 \exp\left\{-d_1\left(\frac{c\varepsilon_n}{\sigma_n N_\alpha}\right)^q\right\}.$$

We can then lower bound the product over indices in $T_n \subset \{0, \dots, N_\alpha - 1\}$ as

$$II \geq \left(1 - 2d_0 \exp\left\{-d_1\left(\frac{\varepsilon_n}{\sigma_n N_\alpha}\right)^q\right\}\right)^{N_\alpha},$$

which, provided $\varepsilon_n/(\sigma_n N_\alpha) \gtrsim \log^{1/q} n$, can be shown to remain bounded away from zero. Indeed, it is equivalent to show that the negative logarithm remains bounded, which can be verified under the previously mentioned condition, using the inequality $\log(1 - x) \geq -x/\sqrt{1 - x}$, for $0 \leq x < 1$.

Finally for the term I , assuming without loss of generality (due to the symmetry of h) that $w_k^* > 0$, using (5) we have

$$\Pi(|w_k - w_k^*| \leq c\varepsilon_n/N_\alpha) \geq \frac{1}{\sigma_n} \int_{w_k^*}^{w_k^* + \frac{c\varepsilon_n}{N_\alpha}} h(x/\sigma_n) dx \geq cc_0 \frac{\varepsilon_n}{\sigma_n N_\alpha} \exp(-c_1 |w_k^* + c\varepsilon_n/N_\alpha|^p \sigma_n^{-p}),$$

so that, since the cardinality of S_n is N_β ,

$$I = \prod_{k \in S_n} \Pi(|w_k - w_k^*| \leq c\varepsilon_n/N_\alpha) \geq \left(c'_0 \frac{\varepsilon_n}{N_\alpha \sigma_n} \right)^{N_\beta} \exp \left\{ -c_1 \sigma_n^{-p} \sum_{k \in S_n} |w_k^* + c\varepsilon_n/N_\alpha|^p \right\}.$$

We study the sum in the exponent. Using the available bounds for w_k^* , combined with $|a + b|^p \leq |a|^p + |b|^p$ valid for any $p \in (0, 1]$, we can bound the sum as

$$\begin{aligned} \sum_{k \in S_n} |w_k^* + c\varepsilon_n/N_\alpha|^p &\lesssim |w_0|^p + \sum_{k \in S_n \setminus \{0\}} |w_k^*|^p + N_\beta \frac{\varepsilon_n^p}{N_\alpha^p} \\ &\lesssim N_\beta^{p(1-\beta)+} + N_\beta^{1+(1-\beta)p} + N_\beta \frac{\varepsilon_n^p}{N_\alpha^p}. \end{aligned} \quad (51)$$

We next note that the third term is dominated by the second if and only if $\varepsilon_n/N_\alpha \ll N_\beta^{1-\beta}$. Since $N_\alpha > N_\beta$ (recall $\alpha > \beta$), for this condition to hold it suffices that $\varepsilon_n \ll N_\beta^{2-\beta}$, which is always the case since $\beta \leq 2$. Hence the second term always dominates the third. Recalling $p \in (0, 1]$, we compare the first and second terms to find that:

1. for $\beta \in (1, 2]$, we have $1 + p - \beta p \geq (2 - \beta)p \geq 0$, hence the second term dominates the first and overall in the right hand side of the bound;
2. for $\beta \in (0, 1]$, we have $1 + p - \beta p > p(1 - \beta)_+$, hence again the second term dominates the first and overall in the right hand side of the bound.

For any $\beta \in (0, 2]$ we thus get that

$$I \geq \exp(-N_\beta(c_2 + \log(\frac{N_\alpha \sigma_n}{\varepsilon_n})) - c_3 \frac{N_\beta^{1+(1-\beta)p}}{\sigma_n^p})$$

and combining with the bounds for the previous terms, we obtain that under the assumption

$$\varepsilon_n/(\sigma_n N_\alpha) \gtrsim \log^{1/q} n, \quad (52)$$

$$\Pi(\|f - f_0\|_\infty \leq \varepsilon_n) \geq \exp \left(-N_\beta(c_2 + \log(\frac{N_\alpha \sigma_n}{\varepsilon_n})) - c_3 \frac{N_\beta^{1+(1-\beta)p}}{\sigma_n^p} - c_4 - n\varepsilon_n^2 \right).$$

The latter lower bound, is in turn lower bounded by $\exp(-c_5 n\varepsilon_n^2)$ for a large enough constant $c_5 > 0$, provided the following hold

1. $N_\beta \lesssim n\varepsilon_n^2$, or equivalently $\varepsilon_n \gtrsim \varepsilon_n^*$ which always holds;

2. $N_\beta \log(N_\alpha \sigma_n / \varepsilon_n) \lesssim n \varepsilon_n^2$ or equivalently

$$\log(N_\alpha \sigma_n / \varepsilon_n) \lesssim (\varepsilon_n / \varepsilon_n^*)^2,$$

which for $\sigma_n \leq 1$, $\varepsilon_n \gtrsim n^{-1/2}$ and N_α as defined above, holds if, for some $\delta > 0$,

$$\varepsilon_n \gtrsim \varepsilon_n^* \log^{1/2+\delta}(n), \quad (53)$$

3. $N_\beta^{1+(1-\beta)p} \lesssim \sigma_n^p n \varepsilon_n^2$, or equivalently

$$\varepsilon_n \gtrsim \varepsilon_n^* \left(\frac{N_\beta^{1-\beta}}{\sigma_n} \right)^{p/2}. \quad (54)$$

Hence, it suffices that ε_n satisfies (52) and (54) to obtain the prior mass bound (48). For the oracle choice of σ_n : we optimize the choice of σ_n based on (52), (54), and then check that (53) also holds. Since (52), (54) imply that

$$\varepsilon_n \gtrsim \{\sigma_n N_\alpha \log^{1/q} n\} \vee \{\varepsilon_n^* \left(\frac{N_\beta^{1-\beta}}{\sigma_n} \right)^{p/2}\}, \quad (55)$$

where the first term in the maximum improves with a faster decay of σ_n while the second deteriorates, we choose σ_n to balance the two terms, resulting in

$$\sigma_n \asymp N_\alpha^{-\frac{2}{2+p}} N_\beta^{\frac{p}{2+p}-\beta} \log^{-\frac{2}{q(2+p)}} n.$$

This results in

$$\varepsilon_n \gtrsim N_\alpha^{\frac{p}{2+p}} N_\beta^{\frac{p}{2+p}-\beta} \log^{\frac{p}{q(2+p)}} n = \varepsilon_n^* (N_\alpha N_\beta)^{\frac{p}{2+p}} \log^{\frac{p}{q(2+p)}} n.$$

This ε_n also satisfies (53), as required.

For the non-oracle choice of σ_n , choosing $\sigma_n = \varepsilon_n^+ / N_\alpha$, where $\varepsilon_n^+ = n^{-2/5}$ is the minimax rate for $\beta = 2$ (that is, for the highest considered smoothness), we have that the inequality arising from the first term in the maximum (55) becomes trivial and the admissible choices of ε_n are determined solely by the second term in the maximum. The resulting constraint is

$$\varepsilon_n \geq \varepsilon_n^* n^{\frac{p}{2} \left(\frac{1-\beta}{1+2\beta} + \frac{2}{5} + \frac{1}{1+2\alpha} \right)},$$

which as p becomes smaller, approaches ε_n^* . □

B.6. Proof of Theorem 5

Proof. The contraction in $L_2(P_X)$ -loss is directly obtained from the contraction in Rényi loss using Lemma 5. For the Rényi contraction result, the proof proceeds similarly to the proof of Theorem 4, but with careful tracking of the dependence of the constants on p using a simplification of the techniques employed for series priors with varying- p tails.

Indeed, up to (50) the proof is identical to the one of Theorem 4, and so is the handling of term III.

We study term II , the product over indices in T_n (defined in (49)) where $w_k^* = 0$. Let us recall that $\alpha = 0$, so that $N_\alpha = N_0 = \kappa n$, $\kappa \in [1/\sqrt{2}, \sqrt{2}]$ (recall that the definition of N_0 is 2^{m_0} , where m_0 is the closest integer solution to $2^{m_0} = n$). We need to lower bound

$$\Pi(|w_k| \leq c\varepsilon_n/n) = 1 - 2\overline{H}_{p_n}\left(\frac{c\varepsilon_n}{\sigma_n N_0}\right),$$

where $\overline{H}_{p_n}$ is the survival function of the p_n -exponential distribution defined in (3). For $\sigma_n \lesssim n^{-s}$, $s > 7/5$, it holds that for large enough n , $x_n := \frac{c\varepsilon_n}{\sigma_n N_0} \geq \frac{c\varepsilon_n}{\sqrt{2}\sigma_n n} \geq 1$ for any $\varepsilon_n \geq \varepsilon_n^+ = n^{-2/5}$, where the latter is the minimax rate for the highest considered regularity of the truth $\beta = 2$. In particular, for large enough n , $x_n \geq 1$ for ε_n as in the statement. This allows us to use the bound on the p_n -exponential cumulative distribution function from Lemma 9. In addition, our choice $\sigma_n = n^{-t}$ with $t > 2.5 > 7/5 + \log(8)/2$ combined with the choice $p_n = 2/\log n$, secure that for large n it holds $x_n^{p_n} \geq 8$, since $x_n \geq n^\tau$, with $\tau > \log(8)/2$. These considerations give that

$$\begin{aligned} \overline{H}_{p_n}(x_n) &\lesssim \frac{e^{\frac{1}{p_n}}}{\sqrt{p_n}} x_n e^{-x_n^{p_n}/p_n} \lesssim \exp\left\{\frac{1}{2} \log \frac{1}{p_n} + \frac{1}{p_n} + \log x_n - \frac{x_n^{p_n}}{p_n}\right\} \\ &\lesssim \exp\left\{\frac{2}{p_n} - \frac{x_n^{p_n}}{2p_n}\right\} \lesssim \exp\left\{-\frac{x_n^{p_n}}{4p_n}\right\}, \end{aligned} \quad (56)$$

where in the top line we used the bound from Lemma 9 and for the second line we first used $\frac{1}{2} \log \frac{1}{p_n} \leq \frac{1}{p_n}$ and $\log x_n = \frac{1}{p_n} \log x_n^{p_n} \leq \frac{1}{2p_n} x_n^{p_n}$, and then $x_n^{p_n} \geq 8$. Using $\log(1 - 2y) \geq -4y$, available for $y \leq 1/4$, and since based on the last bound $\overline{H}_{p_n}(x_n) \rightarrow 0$, for large enough n , we get

$$1 - 2\overline{H}_{p_n}(x_n) = \exp(\log(1 - 2\overline{H}_{p_n}(x_n))) \geq \exp\left(-c' \exp\left(-\frac{x_n^{p_n}}{4p_n}\right)\right),$$

for some constant $c' > 0$, which in turn gives

$$\begin{aligned} II &\geq \prod_{k \in T_n} \exp\left(-c' \exp\left(-\frac{x_n^{p_n}}{4p_n}\right)\right) = \exp\left(-c' \sum_{k \in T_n} \exp\left(-\frac{x_n^{p_n}}{4p_n}\right)\right) \\ &\geq \exp\left(-c'(\sqrt{2}n - N_\beta) \exp\left(-\frac{x_n^{p_n}}{4p_n}\right)\right). \end{aligned}$$

The latter remains bounded away from zero, since, by $x_n^{p_n} \geq 8$, we have $\frac{x_n^{p_n}}{4p_n} \geq \frac{2}{p_n} = \log n$.

We next study term I , the product over indices in S_n (defined in (49)). Assume without loss of generality that $w_k^* > 0$. The lower bound on the p_n -exponential density in Lemma 9 provides

$$\Pi\left(|w_k - w_k^*| \leq \frac{c\varepsilon_n}{N_0}\right) \geq \frac{1}{\sigma_n} \int_{w_k^*}^{w_k^* + \frac{c}{\sqrt{2}}\varepsilon_n/n} h_{p_n}\left(\frac{t}{\sigma_n}\right) dt \gtrsim \frac{\sqrt{p_n}}{\sigma_n} e^{\frac{1}{p_n} - \frac{p_n}{12}} \int_{w_k^*}^{w_k^* + \frac{c}{\sqrt{2}}\varepsilon_n/n} e^{-\frac{1}{p_n} \frac{|t|^{p_n}}{\sigma_n^{p_n}}} dt.$$

Noticing that $1/p_n - p_n/12 \geq c_0/p_n$ with $c_0 = 11/12$, we further bound

$$\Pi\left(|w_k - w_k^*| \leq \frac{c\varepsilon_n}{N_0}\right) \gtrsim e^{c_0/p_n} \frac{\sqrt{p_n}\varepsilon_n}{\sigma_n n} \exp\left\{-\frac{\sigma_n^{-p_n}}{p_n} \left|w_k^* + \frac{c\varepsilon_n}{\sqrt{2}n}\right|^{p_n}\right\},$$

so that, using $|S_n| = N_\beta$, we get for some constant $c_1 > 0$

$$I = \prod_{k \in S_n} \Pi \left(|w_k - w_k^*| \leq \frac{c\varepsilon_n}{N_0} \right) \gtrsim \left(c_1 \frac{\sqrt{p_n} e^{c_0/p_n} \varepsilon_n}{\sigma_n n} \right)^{N_\beta} \exp \left\{ -\frac{\sigma_n^{-p_n}}{p_n} \sum_{k \in S_n} \left| w_k^* + \frac{c\varepsilon_n}{\sqrt{2}n} \right|^{p_n} \right\}.$$

We study the sum, using $|a + b|^p \leq |a|^p + |b|^p$, for all $a, b \in \mathbb{R}$ and the bounds on the weights w_k^* established in the proof of Theorem 4:

$$\begin{aligned} \sum_{k \in S_n} \left| w_k^* + \frac{c\varepsilon_n}{\sqrt{2}n} \right|^{p_n} &\lesssim |w_0|^{p_n} + \sum_{k \in S_n \setminus \{0\}} |w_k^*|^{p_n} + N_\beta \frac{\varepsilon_n^{p_n}}{n^{p_n}} \\ &\lesssim L^{p_n} N_\beta^{p_n(1-\beta)+} + (2L)^{p_n} N_\beta^{p_n(1-\beta)+1} + \frac{N_\beta \varepsilon_n^{p_n}}{n^{p_n}} \\ &\lesssim N_\beta^{p_n(1-\beta)+} + N_\beta^{p_n(1-\beta)+1} + \frac{N_\beta \varepsilon_n^{p_n}}{n^{p_n}}. \end{aligned}$$

As in the proof of Theorem 4, it is straightforward to check that the second term dominates in the right hand side, using $p_n \leq 1$ and $\beta \leq 2$. Hence, for some $c'' > 0$,

$$\begin{aligned} I &\geq \exp \left(-c'' N_\beta \left(1 + \log \left(\frac{e^{-c_0/p_n}}{\sqrt{p_n}} \right) + \log(\sigma_n n / \varepsilon_n) + \frac{\sigma_n^{-p_n}}{p_n} N_\beta^{(1-\beta)p_n} \right) \right) \\ &\geq \exp \left(-c'' \left(N_\beta + \frac{\sigma_n^{-p_n}}{p_n} N_\beta^{1+(1-\beta)p_n} \right) \right), \end{aligned}$$

where for the last bound we used that, under our assumptions on $p_n, \sigma_n, \varepsilon_n$, the two logarithms in the exponent in the top line are negative.

Combining the bounds on terms I, II and III, we get, for some constant $c''' > 0$ and for sufficiently large n ,

$$\Pi(\|f - f_0\|_\infty \leq \varepsilon_n) \geq \exp \left(-c''' \left(N_\beta + \frac{\sigma_n^{-p_n}}{p_n} N_\beta^{1+(1-\beta)p_n} \right) \right),$$

which in turn is lower bounded by $\exp(-Cn\varepsilon_n^2)$ for large enough $C > 0$, provided

1. $N_\beta \lesssim n\varepsilon_n^2$ or equivalently $\varepsilon_n \gtrsim \varepsilon_n^*$;
2. it holds

$$\frac{\sigma_n^{-p_n}}{p_n} N_\beta^{1+(1-\beta)p_n} \lesssim n\varepsilon_n^2$$

or equivalently

$$\varepsilon_n \gtrsim \varepsilon_n^* \frac{n^{(\frac{1-\beta}{1+2\beta}+t)p_n/2}}{\sqrt{p_n}}.$$

The rate $\varepsilon_n \geq \varepsilon_n^* \sqrt{\log n}$ in the statement trivially satisfies the first condition, while for the second one, under our assumptions $n^{(\frac{1-\beta}{1+2\beta}+t)p_n/2}$ is bounded, so it is again satisfied. $\square$

Remark 3. It is easy to verify that the proof of Theorem 5 goes through as well with minor modifications with a pair (σ_n, p_n) similar in spirit to the second choice in Corollary 1, namely $\sigma_n = \exp(-a(\log n)^{1+\gamma})$ and $p_n = b/(\log n)^{1+\gamma}$, for $\gamma > 0$ and $a, b > 0$ such that $ab > \log 8$. Indeed, the condition $1/p_n - p_n/12 \geq c_0/p_n$ used in the proof above is also satisfied for this choice. The only difference with the proof above is that term I gives a slightly different condition: the final rate ε_n should be chosen such that $\varepsilon_n \gtrsim \varepsilon_n^*(\log n)^{(1+\gamma)/2}$, which again gives the minimax rate up to a (slightly different) logarithmic factor.

Remark 4. Theorem 5 is in fact proved for $\sigma_n = n^{-t}$ under a slightly weaker condition on t , namely $t > 7/5 + \log(8)/2$. While this result is very attractive from the theoretical point of view since it leads to full adaptation (up to $\sqrt{\log n}$) of pseudo-posteriors over β -Hölder spaces with $\beta \in (0, 2]$, in practice, the use of a neural network with width n can become expensive for large sample sizes. To address this, one can consider networks of width N_α , for some fixed $0 \leq \alpha < 2$ (recall $N_\alpha \asymp n^{\frac{1}{1+2\alpha}}$, hence the choice $\alpha = 1/2$ would lead to $\sqrt{n}$ width). In that case, with the same proof techniques, one can show that for scalings $\sigma_n = n^{-t}$, $t > 2/5 + \frac{1}{1+2\alpha} + \log(8)/2$ and tails $p_n = 1$ for $n \in \{1, \dots, \lceil e^{2(1+2\alpha)} \rceil - 1\}$ and $p_n = 2(1+2\alpha)/\log n$ for $n \geq \lceil e^{2(1+2\alpha)} \rceil$, the prior mass bound underlying the proof is satisfied for $\beta \in [\alpha, 2]$, thus (near) adaptation is again achieved, however in this α -restricted range. Aside from the computational benefits due to the smaller width, the scalings σ_n and the tails p_n are larger compared to the ones in Theorem 5, which can also be advantageous during posterior sampling. In addition, the constant $\log(8)/2$ appearing in the scaling σ_n results in scalings which for moderately large sample sizes n (such as 400 and 4000 that we use in our simulations in Section 4) are too small and lead to too smooth posteriors. An inspection of the proof, shows that one can use a smaller constant, say $\tau > 0$, in the condition on t , $t > 2/5 + \frac{1}{1+2\alpha} + \tau$, which needs to be counteracted by a larger constant say $g > 0$ in the numerator of p_n , $p_n = g/\log n$, without this affecting the rate. In fact, choosing $g = g_n = \log \log n$ and $\tau = \tau_n \rightarrow 0$ so that $g_n \tau_n \rightarrow \infty$ and $g_n/e^{\tau_n g_n} \rightarrow 0$, leads to the same rates up to logarithmic terms. For further details on these considerations see Remark 5 below. In the simulations Section 4 we used $\tau = 0$ and $g = 1$, which seems to work well for all considered sample sizes.

Remark 5. We provide some details on the modifications to the proof of Theorem 5, needed to enable the use of smaller exponents t in $\sigma_n = n^{-t}$, as discussed in Remark 4. Set $p_n = g/\log n$ and $\sigma_n = n^{-t}$ for $t > 7/5 + \tau$, $\tau > 0$. Then $x_n \geq n^{-7/5+t} = n^\tau$ and $x_n^{p_n} \geq n^{\frac{\tau g}{\log n}} = e^{\tau g}$. To bound the survival function as in (56), it suffices to establish a bound of the form $2 + \log x_n^{p_n} - x_n^{p_n} \leq -m x_n^{p_n}$ for some $m > 0$ which is such that $\exp(-m x_n^{p_n}/p_n) \leq n^{-4/5}$, so that term II is lower bounded by something of lower (or same) order as $\exp(-n \varepsilon_n^2)$, for all $\varepsilon_n \geq \varepsilon_n^+ = n^{-2/5}$. Equivalently, for m we need $m \geq 4g/(5e^{\tau g})$. To get a bound of the form above, we can use the inequality $\log x \leq \frac{\log \lambda}{\lambda} x$ valid for $x \geq \lambda \geq e$, to get that it suffices that

$$2 + (\log(\lambda)/\lambda - 1)x_n^{p_n} \leq -m x_n^{p_n}$$

or equivalently

$$e^{\tau g} \geq \frac{2}{1 - \frac{\log \lambda}{\lambda} - m}, \quad (57)$$

where $e^{\tau g} \geq \lambda$, $1 - \log \lambda/\lambda > m \geq 4g/(5e^{\tau g})$. For example, choosing τ, g so that $e^{\tau g} = 8$, we can get a bound with $g/10 \leq m \leq 3/4 - \log 8/8 \approx 0.49$ so that any $g \leq 4.9$ works. The choice of g then

determines τ via $\tau = \log 8/g$. This does not affect the final rate, since $\frac{n^{(\frac{1-\beta}{1+2\beta}+t)p_n/2}}{\sqrt{p_n}}$ remains of order $\sqrt{\log n}$ as in the end of the proof of Theorem 5.

Another possibility is to choose $g = g_n = \log \log n$ and $\tau = \tau_n \rightarrow 0$ such that $g_n \tau_n \rightarrow \infty$ and $g_n/e^{\tau_n g_n} \rightarrow 0$, in which case for any $\lambda > 0$ for sufficiently large n we have $x_n^{p_n} \geq e^{\tau_n g_n} \geq \lambda$, and it is straightforward to see that any $m < 1$ works, since (57) is satisfied as long as $m < 1 - \log \lambda/\lambda$, where λ can be chosen as large as we wish, and $m \geq 4g_n/(5e^{\tau_n g_n})$ is trivial by the condition $g_n/e^{\tau_n g_n} \rightarrow 0$. With these choices the final rate is affected, since at the end of the proof there is the requirement

$$\varepsilon_n \gtrsim \varepsilon_n^* \frac{n^{(\frac{1-\beta}{1+2\beta}+t)p_n/2}}{\sqrt{p_n}},$$

where $n^{(\frac{1-\beta}{1+2\beta}+t)p_n/2} \leq e^{r g_n} = (\log n)^r$, for some $r \geq (\frac{1-\beta}{1+2\beta} + t)/2$, recalling that $t \rightarrow 7/5$ since $\tau_n \rightarrow 0$. Thus ε_n needs to satisfy

$$\varepsilon_n \geq \varepsilon_n^* (\log n)^{1+r} / \log \log n.$$

Finally, other possibilities such as $p_n = 2/\log^s n$ and $\tau_n = \log^{s-1} n$ can be treated similarly.

Appendix C: Additional results

C.1. Handling other statistical models

We briefly explain how to derive, from the prior mass results obtained in the present paper, contraction rates in other statistical models (above we restricted for simplicity to regression models). Suppose for instance to fix ideas that one wishes to derive a posterior contraction rate for the neural network (log)-priors of Section 3 in density estimation. This is done in a similar way as for the models investigated in Agapiou and Castillo [2024]. For instance, results for nonparametric binary classification can be derived similarly as well, or in any model where one can appropriately link Kullback–Leibler neighborhoods to $\|\cdot\|_\infty$ -neighborhoods.

In the case of density estimation, one observes iid data $X_1, \dots, X_n$ of unknown density f on $[0, 1]$. Given one of the neural network priors of Section 3, one can use the exponential transform $g \rightarrow e^g / \int_0^1 e^g$ to induce a prior Π on density functions.

Using the lower bound on the prior mass in the $\|\cdot\|_\infty$ -norm investigated in the proofs of Theorems 4–5, one can use a generic result on ρ -posterior contraction such as Theorem 3 in L’Huillier et al. [2023]. The latter requires to bound from below the prior mass of a Kullback–Leibler type neighborhood: in density estimation for log-density priors this mass can be bounded from below by the prior mass of a $\|\cdot\|_\infty$ -neighborhood (see e.g. Lemma C.2 in Agapiou and Castillo [2024]). Theorem 4.1 in L’Huillier et al. [2023] then gives posterior contraction in terms of the ρ -Rényi divergence: in density estimation the latter is bounded from below by a constant times the L^1 -norm: this gives posterior contraction in the L^1 -norm in the density estimation model as desired.

C.2. Series prior: Upper bound for Sobolev truths

Define the L_2 -Sobolev-type ball, for any $\beta, L > 0$, as

$$\mathcal{S}^\beta(L) := \left\{ f = (f_k) : \sum_{k \geq 1} k^{2\beta} |f_k|^2 \leq L^2 \right\}.$$

Theorem 6. *Let $p \in (0, 1)$ and let $\alpha > \beta > 0$. Suppose $f_0 \in \mathcal{S}^\beta(L)$ for some $L > 0$ and assume $X^{(n)} \sim P_{f_0}^{(n)}$ from the model (7). Then, for any $\rho \in (0, 1)$, starting from the prior $\Pi = \Pi(p, \alpha)$ defined in (9)–(10), as $n \rightarrow \infty$, we have*

$$E_{f_0} \Pi_\rho [\|f - f_0\|_2 \leq M\varepsilon_n \mid X] \rightarrow 1,$$

where ε_n is given in (13) and $M > 0$ is a large enough constant.

Proof. As in the proof of Theorem 1, we recall that it suffices to show, for some $C, D > 0$

$$\Pi [\|f - f_0\|_2 \leq D\varepsilon_n] \geq \exp(-Cn\varepsilon_n^2).$$

Recall N_γ is the closest integer to $n^{1/(1+2\beta+p(\alpha-\beta))}$. From $f_0 \in \mathcal{S}^\beta(L)$, there exists $D > 0$, such that, as n gets large enough

$$\{f : 1 \leq k \leq N_\gamma, |f_{0,k} - f_k| \leq 1/\sqrt{n}\} \cap \{f : \forall k > N_\gamma, |f_k| \leq k^{-\beta-1/2}\} \subset \{f : \|f - f_0\|_2^2 \leq D\varepsilon_n^2\}.$$

Indeed, suppose f belongs to the left hand side set displayed just above. Using first Parseval's equality and $(a - b)^2 \leq 2a^2 + 2b^2$, we have a constant $D > 0$, such that, for n large enough

$$\begin{aligned} \|f - f_0\|_2^2 &\leq \sum_{k=1}^{N_\gamma} |f_k - f_{0,k}|^2 + 2 \sum_{k > N_\gamma} |f_k|^2 + 2 \sum_{k > N_\gamma} |f_{0,k}|^2 \\ &\leq \frac{N_\gamma}{n} + 2 \sum_{k > N_\gamma} k^{-2\beta-1} + 2 \sum_{k > N_\gamma} k^{-2\beta} k^{2\beta} |f_{0,k}|^2 \\ &\leq \frac{N_\gamma}{n} + \beta^{-1} N_\gamma^{-2\beta} + 2L^2 N_\gamma^{-2\beta} \leq D\varepsilon_n^2. \end{aligned}$$

Using independence of (f_k) under the prior, leads to

$$\Pi[\|f - f_0\|_2^2 \leq D\varepsilon_n^2] \geq \prod_{k=1}^{N_\gamma} \Pi[|f_k - f_{0,k}| \leq 1/\sqrt{n}] \times \prod_{k \geq N_\gamma} \Pi[|f_k| \leq k^{-1/2-\beta}].$$

For the first product, we lower bound the integrand with its minimal value

$$\Pi[|f_k - f_{0,k}| \leq 1/\sqrt{n}] \geq c_0 \int_{\sigma_k^{-1}(f_{0,k}-1/\sqrt{n})}^{\sigma_k^{-1}(f_{0,k}+1/\sqrt{n})} \exp(-c_1|x|^p) dx \geq \frac{2c_0}{\sigma_k \sqrt{n}} \exp\left(-c_1 \frac{|f_{0,k} + 1/\sqrt{n}|^p}{\sigma_k^p}\right).$$

Using $|a + b|^p \leq |a|^p + |b|^p$, available for $p \in (0, 1)$, we get

$$\prod_{k=1}^{N_\gamma} \Pi[|f_k - f_{0,k}| \leq 1/\sqrt{n}] \geq \prod_{k=1}^{N_\gamma} \frac{2c_0}{\sigma_k \sqrt{n}} \times \exp \left(-c_1 \sum_{k=1}^{N_\gamma} \frac{|f_{0,k}|^p + n^{-p/2}}{\sigma_k^p} \right).$$

Since $p < 1$, one notes that $\gamma \leq \alpha$ by definition, so that $\sigma_k^{-1} \geq k^{1/2+\gamma}$. Lemma 6 (applied with α therein replaced by γ) then implies

$$\prod_{k=1}^{N_\gamma} \frac{2c_0}{\sqrt{n}\sigma_k} \geq e^{-(1/2+\gamma-\log(2c_0))N_\gamma} \geq e^{-C_0 N_\gamma}.$$

Furthermore, noting that $n^{-p/2} \lesssim k^{-1/2-\beta}$, for $k \leq N_\gamma$ and $\alpha > \beta$, we get

$$\sum_{k=1}^{N_\gamma} \frac{|f_{0,k}|^p + n^{-p/2}}{\sigma_k^p} \lesssim \sum_{k=1}^{N_\gamma} k^{p(\alpha+1/2)} |f_{0,k}|^p + \sum_{k=1}^{N_\gamma} k^{p(\alpha-\beta)} \lesssim \sum_{k=1}^{N_\gamma} k^{p(\alpha+1/2)} |f_{0,k}|^p + N_\gamma^{p(\alpha-\beta)+1}.$$

To take care of the last sum we use Holder's inequality with exponents $a = 2/p > 1$ and $b = 2/(2-p)$ (such that $1/a + 1/b = 1$), as well as $f_0 \in \mathcal{S}_\beta(L)$, to obtain

$$\sum_{k=1}^{N_\gamma} k^{p(\alpha+1/2)} |f_{0,k}|^p \leq N_\gamma^{p(\alpha-\beta+1/2)} \sum_{k=1}^{N_\gamma} k^{p\beta} |f_{0,k}|^p \leq N_\gamma^{p(\alpha-\beta+1/2)} N_\gamma^{1-p/2} L^p \lesssim N_\gamma^{p(\alpha-\beta)+1}.$$

Finally, noting that for n large enough, $N_\gamma \leq N_\gamma^{p(\alpha-\beta)+1} \leq n\varepsilon_n^2$, we obtain large enough constants $C_1, \tilde{C}_1 > 0$, such that

$$\prod_{k=1}^{N_\gamma} \Pi[|f_k - f_{0,k}| \leq 1/\sqrt{n}] \geq \exp(-C_0 N_\gamma - C_1 N_\gamma^{p(\alpha-\beta)+1}) \geq \exp(\tilde{C}_1 n\varepsilon_n^2).$$

For the second product, since ζ_k 's are symmetric, have density h and survival function $\bar{H}$,

$$\prod_{k \geq N_\gamma} \Pi[|f_k| \leq k^{-1/2-\beta}] = \prod_{k \geq N_\gamma} \Pi[|\zeta_k| \leq k^{\alpha-\beta}] = \prod_{k \geq N_\gamma} (1 - 2\bar{H}(k^{\alpha-\beta})).$$

Using Condition (6), and the inequality $\log(1 - 2x) \geq -4x$, valid for $x \in [0, 1/4]$, we obtain

$$\prod_{k \geq N_\gamma} (1 - 2\bar{H}(k^{\alpha-\beta})) \geq \exp \sum_{k > N_\gamma} \log(1 - 2d_0 e^{-d_1 k^{q(\alpha-\beta)}}) \geq \exp\{-4d_0 \sum_{k > N_\gamma} e^{-d_1 k^{q(\alpha-\beta)}}\}.$$

Since the series $\sum_k e^{-ck^\delta}$ converges for any given constants $c, \delta > 0$, one deduces that the last display converges to 1 as $n \rightarrow \infty$ and in particular is bounded from below by $1/2$ for n large enough. Gathering the previous bounds for both products provides a constant $C > 0$, such that for n large enough, one obtains

$$\Pi[\|f - f_0\|_2^2 \leq D\varepsilon_n^2] \geq \exp(-Cn\varepsilon_n^2). \quad \square$$

C.3. Series prior: Lower bound (case $p = 1$)

The following is the special case of Theorem 2 where $p = 1$ (Laplace priors on coefficients).

Theorem 7. *Let $\alpha > \beta > 0$. Suppose the data X follows the white noise model for some true function f_0 , and let the prior on f as in (9) be defined by taking ζ_k s to be drawn iid from a standard Laplace distribution. There exists a function $f_0 \in \mathcal{F}(\beta, L)$ such that, if*

$$\zeta_n = \varepsilon_n(1, \alpha, \beta) = n^{-\frac{\beta}{\alpha+\beta+1}},$$

then for any $\rho \in (0, 1]$, for $m > 0$ small enough, as $n \rightarrow \infty$,

$$E_{f_0} \Pi_\rho[\|f - f_0\|_2 < m\zeta_n \mid X] \rightarrow 0.$$

Proof. To simplify the notation, we give the proof first for the standard posterior $\rho = 1$. Let us choose f_0 as the function in $\mathcal{F}(\beta, L)$ defined through its basis coefficients by $f_{0,k} = Lk^{-1/2-\beta}$ for $\beta > 0$. Let us recall the definitions of γ in (12) and N_γ in (15) and let us set $p = 1$. With this notation we have $\zeta_n = \varepsilon_n(1, \alpha, \beta) = N_\gamma^{-\beta}$ by definition. Denoting, for any square-integrable function g , by $\|g\|_{N_\gamma}^2 = \sum_{k=1}^{N_\gamma} g_k^2$, it is enough to prove, for small $m > 0$ to be chosen and $n_\gamma = dN_\gamma$ for some small enough constant d to be chosen below, that

$$E_{f_0} \Pi[\|f - f_0\|_{n_\gamma} \geq m\zeta_n \mid X] \rightarrow 1.$$

Under the prior distribution, coefficients f_k have distribution $\sigma_k \text{Lap}(1)$. By writing the Laplace distribution as a mixture of two exponential distributions (one for the positive part, one for the negative part), and using Bayes' formula, one can write the posterior distribution $\mathcal{L}(f_k \mid X)$ of the k th coefficient f_k as a mixture

$$\mathcal{L}(f_k \mid X) = w_k^+ \mathcal{N}(\mu_k, 1/n)_+ + (1 - w_k^+) \mathcal{N}(\nu_k, 1/n)_-, \quad (58)$$

where $\mathcal{N}(\mu, \sigma^2)_+$ denotes the distribution of $Z \vee 0$ if $Z \sim \mathcal{N}(\mu, \sigma^2)$ and $\mathcal{N}(\mu, \sigma^2)_-$ the one of $Z \wedge 0$, and where we have set

$$\mu_k = X_k - \frac{1}{n\sigma_k}, \quad \nu_k = X_k + \frac{1}{n\sigma_k} \quad (59)$$

$$w_k^+ = \frac{e^{n\mu_k^2/2} \Phi(\sqrt{n}\mu_k)}{e^{n\nu_k^2/2} \bar{\Phi}(\sqrt{n}\nu_k) + e^{n\mu_k^2/2} \Phi(\sqrt{n}\mu_k)}, \quad (60)$$

where Φ is the distribution function of the standard normal distribution and $\bar{\Phi} = 1 - \Phi$.

The triangle inequality gives, denoting $\mu = (\mu_k)$ for μ_k as in (59), that $\|f_0 - \mu\|_{n_\gamma} \leq \|f_0 - f\|_{n_\gamma} + \|f - \mu\|_{n_\gamma}$. This implies

$$\Pi[\|f - f_0\|_{n_\gamma} \geq m\zeta_n \mid X] \geq \Pi[\|f - \mu\|_{n_\gamma} \leq m\zeta_n \mid X] \cdot \mathbb{1}\{\|f_0 - \mu\|_{n_\gamma} \geq 2m\zeta_n\}.$$

It now suffices to show that each term of the product of the right hand side of the last display goes to 1 in probability under P_{f_0} .

Starting with the indicator, its expectation under P_{f_0} equals, denoting $X = (X_k)$, and using $\|\mu - f_0\|_{n_\gamma} \geq \|\mu - X\|_{n_\gamma} - \|X - f_0\|_{n_\gamma}$ by the triangle inequality,

$$P[\|\mu - f_0\|_{n_\gamma} \geq 2m\zeta_n] \geq P[\|X - f_0\|_{n_\gamma} \leq m\zeta_n] \cdot \mathbb{1}\{\|\mu - X\|_{n_\gamma} \geq 3m\zeta_n\}.$$

By definitions of μ and (σ_k) , we have, for a constant $C_\alpha > 0$ depending only on α ,

$$\|\mu - X\|_{n_\gamma}^2 = \sum_{k=1}^{n_\gamma} \frac{\sigma_k^{-2}}{n^2} \geq C_\alpha \frac{n_\gamma^{2+2\alpha}}{n^2} = C' \zeta_n^2.$$

for $C' = C'(\alpha, d)$ and d the constant such that $n_\gamma = dN_\gamma$. Hence for m small enough so that $3m < C'$, the indicator in the last but one display equals 1. Also, $\|X - f_0\|_{n_\gamma}^2 = \sum_{k=1}^{n_\gamma} \varepsilon_k^2/n$. The later quantity has expectation $n_\gamma/n = n^{-(\alpha+\beta)/(1+\alpha+\beta)} = o(\zeta_n^2)$ since $\alpha > \beta$ by assumption. A standard concentration argument (e.g. using Tchebychev's inequality, or a more precise exponential concentration bound for the χ^2 distribution) then gives $P[\|X - f_0\|_{n_\gamma} \leq m\zeta_n] = 1 + o(1)$.

To conclude the proof, it suffices to check that $\Pi[\|f - \mu\|_{n_\gamma} \leq m\zeta_n \mid X]$ goes to 1 in probability under P_{f_0} . By Markov's inequality, for $\mathcal{A}_n$ the event as in (61),

$$\begin{aligned} \Pi[\|f - \mu\|_{n_\gamma} > m\zeta_n \mid X] \mathbb{1}_{\mathcal{A}_n} &\leq \frac{1}{(m\zeta_n)^2} \int \|f - \mu\|_{n_\gamma}^2 d\Pi(f \mid X) \mathbb{1}_{\mathcal{A}_n} \\ &\leq \frac{\mathbb{1}_{\mathcal{A}_n}}{(m\zeta_n)^2} \sum_{k=1}^{n_\gamma} \left[w_k^+ \int (f_k - \mu_k)^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k) + (1 - w_k^+) \int (f_k - \mu_k)^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \right] \\ &\leq \frac{\mathbb{1}_{\mathcal{A}_n}}{(m\zeta_n)^2} \sum_{k=1}^{n_\gamma} \left[\int (f_k - \mu_k)^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k) + c_1 e^{-c_2 n^{\frac{\alpha-\beta}{2\gamma+1}}} \int (f_k - \mu_k)^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \right], \end{aligned}$$

where for the last inequality we use $w_k^+ \leq 1$ and the uniform bound on $(1 - w_k^+)$ obtained in Lemma 1, and $2\gamma = \alpha + \beta$. The first integral on the last line can be written

$$\int (u - \mu_k)^2 \sqrt{n} \phi(\sqrt{n}(u - \mu_k)) \mathbb{1}\{u \geq 0\} du / \int \sqrt{n} \phi(\sqrt{n}(u - \mu_k)) \mathbb{1}\{u \geq 0\} du.$$

The denominator equals $\bar{\Phi}(-\sqrt{n}\mu_k) = \Phi(\sqrt{n}\mu_k)$. By Lemma 1, on the event $\mathcal{A}_n$ (see (61)) of overwhelming probability, it holds $\sqrt{n}\mu_k \geq \sqrt{n}f_{0,k}/4$, which is bounded away from 0 for $k \leq n_\gamma$, so that $\Phi(\sqrt{n}\mu_k) \geq 1/2$ for such k 's. One can then bound the numerator in the last display from above by

$$\int (u - \mu_k)^2 \sqrt{n} \phi(\sqrt{n}(u - \mu_k)) du = \frac{1}{n} \int u^2 \phi(u) du = 1/n.$$

We now bound the second integral in the former display on $\mathcal{A}_n$, using first $(f_k - \mu_k)^2 \leq 2f_k^2 + 2\mu_k^2 \leq 2f_k^2 + 2\nu_k^2$ using $0 \leq \mu_k \leq \nu_k$ on $\mathcal{A}_n$ and then

$$\int f_k^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \leq E[(Z + \nu_k)_-^2] \leq 2E[Z^2] + 2\nu_k^2 \leq 2/n + 2\nu_k^2,$$

with $Z \sim \mathcal{N}(0, 1/n)$, where we have used $E[Y_-^2] \leq E[Y^2]$ for any variable Y (here $Y = Z + \nu_k$). Also,

$$E_{f_0}[\nu_k^2] \leq 2E_{f_0}[X_k^2] + 2/(n\sigma_k)^2 \leq 4f_{0,k}^2 + 2/n + 2/(n\sigma_k)^2.$$

This leads to $\sum_{k=1}^{n_\gamma} E_{f_0}[\nu_k^2] \lesssim C + n_\gamma/n + \zeta_n^2$, where we have used that f_0 is squared-integrable and that $\sum_{k=1}^{n_\gamma} 1/(n\sigma_k)^2 \lesssim \zeta_n^2$. By gathering the previous bounds one obtains

$$\begin{aligned} E_{f_0} [\Pi[\|f - \mu\|_{n_\gamma} > m\zeta_n \mid X] \mathbb{1}_{\mathcal{A}_n}] &\leq \frac{1}{(m\zeta_n)^2} \left[2\frac{n_\gamma}{n} + c_1 e^{-c_2 n^{\frac{\alpha-\beta}{2\gamma+1}}} C \left\{ 1 + n_\gamma/n + \zeta_n^2 \right\} \right] \\ &\lesssim (n_\gamma/n)\zeta_n^{-2} + \zeta_n^{-2} e^{-c_2 n^{\frac{\alpha-\beta}{2\gamma+1}}} = o(1), \end{aligned}$$

using that $(n_\gamma/n)\zeta_n^{-2} = o(1)$ by the definitions of n_γ, ζ_n and using $\alpha > \beta$ by assumption. Since $P_{f_0}[\mathcal{A}_n] = 1 + o(1)$ by Lemma 1, this implies $E_{f_0} [\Pi[\|f - \mu\|_{n_\gamma} > m\zeta_n \mid X]] = o(1)$ as desired, which concludes the proof for the usual posterior.

For the ρ -posterior with $\rho < 1$, the proof is mostly the same: first one notes that the ρ -posterior on the k th coordinate is the mixture distribution

$$w_k^+ \mathcal{N}(\mu_k, 1/n')_+ + (1 - w_k^+) \mathcal{N}(\nu_k, 1/n')_-,$$

where we have set $n' = n\rho$ and with the updated definitions

$$\begin{aligned} \mu_k &= X_k - \frac{1}{n'\sigma_k}, \quad \nu_k = X_k + \frac{1}{n'\sigma_k} \\ w_k^+ &= \frac{e^{n'\mu_k^2/2} \Phi(\sqrt{n'}\mu_k)}{e^{n'\nu_k^2/2} \Phi(\sqrt{n'}\nu_k) + e^{n'\mu_k^2/2} \Phi(\sqrt{n'}\mu_k)}. \end{aligned}$$

The above proof and that of Lemma 1 for $\rho = 1$ both go through with n replaced by the updated ‘effective’ sample size $n' = n\rho$; since $\rho \in (0, 1]$ is fixed, this only changes the constants in the obtained rates, which concludes the proof. $\square$

Lemma 1. Let f_0 be defined by $f_{0,k} = Lk^{-1/2-\beta}$ for $\beta > 0$ and let $\sigma_k = k^{-1/2-\alpha}$ for $\alpha > 0$.

1. Let $\mathcal{A}_n$ be the event defined by, for μ_k as in (59) and $n_\gamma = dN_\gamma$,

$$\mathcal{A}_n = \{ \mu_k \geq f_{0,k}/4, \text{ for all } k = 1, \dots, n_\gamma \}. \quad (61)$$

Then, for a small enough constant $d > 0$ above, one has $P_{f_0}[\mathcal{A}_n] = 1 + o(1)$ as $n \rightarrow \infty$.

2. There exist constants $c_1, c_2 > 0$ such that, for w_k^+ as in (60), on the event $\mathcal{A}_n$ as in (61),

$$\max_{1 \leq k \leq n_\gamma} (1 - w_k^+) \leq c_1 e^{-c_2 n^{(\alpha-\beta)/(\alpha+\beta+1)}}.$$

Proof. One first notes that for small enough d , for any $k \leq n_\gamma = dN_\gamma$, one has $f_{0,k}/2 \geq (n\sigma_k)^{-1}$, by definition of $f_{0,k}$. Since under P_{f_0} we have $\mu_k = X_k - (n\sigma_k)^{-1} = f_{0,k} + \varepsilon_k/\sqrt{n} - (n\sigma_k)^{-1}$, it holds $\mu_k \geq f_{0,k}/2 + \varepsilon_k/\sqrt{n} \geq f_{0,k}/2 - \sqrt{2\log n/n}$ on the event

$$\mathcal{B}_n = \{ |\varepsilon_k| \leq \sqrt{2\log n}, \text{ for all } k = 1, \dots, n_\gamma \}.$$

A union bound shows that $P_{f_0}[\mathcal{B}_n^c] = o(1)$. Also, by definition of $f_{0,k}$ and n_γ , we have $\sqrt{2\log n/n} \leq f_{0,k}/4$ so that $\mathcal{B}_n \subset \mathcal{A}_n$ which implies $P_{f_0}[\mathcal{A}_n] = 1 + o(1)$ as $n \rightarrow \infty$.

On the other hand, note that $nf_{0,k}^2 \geq nn_\gamma^{-1-2\beta} \gtrsim n^{(\alpha-\beta)/(2\gamma+1)}$, so that $n\mu_k^2 \gtrsim n^{(\alpha-\beta)/(2\gamma+1)}$ on the event $\mathcal{A}_n$. This shows that $e^{n\mu_k^2/2}\Phi(\sqrt{n}\mu_k) \gtrsim e^{c_2n^{(\alpha-\beta)/(2\gamma+1)}}$ uniformly over $k = 1, \dots, n_\gamma$, for some $c_2 > 0$.

Also, by definition, $\nu_k \geq \mu_k$ and the latter is positive on $\mathcal{A}_n$, so since $e^{u^2/2}\bar{\Phi}(u) \lesssim 1/u$ for $u > 0$, one obtains $e^{n\nu_k^2/2}\bar{\Phi}(\sqrt{n}\nu_k) \leq 1/(\sqrt{n}\nu_k)$ uniformly over $k \leq n_\gamma$, which is bounded from above by a constant (since $\sqrt{n}\nu_k \geq \sqrt{n}\mu_k$ goes to infinity on $\mathcal{A}_n$).

Putting together the two previously obtained bounds, one obtains on $\mathcal{A}_n$

$$1 - w_k^+ \leq \frac{e^{n\nu_k^2/2}\bar{\Phi}(\sqrt{n}\nu_k)}{e^{n\mu_k^2/2}\Phi(\sqrt{n}\mu_k)} \lesssim e^{-c_2n^{(\alpha-\beta)/(2\gamma+1)}}/(\sqrt{n}\nu_k) \lesssim e^{-c_2n^{(\alpha-\beta)/(2\gamma+1)}},$$

uniformly over $k = 1, \dots, n_\gamma$, which concludes the proof. $\square$

C.4. Series prior: Upper bound ($p = 1$) for classical posteriors $\rho = 1$

The next result provides an example of extension of Theorem 1 in the main paper (which considers ρ -posteriors, $\rho < 1$) to classical posteriors ($\rho = 1$). It focuses on the case of Laplace tails $p = 1$ for simplicity, although a similar result is expected to hold for other p 's as well, albeit with more technical proofs, so for clarity we focus on $p = 1$ here.

Theorem 8. *Let $\alpha > \beta > 0$. Suppose the data X follows the white noise model for some true function f_0 , and let the prior on f be as in (9) and taking ζ_k 's to be drawn iid from a standard Laplace distribution. For any function $f_0 \in \mathcal{F}(\beta, L)$, if*

$$\varepsilon_n = \varepsilon_n(1, \alpha, \beta) = n^{-\frac{\beta}{\alpha+\beta+1}},$$

then for $M > 0$ large enough, for some $b > 0$, as $n \rightarrow \infty$,

$$E_{f_0} \Pi[\|f - f_0\|_2 > M\varepsilon_n \mid X] \rightarrow 0.$$

Proof. Let $K_n := DN_\gamma$, where D is a large enough constant to be chosen below. We write $\|f - f_0\|^2 = \sum_{k \geq 1} (f_k - f_{0,k})^2 = (\sum_{k \leq K_n} + \sum_{k > K_n}) (f_k - f_{0,k})^2$ and distinguish two cases: $k \leq K_n$ and $k > K_n$.

We deal first with the indices $k > K_n$. By definition of N_γ, ε_n and for $f_0 \in \mathcal{F}(\beta, L)$, it holds

$$\sum_{k > K_n} f_{0,k}^2 \leq L^2 K_n^{-2\beta} = L^2 N_\gamma^{-2\beta} \lesssim \varepsilon_n^2.$$

As $\|f^{[K_n^c]} - f_0^{[K_n^c]}\|_2 \leq \|f^{[K_n^c]}\|_2 + \|f_0^{[K_n^c]}\|_2$, provided we choose M sufficiently large with respect to D, L , it suffices to show that $E_{f_0} \Pi[\|f^{[K_n^c]}\|_2 > M\varepsilon_n/2 \mid X] \rightarrow 0$. To do so, by Markov's inequality, it suffices to show that, on an event of high probability,

$$\sum_{k > K_n} \int f_k^2 d\Pi(f_k \mid X_k) = o(\varepsilon_n^2). \quad (62)$$

Let us consider the event

$$\mathcal{A} := \bigcup_{j \geq 0} \left\{ \max_{jn < k \leq (j+1)n} |\varepsilon_k| \leq \sqrt{2 \log \{(j+1)^2 n\}} \right\}.$$

A union bound argument shows that $P[\mathcal{A}^c] = o(1)$. By definition of K_n, σ_k and the fact that $f_0 \in \mathcal{F}(\beta, L)$, we have, for any $k > K_n$ that $|f_{0,k}| \leq (1/2)(n\sigma_k)^{-1}$, provided the constant D is chosen large enough. This implies, on the event $\mathcal{A}$, that for any $K_n \leq k \leq n$,

$$\mu_k = X_k - \frac{1}{n\sigma_k} \leq -\frac{1}{2n\sigma_k} + \sqrt{\frac{2 \log n}{n}}.$$

Since $\alpha > \beta$ we have $\sqrt{\log n/n} = o(1/(n\sigma_k))$ for $k > K_n$, so that for $K_n < k \leq n$, on $\mathcal{A}$,

$$\mu_k \leq -\frac{1}{4n\sigma_k}.$$

Since the growth of σ_k^{-1} is polynomial in k , this also implies that the inequality in the last display also holds for large enough n and any index $k > n$, so that the inequality holds for all $k > K_n$. Similarly, we have, on the event $\mathcal{A}$ and for any $k > K_n$,

$$\nu_k \geq \frac{1}{4n\sigma_k}.$$

Now bounding w_k^+ and $1 - w_k^+$ by 1, one can bound from above, on the event $\mathcal{A}$,

$$\int f_k^2 d\Pi(f_k | X_k) \leq \int f_k^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k) + \int f_k^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \leq \frac{2}{n^2} \frac{1}{\mu_k^2} + \frac{2}{n^2} \frac{1}{\nu_k^2},$$

where the second inequality follows from Lemma 2 with $\mu = \mu_k, \nu = \nu_k$ and $\sigma^2 = 1/n$. One deduces that on $\mathcal{A}$, the last display is bounded from above by $C\sigma_k^2$. Since $\sum_{k > K_n} \sigma_k^2 \lesssim K_n^{-2\alpha} = o(K_n^{-2\beta})$ since $\alpha > \beta$, and next using that $K_n^{-2\beta} = O(\varepsilon_n^2)$, one concludes that $\sum_{k > K_n} \int f_k^2 d\Pi(f_k | X_k) = o_P(\varepsilon_n^2)$ as desired.

It now remains to deal with the indices $k \leq K_n$. Here one can follow the final bounds in the lower bound argument in Theorem 7 and extend these to any function $f_0 \in \mathcal{F}(\beta, L)$. Denoting by $\|\cdot\|_{K_n}$ the L^2 -norm truncated to the first K_n coefficients, let us define a ‘centering’ function $h = h(X)$ from its basis coefficients (h_k) as follows: $h_k = 0$ for $k > K_n$ and, for $k \leq K_n$,

$$h_k = \begin{cases} \mu_k := X_k - 1/(n\sigma_k) & \text{if } \mu_k > (\log n)/\sqrt{n}, \\ \nu_k := X_k + 1/(n\sigma_k) & \text{if } \nu_k < -(\log n)/\sqrt{n}, \\ 0 & \text{otherwise.} \end{cases} \quad (63)$$

Note that the first two cases in the above definition are mutually exclusive, as $\mu_k \leq \nu_k$ by definition. Now one can further write, for M a large constant to be chosen below,

$$\begin{aligned} \Pi[\|f - f_0\|_{K_n} \geq M\varepsilon_n | X] \\ = \Pi[\|f - f_0\|_{K_n} \geq M\varepsilon_n | X] \mathbb{I}\{\|f_0 - h\|_{K_n} \leq M\varepsilon_n/2\} \end{aligned}$$

$$\begin{aligned}
& + \Pi[\|f - f_0\|_{K_n} \geq M\varepsilon_n \mid X] \mathbb{1}\{\|f_0 - h\|_{K_n} > M\varepsilon_n/2\} \\
& \leq \Pi[\|f - h\|_{K_n} \geq M\varepsilon_n/2 \mid X] + \mathbb{1}\{\|f_0 - h\|_{K_n} > M\varepsilon_n/2\},
\end{aligned}$$

where one uses the triangle inequality and that indicators and probabilities are bounded from above by 1. It now suffices to show that the expectation under $P_{f_0}^{(n)}$ of the last display goes to 0.

Starting first with the indicator, and denoting by $e(\cdot)$ the function with coefficients $e_k = \varepsilon_k$ for $k \leq K_n$ and 0 otherwise, applying the triangle inequality gives

$$P_{f_0}^{(n)}[\|f_0 - h\|_{K_n} > M\varepsilon_n/2] \leq P[\|e\|_{K_n}/\sqrt{n} + \|(1/(n\sigma_k))\|_{K_n} > M\varepsilon_n/2].$$

By our choice of K_n , we have $\|(1/(n\sigma_k))\|_{K_n}^2 = n^{-2} \sum_{k=1}^{KN_\gamma} \sigma_k^{-2} \lesssim \varepsilon_n^2$. Hence for M large enough the last norm in the above display is less than $M\varepsilon_n/4$. It now suffices to bound $P[\|e\|_{K_n}/\sqrt{n} > M\varepsilon_n/4]$. By Markov's inequality, this is bounded by a multiple of $(n\varepsilon_n^2)^{-1} E[\|e\|_{K_n}^2] = (n\varepsilon_n^2)^{-1} K_n^2 = o(1)$, by definition of ε_n, K_n .

Finally, it now remains to deal with the term $E_{f_0} \Pi[\|f - h\|_{K_n} \geq M\varepsilon_n/2 \mid X]$. By Markov's inequality, to show that this term is a $o(1)$, it suffices to check that $E_{f_0} \int \|f - h\|_{K_n}^2 d\Pi(f \mid X) = o(\varepsilon_n^2)$, which is done in Lemma 3 below. This concludes the proof of Theorem 8. $\square$

Let us recall that $\mathcal{N}(\mu, \sigma^2)_+$ denotes the distribution of $Z \vee 0$ if $Z \sim \mathcal{N}(\mu, \sigma^2)$ and $\mathcal{N}(\mu, \sigma^2)_-$ the one of $Z \wedge 0$.

Lemma 2. *For any $\mu < 0$ and $\sigma^2 > 0$, the following bound holds*

$$\int x^2 d\mathcal{N}(\mu, \sigma^2)_+(x) \leq 2 \frac{\sigma^4}{\mu^2}.$$

Similarly, for any $\nu < 0$ and $\sigma^2 > 0$, it holds $\int x^2 d\mathcal{N}(\nu, \sigma^2)_-(x) \leq 2\sigma^4/\nu^2$.

Proof. The integral in the display of the lemma equals, with $z = -\mu > 0$,

$$\mathcal{I} := \int_0^\infty x^2 \exp\{-(x+z)^2/(2\sigma^2)\} dx / \mathcal{D},$$

where $\mathcal{D} = \int_0^\infty \exp\{-(x+z)^2/(2\sigma^2)\} dx$. Integrating by part once gives

$$\mathcal{D} \cdot \mathcal{I} = \int_0^\infty \frac{2x}{x+z} \sigma^2 \exp\{-(x+z)^2/(2\sigma^2)\} dx,$$

where the bracket term vanishes both at 0 and infinity. Integrating by part once more,

$$\mathcal{D} \cdot \mathcal{I} = \int_0^\infty \frac{2}{(x+z)^2} \sigma^4 \exp\{-(x+z)^2/(2\sigma^2)\} dx.$$

Using $x+z \geq z > 0$ and $z^2 = \mu^2$, the last display is bounded from above by $2\sigma^4 \mathcal{D}/z^2$, which implies the first bound of the lemma. The second part follows by symmetry. $\square$

Lemma 3. *Let h be the random function defined by (63). Then for any $D > 0$ and $K_n := DN_\gamma$, it holds*

$$E_{f_0} \int \|f - h\|_{K_n}^2 d\Pi(f | X) = o(\varepsilon_n^2),$$

where ε_n is the rate in the statement of Theorem 8.

Proof. We use the expression (58) of the posterior and write the integral in the Lemma

$$\mathcal{I} := \sum_{k=1}^{K_n} \left[w_k^+ \int (f_k - h_k)^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k) + (1 - w_k^+) \int (f_k - h_k)^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \right].$$

Let us distinguish three cases depending on the split of indices in the definition (63) of h , namely $\mathcal{C}_1 = \{k \leq K_n : \mu_k > (\log n)/\sqrt{n}\}$ (Case 1), $\mathcal{C}_2 = \{k \leq K_n : \nu_k < -(\log n)/\sqrt{n}\}$ (Case 2) and $\mathcal{C}_3 := \{k \leq K_n : \mu_k \leq (\log n)/\sqrt{n}, \nu_k \geq -(\log n)/\sqrt{n}\}$ (Case 3). Note that the corresponding sets of indices is random, so has random cardinality, but in all cases once we bound the corresponding quantities by convenient upper-bounds, we will eventually just bound the sum over the corresponding k 's (which is over a random set) simply by that over all $k \leq K_n$.

Starting with Case 3, since $h_k = 0$ in that case and using $0 \leq w_k^+ \leq 1$, it is enough to bound from above

$$\sum_{k=1}^{K_n} \left[\int f_k^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k) + \int f_k^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \right] \mathbb{1}\{k \in \mathcal{C}_3\}.$$

By symmetry it is enough to deal with the first integral in the last display, the other being dealt with similarly. We thus focus on bounding from above $I_k := \int f_k^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k)$. The following simple bound always holds $I_k = E[Z_+^2] \leq E[Z^2] \leq 2\mu_k^2 + 2/n$. In case $\mu_k \geq 0$, for $k \in \mathcal{C}_3$ by definition this is then further bounded by $2(\log n)^2/n + 2/n$. In case $\mu_k < 0$, on top of the previous bound we can also now use Lemma 2 to get $I_k \leq 2/(n\mu_k)^2$. This means that in that case $I_k/2 \leq \min(\mu_k^2 + 1/n, 1/(n\mu_k)^2) \leq 2/n$ (by comparing the bound to the case $\mu_k = 1/\sqrt{n}$). Hence

$$\sum_{k=1}^{K_n} \int f_k^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k) \mathbb{1}\{k \in \mathcal{C}_3\} \leq \sum_{k=1}^{K_n} 4(\log^2 n)/n = 4K_n(\log^2 n)/n = o(\varepsilon_n^2),$$

where for the last comparison we use $\alpha > \beta$.

Now dealing with Case 1, we use the expression of the weight w_k^+ to get

$$1 - w_k^+ \leq \frac{e^{n\nu_k^2/2} \bar{\Phi}(\sqrt{n}\nu_k)}{e^{n\mu_k^2/2} \Phi(\sqrt{n}\mu_k)} \leq \frac{2}{\sqrt{n}\nu_k} e^{-n\mu_k^2/2},$$

by using $\Phi(\sqrt{n}\mu_k) \geq \Phi(0) = 1/2$ and the bound $\bar{\Phi}(y) \leq \phi(y)/y$ for $y > 0$. Since in Case 1 one both has $\nu_k \geq \mu_k > 0$ and $\mu_k > (\log n)/\sqrt{n}$, one gets, for $k \in \mathcal{C}_1$,

$$1 - w_k^+ \leq \frac{2}{\log n} e^{-(\log n)^2/2}.$$

This implies the bound

$$\begin{aligned} \mathcal{I}_1 &:= \sum_{k=1}^{K_n} \left[w_k^+ \int (f_k - h_k)^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k) + (1 - w_k^+) \int (f_k - h_k)^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \right] \mathbb{1}_{k \in \mathcal{C}_1} \\ &\leq \sum_{k=1}^{K_n} \left[\int (f_k - \mu_k)^2 d\mathcal{N}(\mu_k, 1/n)_+(f_k) + C e^{-(\log n)^2/2} \int (f_k - \mu_k)^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \right] \mathbb{1}_{k \in \mathcal{C}_1} \end{aligned}$$

Both terms are now bounded in a similar way as in the proof of Theorem 7. The first integral in the last line is bounded from above by $\int (u - \mu_k)^2 \sqrt{n} \phi(\sqrt{n}(u - \mu_k)) \mathbb{1}_{u \geq 0} du / \Phi(0)$, using $\Phi(\sqrt{n}\mu_k) \geq \Phi(0) = 1/2$ since $\sqrt{n}\mu_k > 0$, so that the integral is at most $2/n$. One deduces

$$\mathcal{I}_1 \leq \sum_{k=1}^{K_n} \left[\frac{2}{n} + C e^{-(\log n)^2/2} \int (f_k - \mu_k)^2 d\mathcal{N}(\nu_k, 1/n)_-(f_k) \right].$$

Now note that the expectation under E_{f_0} of the integral in the last display has been bounded from above in the proof of Theorem 7 by a constant C (n_γ therein is replaced by K_n , which does not change the bound, up to a multiplicative constant). This implies

$$\mathcal{I}_1 \leq 2K_n/n + CK_n e^{-(\log n)^2/2} = o(\varepsilon_n).$$

Finally Case 2 is handled exactly as Case 1, by symmetry. Putting the three obtained bounds together concludes the proof. $\square$

C.5. SNN: prior with $p > 1$

The following theorem complements Theorem 4 in the lighter than Laplace tails case ($p > 1$).

Theorem 9. *Consider the setting of Theorem 4 but with $p > 1$. For any $\rho \in (0, 1)$, denoting D_ρ the Rényi divergence (26), there exists a large enough constant $M > 0$, such that, as $n \rightarrow \infty$,*

$$E_{f_0} \Pi_\rho \left[\left\{ f : \frac{1}{n} D_\rho(P_f^n, P_{f_0}^n) \geq M \varepsilon_n^2 \right\} \mid X, Y \right] \rightarrow 0,$$

where, letting $\varepsilon_n^* = n^{-\frac{\beta}{1+2\beta}}$ and $\varepsilon_n^+ = n^{-2/5}$ (equal to ε_n^* for $\beta = 2$), ε_n is given as follows:

a) if $\beta \in (0, 1 + \frac{1}{p}]$

$$i) \text{ (Oracle } \sigma_n) \text{ for } \sigma_n = N_\alpha^{-\frac{2}{2+p}} N_\beta^{\frac{2}{2+p} - \beta} \log^{-\frac{2}{q(2+p)}}(n),$$

$$\varepsilon_n = \varepsilon_n^* (N_\alpha N_\beta)^{\frac{p}{2+p}} \log^{\frac{p}{q(2+p)}}(n);$$

ii) (Non-oracle σ_n) for $\sigma_n = \varepsilon_n^+ / N_\alpha$

$$\varepsilon_n = \varepsilon_n^* n^{\frac{p}{2}(\frac{1-\beta}{1+2\beta} + \frac{2}{5} + \frac{1}{1+2\alpha})};$$

b) if $\beta \in (1 + \frac{1}{p}, 2]$

i) (Oracle σ_n) for $\sigma_n = (\varepsilon_n^*)^{\frac{2}{2+p}} N_\alpha^{-\frac{2}{2+p}} N_\beta^{-\frac{1}{2+p}} \log^{-\frac{2}{q(2+p)}}(n)$,

$$\varepsilon_n = \varepsilon_n^* N_\alpha^{\frac{p}{2+p}} N_\beta^{\frac{\beta p - 1}{2+p}} \log^{\frac{p}{q(2+p)}} n;$$

ii) (Non-oracle σ_n) for $\sigma_n = \varepsilon_n^+ / N_\alpha$

$$\varepsilon_n = \varepsilon_n^* n^{\frac{p}{5} - \frac{1}{2(1+2\beta)}};$$

Proof. The proof is identical to the proof of Theorem 4 up to (51), which we can still get using $|a + b|^p \leq c(p)(|a|^p + |b|^p)$, valid for any $p > 0$. We next note that in the case $p > 1$

1. for $\beta \in (0, 1]$, we have $1 + p - \beta p > p(1 - \beta)_+$, hence again the second term dominates the first in (51) and overall in the right hand side of the bound;
2. for $\beta \in (1, 2]$, the second term dominates the first for $p \leq (\beta - 1)^{-1}$, otherwise the first term dominates in the right hand side of the bound.

For $\beta \in (0, 1 + \frac{1}{p}]$, the remaining of the proof is still identical to the proof of Theorem 4, and we only need to deal with the case $\beta \in (1 + \frac{1}{p}, 2]$, in which the first term dominates in (51) and, in fact, the sum is bounded by a constant. We hence get, for some constant c_3 ,

$$I \geq \exp\left(-N_\beta(c_2 + \log(\frac{N_\alpha \sigma_n}{\varepsilon_n})) - c_3 \sigma_n^{-p}\right)$$

and, under assumption (52), it holds

$$\Pi(\|f - f_0\|_\infty \leq \varepsilon_n) \geq \exp\left(-N_\beta(c_2 + \log(\frac{N_\alpha \sigma_n}{\varepsilon_n})) - c_3 \sigma_n^{-p}\right).$$

The latter is lower bounded by $\exp(-c'_4 n \varepsilon_n^2)$ provided ε_n satisfies (53) and

$$\varepsilon_n \gtrsim n^{-1/2} \sigma_n^{-p/2}. \quad (64)$$

Combining, to have the desired prior mass bound, it suffices that (52), (53) and (64) hold.

We optimize the choice of σ_n based on (52), (64), and then check that (53) also holds. Since (52), (64) imply that

$$\varepsilon_n \gtrsim \{\sigma_n N_\alpha \log^{1/q} n\} \vee \{n^{-1/2} \sigma_n^{-p/2}\} \asymp \{\sigma_n N_\alpha \log^{1/q} n\} \vee \{\varepsilon_n^* N_\beta^{-1/2} \sigma_n^{-p/2}\}, \quad (65)$$

where the first term in the maximum improves with a faster decay of σ_n while the second deteriorates, we choose σ_n to balance the two terms, resulting in

$$\sigma_n \asymp (\varepsilon_n^*)^{\frac{2}{2+p}} N_\alpha^{-\frac{2}{2+p}} N_\beta^{-\frac{1}{2+p}} \log^{-\frac{2}{q(2+p)}} n.$$

This results in

$$\varepsilon_n \gtrsim (\varepsilon_n^*)^{\frac{2}{2+p}} N_\alpha^{\frac{p}{2+p}} N_\beta^{-\frac{1}{2+p}} \log^{\frac{p}{q(2+p)}} n = \varepsilon_n^* N_\alpha^{\frac{p}{2+p}} N_\beta^{\frac{\beta p - 1}{2+p}} \log^{\frac{p}{q(2+p)}} n.$$

This ε_n also satisfies (53) (e.g. since $\beta, p > 1$, which guarantees that ε_n polynomially slower than ε_n^*), as required.

Choosing $\sigma_n = \varepsilon_n^+ / (N_\alpha \log^{1/q} n)$, for $\varepsilon_n^+ = n^{-2/5}$ (the minimax rate for $\beta = 2$), again makes the first term trivial and gives rise to the constraint $\varepsilon_n \gtrsim \varepsilon_n^* n^{\frac{p}{5} - \frac{1}{2(1+2\beta)}}$. $\square$

Remark 6. Let $p > 0, \beta \in (0, 2]$ and $0 < \alpha \leq \beta$. Displays (55) and (65), in the proofs of Theorems 4 and 9, respectively, show that for any choice of σ_n , the prior mass bound holds with

$$\varepsilon_n \geq \{\sigma_n N_\alpha \log^{1/q} n\} \vee \{\varepsilon_n^* \left(\frac{N_\beta^{1-\beta}}{\sigma_n} \right)^{p/2}\},$$

for $\beta \in (0, 1 + \frac{1}{1 \vee p}]$, and with

$$\varepsilon_n \geq \{\sigma_n N_\alpha \log^{1/q} n\} \vee \{\varepsilon_n^* N_\beta^{-1/2} \sigma_n^{-p/2}\},$$

for $\beta \in (1 + \frac{1}{1 \vee p}, 2]$ (a regime admissible only for $p > 1$), as long as (53) holds. In particular, for $\beta \in (0, 1 + \frac{1}{1 \vee p}]$, the standard choices $\sigma_n = n^{-1/2}$ and $\sigma_n = n^{-1}$, give rise to the constraints

$$\varepsilon_n \geq n^{\frac{1-2\alpha}{2(1+2\alpha)}} \log^{1/q} n \vee \varepsilon_n^* n^{\frac{p}{2} \left(\frac{1-\beta}{1+2\beta} + \frac{1}{2} \right)}$$

and

$$\varepsilon_n \geq n^{\frac{-2\alpha}{1+2\alpha}} \log^{1/q} n \vee \varepsilon_n^* n^{\frac{p}{2} \left(\frac{1-\beta}{1+2\beta} + 1 \right)},$$

respectively, which are limited by the choice of α (influencing the network's width), even when p is small. Given that the second terms in the maxima tend to ε_n^* for small $p > 0$, the choice $\sigma_n = n^{-1}$ appears to be better, since it improves the first term which is independent of p . Finally, for $p > 1$ and $\beta \in (1 + \frac{1}{1 \vee p}, 2]$, the standard choices $\sigma_n = n^{-1/2}$ and $\sigma_n = n^{-1}$, give rise to the constraints

$$\varepsilon_n \geq n^{\frac{1-2\alpha}{2(1+2\alpha)}} \log^{1/q} n \vee \varepsilon_n^* n^{\frac{p}{4} - \frac{\beta}{1+2\beta}}$$

and

$$\varepsilon_n \geq n^{\frac{-2\alpha}{1+2\alpha}} \log^{1/q} n \vee \varepsilon_n^* n^{\frac{p}{2} - \frac{\beta}{1+2\beta}},$$

respectively, which are similarly limited by the choice of α .

Appendix D: Technical lemmas

D.1. Theory on contraction for ρ -posteriors

Given a statistical model $(P_f^{(n)})_{f \in \mathcal{F}}$ and $f_0 \in \mathcal{F}$, define the Kullback–Leibler neighborhood of f_0

$$\mathcal{B}_n(f_0, \varepsilon) := \left\{ f \in \mathcal{F} : \int \log \frac{dP_{f_0}^{(n)}}{dP_f^{(n)}} dP_{f_0}^{(n)} \leq n\varepsilon^2, \int \log^2 \frac{dP_{f_0}^{(n)}}{dP_f^{(n)}} dP_{f_0}^{(n)} \leq n\varepsilon^2 \right\}. \quad (66)$$

[Need ref for these Lemmas]

Lemma 4. Let $(P_f^{(n)})_{f \in \mathcal{F}}$ be a statistical model and assume data generated as $X^{(n)} \sim P_{f_0}^{(n)}$ for some $f_0 \in \mathcal{F}$. Let (ε_n) be a positive sequence such that $\varepsilon_n \rightarrow 0$ and $n\varepsilon_n^2 \rightarrow \infty$ as $n \rightarrow \infty$, suppose Π is a prior distribution on $\mathcal{F}$, satisfying

$$\Pi[\mathcal{B}_n(f_0, \varepsilon_n)] \geq e^{-n\varepsilon_n^2}. \quad (67)$$

Then, for any $\rho \in (0, 1)$, there exists a large enough constant $M > 0$, such that, as $n \rightarrow \infty$,

$$E_{f_0} \Pi_\rho \left[\left\{ f : \frac{1}{n} D_\rho(P_f^{(n)}, P_{f_0}^{(n)}) \leq M\varepsilon_n^2 \right\} \mid X^{(n)} \right] \rightarrow 1.$$

Lemma 5. Let $f_0 \in \mathcal{F}$, $\mathcal{B}_n(f_0, \varepsilon)$ the set in (66) and $(P_f^{(n)})_{f \in \mathcal{F}}$ be the statistical model given

1. either by Equation (7), in which case for any $\rho \in (0, 1)$,

$$\{f \in \mathcal{F} : \|f - f_0\|_2 \leq \varepsilon\} \subset \mathcal{B}_n(f_0, \varepsilon) \quad \text{and} \quad \frac{1}{n} D_\rho(P_f^{(n)}, P_{f_0}^{(n)}) = \frac{\rho}{2(1-\rho)} \|f - f_0\|_2^2.$$

2. or by Equation (20), and then there exists a constant $C > 0$ such that

$$\{f \in \mathcal{F} : \|f - f_0\|_\infty \leq \varepsilon\} \subset \mathcal{B}_n(f_0, C\varepsilon).$$

Further assuming $\|f\|_\infty \vee \|f_0\|_\infty \leq F$, we have, for any $\rho \in (0, 1)$,

$$\frac{1}{n} D_\rho(P_f^{(n)}, P_{f_0}^{(n)}) \geq \frac{\rho}{2} e^{-2F^2\rho(1-\rho)} \|f - f_0\|_{2, P_X}^2.$$

D.2. Lemmas for series priors

In this Section we regroup different Lemmas used in the proofs of Section 2.

Lemma 6. Let $\alpha > 0$ and let N_α be defined in (15). For $n \geq n_0(\alpha)$, we have

$$\prod_{k=1}^{N_\alpha} \frac{k^{1/2+\alpha}}{\sqrt{n}} \geq e^{-(1/2+\alpha)N_\alpha}.$$

Proof. The product in the statement equals $\exp\{(1/2+\alpha) \sum_{k=1}^{N_\alpha} \log k - (N_\alpha/2) \log n\}$. Using a comparison series/integral, the partial sum is bounded from below by $\int_1^{N_\alpha} \log(x) dx = N_\alpha \log N_\alpha - (N_\alpha - 1)$ by integration by parts. On the other hand,

$$-(N_\alpha/2) \log n = -\frac{1+2\alpha}{2} N_\alpha \log \left(n^{\frac{1}{2\alpha+1}} \right) \geq -\left(\frac{1}{2} + \alpha\right) N_\alpha \log(N_\alpha + 1).$$

Combining the previous two bounds gives, using the inequality $\log(1+x) \leq x$,

$$\begin{aligned} (1/2 + \alpha) \sum_{k=1}^{N_\alpha} \log k - (N_\alpha/2) \log n &\geq (1/2 + \alpha) \left[-N_\alpha \log(1 + N_\alpha^{-1}) - (N_\alpha - 1) \right] \\ &\geq -(1/2 + \alpha) N_\alpha \end{aligned}$$

Taking exponentials on both sides gives the result. $\square$

Lemma 7. *Let $\alpha, \beta, p > 0$ with $p \geq 2$ and $\alpha \geq \beta$. Let γ and N_γ be defined in (12) and (15) respectively. For $n \geq n_0(\alpha, \beta, p)$, we have*

$$\prod_{k=1}^{N_\gamma} \frac{k^{1/2+\alpha}}{\sqrt{n}} \geq e^{-(\gamma-\alpha)-(1/2+\alpha)N_\gamma - N_\gamma^{1+p(\alpha-\beta)}}.$$

Proof. Proceeding as in the proof of Lemma 6, the logarithm of the product in the display of the Lemma is bounded from below by, taking n large enough so that $N_\gamma \geq 1$,

$$\begin{aligned} & (1/2 + \alpha)\{N_\gamma \log N_\gamma - (N_\gamma - 1)\} - N_\gamma(1/2 + \gamma) \log(N_\gamma + 1) \\ & \geq (1/2 + \alpha)\{N_\gamma \log N_\gamma - N_\gamma\} + 1/2 + \alpha - N_\gamma(1/2 + \gamma) \log N_\gamma - N_\gamma(1/2 + \gamma) \log(1 + N_\gamma^{-1}). \end{aligned}$$

The last term is bounded from below by $-(1/2 + \gamma)$, using $\log(1 + x) \leq x$ for $x > 0$. Regrouping the terms, the last display is further bounded from below by

$$-(\gamma - \alpha) - (\gamma - \alpha)N_\gamma \log N_\gamma - (1/2 + \alpha)N_\gamma.$$

With $(\gamma - \alpha)N_\gamma \log(N_\gamma) = N_\gamma \log(N_\gamma^{\gamma-\alpha}) \leq N_\gamma^{1+p(\gamma-\alpha)}$, using $p \geq 2 \geq 1$, $\gamma \geq \alpha$ for $p \geq 2$ and $\log(x) \leq x$ for $x > 0$ the result follows. $\square$

Lemma 8. *Let f_0 be defined by $f_{0,k} = Lk^{-1/2-\beta}$ for $\beta > 0$ and let $\sigma_k = k^{-1/2-\alpha}$ for $\alpha > \beta$. Let $p < 2$ and recall the definitions of γ and N_γ in (12)–(15). Let $n_\gamma := dN_\gamma$ for some constant $d > 0$. In the (projected) Gaussian white noise model (7), define the event*

$$\mathcal{B}_n := \left\{ |\xi_k| \leq \sqrt{2 \log n}, \quad \text{for all } k = 1, \dots, n_\gamma \right\}. \quad (68)$$

We have $P_{f_0}^{(n)}(\mathcal{B}_n) \rightarrow 1$, as $n \rightarrow \infty$. Also, for all $k = 1, \dots, n_\gamma$ and large enough n , the following hold on $\mathcal{B}_n$:

1. $f_{0,k}/2 \leq X_k \leq 3f_{0,k}/2$,
2. $L\sigma_k \leq 2X_k$,
3. for any constant $M = M(p)$, one can choose $d > 0$ small enough in $n_\gamma := dN_\gamma$, such that

$$nX_k^2 \geq M \left(\frac{X_k}{\sigma_k} \right)^p.$$

Proof. A union bound directly shows that $P_{f_0}^{(n)}[\mathcal{B}_n^c] = o(1)$. Also, by definition of $f_{0,k}$ and σ_k the second point $L\sigma_k \leq 2X_k$ immediately follows from the first. Let us check the latter. For $k \leq n_\gamma$ and n large enough, we have

$$f_{0,k} = Lk^{-\beta-\frac{1}{2}} \geq Ln_\gamma^{-\beta-\frac{1}{2}} \geq 2\sqrt{\frac{2 \log n}{n}}.$$

Therefore, on the event $\mathcal{B}_n$,

$$\frac{f_{0,k}}{2} \leq f_{0,k} - \sqrt{\frac{2 \log n}{n}} \leq X_k \leq f_{0,k} + \sqrt{\frac{2 \log n}{n}} \leq \frac{3}{2}f_{0,k}.$$

We are left to check the last point, since $p < 2$, on $\mathcal{B}_n$ we can lower bound, for n large enough,

$$X_k^{2-p} \sigma_k^p \geq 2^{p-2} f_{0,k}^{2-p} \sigma_k^p = (L/2)^{2-p} k^{-(1+2\beta+p(\alpha-\beta))} = (L/2)^{2-p} k^{-1-2\gamma}.$$

Recalling $n_\gamma = dn^{1/(1+2\gamma)}$, for any constant $M = M(p)$ and $k \leq n_\gamma$, one can choose $d > 0$ small enough, such that

$$X_k^{2-p} \sigma_k^p \geq (L/2)^{2-p} d^{-1-2\gamma} n^{-1} \geq Mn^{-1}. \quad \square$$

Lemma 9 (Bounds for the p -exponential distribution). *Let h_p be the density function defined in (3) and let $\bar{H}_p : x \mapsto \int_0^x h_p(t) dt$ be the associated survival function. Denote Z_p the normalizing constant, such that*

$$h_p(t) = \frac{1}{Z_p} \exp \left\{ \frac{|t|^p}{p} \right\}.$$

Direct computation shows that, for Γ for the usual Gamma function, we have

$$Z_p = 2p^{1/p-1} \Gamma(1/p).$$

For any $p > 0$ and $t \in \mathbb{R}$, we have

$$\frac{1}{2\sqrt{2\pi}} \sqrt{p} e^{1/p-1/2} e^{-\frac{|t|^p}{p}} \leq h_p(t) \leq \frac{1}{2\sqrt{2\pi}} \sqrt{p} e^{1/p} e^{-\frac{|t|^p}{p}}.$$

Also, provided $0 < p < 1$, we have, for any $x \geq 1$,

$$\bar{H}_p(x) \leq \frac{1}{2\sqrt{2\pi}} \frac{e^{1/p}}{\sqrt{p}} x^{1-p} e^{-\frac{|x|^p}{p}}.$$

Proof of Lemma 9. From the Stirling approximation of the Gamma function, the following inequalities are available for any $z > 0$ (see Error bounds and exponential improvements for the asymptotic expansions of the gamma function and its reciprocal)

$$\sqrt{\frac{2\pi}{z}} \left(\frac{z}{e}\right)^z \leq \Gamma(z) \leq \sqrt{\frac{2\pi}{z}} \left(\frac{z}{e}\right)^z e^{\frac{1}{12z}},$$

direct algebraic manipulation gives inequalities for Z_p ,

$$2\sqrt{2\pi} \frac{e^{-1/p}}{\sqrt{p}} \leq Z_p \leq 2\sqrt{2\pi} \frac{e^{p/12-1/p}}{\sqrt{p}}, \quad (69)$$

which leads to the required bounds on h_p . For the second part of the Lemma, using the definition of $\bar{H}_p(x)$ and the change of variable $t = (pu)^{1/p}$, we get

$$Z_p \cdot \bar{H}_p(x) = p^{1/p-1} \int_{x^p/p}^{\infty} u^{1/p-1} e^{-u} du = p^{1/p-1} I\left(\frac{1}{p}, \frac{x^p}{p}\right), \quad (70)$$

where for any $z > 0$ and $s > 0$, we define

$$I(s, z) := \int_z^{\infty} e^{-t} t^{s-1} dt.$$

We prove now that if $s > 1$ and $z \geq s$, we have

$$I(s, z) \leq sz^{s-1}e^{-z}. \quad (71)$$

Indeed, from the change of variables $t = (u + 1)z$, we have

$$I(s, z) = e^{-z}z^s \int_0^\infty e^{-uz}(u+1)^{s-1} du.$$

Using in succession the inequalities $(u+1)^{s-1} \leq e^{u(s-1)}$ and $(1-s)s \geq (1-s)z$, available for $s > 1$ and $z \geq s$, we get

$$\int_0^\infty e^{-uz}(u+1)^{s-1} du \leq \int_0^\infty e^{-uz}e^{u(s-1)} du = (z - (s-1))^{-1} \leq sz^{-1}.$$

Combining this inequality with the previous change of variables yields the desired bound on $I(s, z)$. Provided $x \geq 1$, such that $x^p/p \geq 1/p$ inequality (71) applied with $s = 1/p > 1$ and $z = x^p/p$ in (70) leads to

$$Z_p \cdot \bar{H}_p(x) \leq p^{-1}p^{1/p-1}(x^p/p)^{1/p-1}e^{-x^p/p} = p^{-1}x^{1-p}e^{-x^p/p},$$

Using (69) the lower bound on Z_p gives the desired result. $\square$

D.3. Piecewise affine approximation by SNN

Lemma 10. *Let $f \in \mathcal{H}^\beta(L)$ for some $L > 0$ and $\beta \in (0, 2]$. For any integer $M \geq 1$, consider*

$$\begin{cases} w_0 := M \left[f\left(\frac{1}{M}\right) - f(0) \right], \\ w_k := M \left[f\left(\frac{k+1}{M}\right) - 2f\left(\frac{k}{M}\right) + f\left(\frac{k-1}{M}\right) \right] \text{ for } k = 1, \dots, M-1 \end{cases}$$

and

$$f_M : x \mapsto f(0) + \sum_{k=0}^{M-1} w_k \left(x - \frac{k}{M} \right)_+.$$

Then f_M is a shallow network as in (23) (by construction), such that:

- f_M coincides with the piecewise affine function f_M^* changing slope and interpolating f precisely at k/M , $k = 0, \dots, M$;
- $|w_0| \leq LM^{(1-\beta)_+}$ and $|w_k| \leq 2LM^{1-\beta}$ for $k = 1, \dots, M-1$;
- $f_M \in NN_1(M, 2LM^{(1-\beta)_+} \vee 1)$;
- f_M approximates f uniformly

$$\sup_{x \in [0,1]} |f(x) - f_M(x)| \leq 2LM^{-\beta}.$$

Proof. It is clear from its definition that f_M is a shallow ReLU network of width M as in equation (23). We next show that f_M coincides with the piecewise affine function f_M^* changing slope and

interpolating f precisely at k/M , $k = 0, \dots, M$. Indeed, the latter is such that

$$\begin{aligned} f_M^*(x) &= M \left[f\left(\frac{k+1}{M}\right) - f\left(\frac{k}{M}\right) \right] x + (k+1)f\left(\frac{k}{M}\right) - kf\left(\frac{k+1}{M}\right), \\ &= f\left(\frac{k}{M}\right) + M \left[f\left(\frac{k+1}{M}\right) - f\left(\frac{k}{M}\right) \right] \left(x - \frac{k}{M}\right)_+, \\ x \in I_k &:= \left[\frac{k}{M}, \frac{k+1}{M} \right), \quad k = 0, \dots, M-1. \end{aligned}$$

We proceed by induction. For $k = 1$, that is for $x \in I_0$, it is immediate that

$$f_M^*(x) = f(0) + w_0 x_+,$$

where the right hand side above coincides with f_M for $x \in I_0$. Suppose that $f_M^* = f_M$ on I_{n-1} , $n < M$. Then for $x \in I_n = \left[\frac{n}{M}, \frac{n+1}{M} \right)$,

$$\begin{aligned} f_M(x) &= f(0) + \sum_{k=0}^{M-1} w_k \left(x - \frac{k}{M}\right)_+ = f(0) + \sum_{k=0}^n w_k \left(x - \frac{k}{M}\right)_+ \\ &= f(0) + \sum_{k=0}^{n-1} w_k \left(x - \frac{k}{M}\right)_+ + w_n \left(x - \frac{n}{M}\right)_+ \\ &= f(0) + \sum_{k=0}^{n-1} w_k \left(x - \frac{n-1}{M} + \frac{n-1}{M} - \frac{k}{M}\right) + w_n \left(x - \frac{n}{M}\right) \\ &= f(0) + \sum_{k=0}^{n-1} w_k \left(\frac{n-1}{M} - \frac{k}{M}\right) + \sum_{k=0}^{n-1} w_k \left(x - \frac{n-1}{M}\right) + w_n \left(x - \frac{n}{M}\right) \\ &= f_M\left(\frac{n-1}{M}\right) + \sum_{k=0}^{n-1} w_k \left(x - \frac{n-1}{M}\right) + w_n \left(x - \frac{n}{M}\right) \\ &= f\left(\frac{n-1}{M}\right) + \sum_{k=0}^n w_k \left(x - \frac{n}{M}\right) + \sum_{k=0}^{n-1} \frac{w_k}{M}, \end{aligned}$$

where in the last equality we used the induction hypothesis for $x = \frac{n-1}{M} \in I_{n-1}$ and the fact that f_M^* interpolates f . Noticing that for all $m \geq 1$

$$\sum_{k=0}^m \frac{w_k}{M} = f\left(\frac{m+1}{M}\right) - f\left(\frac{m}{M}\right),$$

we get

$$f_M(x) = f\left(\frac{n}{M}\right) + M \left[f\left(\frac{n+1}{M}\right) - f\left(\frac{n}{M}\right) \right] \left(x - \frac{n}{M}\right) = f_M^*(x), \quad \forall x \in I_n,$$

and the claim is proved.

The bounds on w_k , $k = 0, \dots, M-1$ for $\beta \in (0, 1]$ follow from the definition of $\mathcal{H}^\beta(L)$, for $k = 0$ directly, while for $k \geq 1$ after an application of the triangle inequality. For $\beta \in (1, 2]$, using the triangle inequality and the mean value theorem, there exist $c \in [0, \frac{1}{2M}]$ and $d \in [\frac{1}{2M}, \frac{1}{M}]$, such that

$$|w_0| = M \left| f\left(\frac{1}{M}\right) - f\left(\frac{1}{2M}\right) + f\left(\frac{1}{2M}\right) - f(0) \right| = M \left| \frac{f'(c)}{2M} + \frac{f'(d)}{2M} \right| \leq \frac{L}{2} \leq L,$$

where we have used the definition of $\mathcal{H}^\beta(L)$ to upper bound the derivative. For $k = 1, \dots, M-1$, again using the mean value theorem, there exist $c_k \in I_k, d_k \in I_{k-1}$, such that

$$\begin{aligned} |w_k| &= M \left| f\left(\frac{k+1}{M}\right) - f\left(\frac{k}{M}\right) - \left[f\left(\frac{k}{M}\right) - f\left(\frac{k-1}{M}\right) \right] \right| \\ &= M \left| \frac{f'(c_k)}{M} - \frac{f'(d_k)}{M} \right| \leq L \left(\frac{2}{M} \right)^{\beta-1} \leq 2LM^{1-\beta}, \end{aligned}$$

where in the first upper bound we have used the definition of $\mathcal{H}^\beta(L)$, while in the second the fact that $\beta \leq 2$.

The assumption $f \in \mathcal{H}_\beta(L)$ also implies $|f(0)| \leq L$, while $|k/M| \leq 1$ for all $k \in 0, \dots, M-1$, hence, combining with the bounds on $|w_k|$ we get $f_M \in NN_1(M, 2LM^{(1-\beta)+} \vee 1)$, as claimed.

For the last uniform approximation bound, fix $x \in [0, 1]$ and let k be the unique index among $\{0, \dots, M-1\}$ such that $x \in I_k$. Using that $f_M = f_M^*$, we have

$$|f(x) - f_M(x)| = \left| f(x) - f\left(\frac{k}{M}\right) - M \left[f\left(\frac{k+1}{M}\right) - f\left(\frac{k}{M}\right) \right] \left(x - \frac{k}{M}\right) \right|. \quad (72)$$

If $\beta \in (0, 1]$, using the definition of $\mathcal{H}_\beta(L)$, we have

$$|f(x) - f_M(x)| \leq \left| f(x) - f\left(\frac{k}{M}\right) \right| + M \left| f\left(\frac{k+1}{M}\right) - f\left(\frac{k}{M}\right) \right| \left| x - \frac{k}{M} \right| \leq 2LM^{-\beta},$$

since $|x - k/M| \leq M^{-1}$. For $\beta \in (1, 2]$, using the mean value theorem, there exist $c_k, d_k \in I_k$ such that

$$f(x) = f\left(\frac{k}{M}\right) + f'(c_k) \left(x - \frac{k}{M}\right) \quad \text{and} \quad f\left(\frac{k+1}{M}\right) = f\left(\frac{k}{M}\right) + \frac{1}{M} f'(d_k).$$

Plugging the last two identities into (72), using again the definition of $\mathcal{H}_\beta(L)$ and the fact that $|x - k/M| \leq M^{-1}$, we again get the bound

$$|f(x) - f_M(x)| = |f'(c_k) - f'(d_k)| \left| x - \frac{k}{M} \right| \leq LM^{1-\beta} \left| x - \frac{k}{M} \right| \leq LM^{-\beta} \leq 2LM^{-\beta}.$$

□