

Sharp Capacity Thresholds in Linear Associative Memory: From Winner-Take-All to Listwise Retrieval

Nicholas Barnfield¹ Juno Kim² Eshaan Nichani³
 Jason D. Lee² Yue M. Lu^{1*}

¹Harvard University ²University of California, Berkeley ³Princeton University

May 7, 2026

Abstract

How many key-value associations can a $d \times d$ linear memory store? We show that the answer depends not only on the d^2 degrees of freedom in the memory matrix, but also on the retrieval criterion. In an isotropic Gaussian model for the stored pairs, we show that top-1 retrieval, where every signal must beat its largest distractor, requires the logarithmic model-size scale $d^2 \asymp n \log n$. We prove that the correlation matrix memory construction, which stores associations by superposing key-target outer products, achieves this scale through a sharp phase transition, and that the same scaling is necessary for any linear memory. Thus the logarithm is the intrinsic extreme-value price of winner-take-all decoding.

We next consider listwise retrieval, where the correct target need not be the unique top-scoring item but should remain among the strongest candidates. To formalize this regime, we propose the Tail-Average Margin (TAM), a convex upper-tail criterion that certifies inclusion of the correct target in a controlled candidate list. Under this listwise retrieval criterion, the capacity follows the quadratic scale $d^2 \asymp n$. At load $n/d^2 \rightarrow \alpha$, we develop an exact asymptotic theory for the TAM empirical-risk minimizer through a two-parameter scalar variational principle. The theory has a rich phenomenology: in the ridgeless limit it yields a closed-form critical load separating satisfiable and unsatisfiable phases, and it predicts the limiting laws of true scores, competitor scores, margins, and percentile profiles. Finally, a small-tail extrapolation further leads to the conjectural sharp top-1 threshold $d^2 \sim 2n \log n$.

Contents

1	Introduction	3
1.1	Summary of main results	4
1.2	Related Work	6

*Correspondence: yuelu@seas.harvard.edu.

2	Top-1 retrieval and the logarithmic barrier	9
2.1	Achievability via correlation matrix memory	9
2.2	A universal upper bound for top-1 retrieval	11
3	Listwise retrieval and the Tail-Average Margin	14
3.1	The Tail-Average Margin functional	14
3.2	Learning under the TAM criterion	15
3.3	Performance metrics	17
4	Asymptotic theory of TAM	18
4.1	Gradient and Hessian of the TAM loss	19
4.2	Leave-one-out reduction	21
4.2.1	LOO Hessian and rank-one reduction	21
4.2.2	Justification of Result 1	22
4.3	Scalar variational prediction	24
4.4	Justification of Result 2	26
4.4.1	Gaussian law for competitor scores	27
4.4.2	Proximal law for diagonal scores	28
4.4.3	Finite-dimensional scalar equations and variational consistency	30
5	Ridgeless limit and phase transition	32
5.1	Phase transition	33
5.2	Score, margin, and percentile profiles	36
5.2.1	The SAT phase	36
5.2.2	The UNSAT phase	39
5.3	Numerical validation	40
5.4	A small-tail conjecture for top-1 retrieval	43
6	Conclusion	44
A	Correlation matrix memory	50
A.1	A single-coordinate law	50
A.2	Joint distribution of two columns	51
A.3	Proof of Proposition 1	55
A.4	Percentile law for correlation matrix memory	64
B	Auxiliary estimates for the universal top-1 bound	71
C	Taylor expansion of the interior remainder	73
C.1	Taylor identities	73
C.2	Explicit form of the residual	74
C.3	Entry-size bounds	75
D	A priori bounds on the Frobenius norm of $\widehat{W}$	76
D.1	Upper bound	76
D.2	Lower bound	77

E	A coupling lemma for bilinear Gaussian forms	80
E.1	Deterministic matrix	80
E.2	Random matrix with concentrated Frobenius norm	81
F	A trace estimate for the inverse Hessian	82
G	Variational formulation of the scalar equations	88
G.1	The scalar TAM functional	88
G.2	Moreau envelope identities	89
G.3	Saddle-point equations	89
G.4	Maximization in χ and convexity in ν	90

1 Introduction

Associative memory refers broadly to the ability of a system to store and retrieve relationships between objects. Given a collection of input-output pairs, one would like the system to recover the correct target when presented with a corresponding cue. This basic mechanism appears in many forms, ranging from the first neural network models for pattern learning [Kohonen, 1972, Amari, 1972, Anderson, 1972, Hopfield, 1982, 1984], to modern key-value retrieval systems [Vaswani et al., 2017, Ramsauer et al., 2021, Gershman et al., 2025]. Associative memory has also recently attracted attention as a model of factual recall in large language models [Cabannes et al., 2024a,b, Jiang et al., 2024, Nichani et al., 2024] as trained transformer layers have been observed to function as key-value memories [Geva et al., 2021, Meng et al., 2022, Bietti et al., 2023].

A central theoretical question in associative memory is: given n associations, how large must the memory model be in order to retrieve them under a prescribed criterion? In this paper, we study this question in the simple but already nontrivial setting of *linear associative memory*. Given pairs of key vectors (embeddings) v_i and associated targets (unembeddings) u_i , we consider a matrix memory model $W \in \mathbb{R}^{d \times d}$ and define the score matrix $S = (s_{j,i})_{1 \leq i,j \leq n}$ as

$$s_{j,i} = u_j^\top W v_i, \quad \{(u_i, v_i)\}_{i=1}^n \subset \mathbb{R}^d \times \mathbb{R}^d, \quad W \in \mathbb{R}^{d \times d}.$$

This score matrix encodes the predictions for the target when decoding to a fixed vocabulary of size n . Rather than requiring the output $W v_i$ of the memory model on an input v_i to be exactly its associated target u_i , as in classical models for associative memory [Hopfield, 1982], we will consider methods which compare the score for the true target $s_{i,i}$ to its competitors $(s_{j,i})_{j \neq i}$. One main message of the paper is that the answer to the above question depends sharply on the retrieval criterion: as we will see next, controlling the single largest distractor is fundamentally more expensive than controlling the average strength of a fixed upper tail.

A standard baseline in the associative memory literature is *top-1 retrieval*, also called exact retrieval [Kohonen, 1972, Cabannes et al., 2024a,b, Nichani et al., 2024]. Under this criterion, each key must retrieve its matched target with the largest score:

$$s_{i,i} > s_{j,i}, \quad \forall j \neq i, \quad \forall i \in [n]. \quad (1)$$

Operationally, this is the natural criterion for final winner-take-all decoding. It is also the hard-attention, or low-temperature, limit of softmax attention in transformer-based architectures [Vaswani et al., 2017]; at fixed temperature, the same concentration occurs when the leading score gap is large.

We also study a complementary retrieval regime: *listwise recoverability*. In many memory architectures, a memory access is an intermediate computation rather than a final decision; its role is to keep the correct association salient among high-scoring candidates for subsequent layers, heads, or rerankers to refine. The most direct listwise condition would require the signal to beat the k -th largest competitor. This rank condition is non-convex in the score matrix. We instead use a convex certificate [Rockafellar and Uryasev, 2000, 2002]: we compare the signal score with the upper-tail average of its competitors. For query i , let

$$s_{(1),i} \geq s_{(2),i} \geq \cdots \geq s_{(n-1),i}$$

denote the ordered competing scores, and set $k = \lceil r(n-1) \rceil$ for a tail fraction r . We call the resulting upper-tail comparison the *Tail-Average Margin* (TAM) criterion:

$$s_{i,i} > \text{TAM}_r((s_{j,i})_{j \neq i}) := \frac{1}{k} \sum_{\ell=1}^k s_{(\ell),i}, \quad \forall i \in [n]. \quad (2)$$

In other words, for each query, the target score $s_{i,i}$ is required to beat the average of its top- k competitors. Since $\text{TAM}_r((s_{j,i})_{j \neq i}) \geq s_{(k),i}$, a positive TAM margin certifies that the matched target lies in the top- k candidate list. The parameter r tunes the retrieval statistic: when $r = 1/(n-1)$, TAM reduces to top-1 retrieval, while $r = 1$ compares the signal with the average competitor score. Thus top-1 retrieval controls the single largest distractor, whereas TAM controls the average strength of the upper tail of the competitor distribution.

This listwise perspective is in the same spirit as list decoding in coding theory, where the decoder is allowed to retain a controlled list of plausible messages rather than make a unique decision [Elias, 1957, Guruswami, 2007]. It is also aligned with listwise ranking losses in information retrieval, which optimize criteria defined on ranked candidate lists rather than on a single pairwise comparison [Cao et al., 2007].

For the two criteria studied in this paper—top-1 retrieval and TAM—we ask how large the ambient dimension, and hence the d^2 parameters of the linear memory, must be in order to retrieve n stored associations. In full generality, the answer depends on the geometry of the stored vectors and on the rule used to construct the memory W . To obtain a sharp high-dimensional theory, we study the isotropic Gaussian model

$$u_i, v_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, I_d/d), \quad i \in [n]. \quad (3)$$

This model isolates the geometric effect of high-dimensional interference, without imposing additional structure on the stored associations.

1.1 Summary of main results

Top-1 retrieval. For top-1 retrieval (1), we rigorously prove in Section 2 that the required model size is governed by the scale

$$d^2 \asymp n \log n. \quad (4)$$

The positive direction builds on the correlation matrix memory construction, which stores the associations as a superposition of outer products [Kohonen, 1972, Elhage et al., 2022]. Previous work of Nichani et al. [2024] showed that this special memory achieves top-1 retrieval with high probability when $d^2 \geq Cn(\log n)^4$ for a sufficiently large constant C . We improve this result by proving a sharp transition for correlation matrix memory: Proposition 1 shows that it succeeds when $d^2 > (8 + \epsilon)n \log n$ and fails when $d^2 < (8 - \epsilon)n \log n$.

The main top-1 result is the negative direction. We prove that the logarithmic factor is fundamental, not an artifact of the correlation matrix construction: Theorem 1(ii) shows that, for any $C < 2/\pi$, if $d^2 \leq Cn \log n$, then no memory $W \in \mathbb{R}^{d \times d}$ can attain top-1 retrieval with high probability. Thus the logarithmic factor in (4) is unavoidable: it is the inherent price of asking every signal to beat the largest element in a cloud of $n - 1$ competitors.

Tail-Average Margin. For the listwise criterion (2), the picture changes qualitatively. TAM replaces the maximum competitor, governed by extreme-value fluctuations, by the average of the largest $k = \lceil r(n - 1) \rceil$ competing scores. This upper-tail average is no longer controlled by a single maximum when the tail fraction r is fixed, and hence removes the bottleneck responsible for the logarithmic factor in top-1 retrieval, while still certifying that the correct target lies in a controlled top- k candidate list. The natural model-size scale for TAM is therefore

$$d^2 \asymp n.$$

We learn the memory matrix for this criterion through empirical risk minimization. With logistic loss $\ell(t) = \log(1 + e^{-t})$, ridge parameter $\lambda \geq 0$, and the TAM functional TAM_r , the estimator is

$$\widehat{W} \in \arg \min_{W \in \mathbb{R}^{d \times d}} \left\{ \frac{1}{n} \sum_{i=1}^n \ell \left(s_{i,i}(W) - \text{TAM}_r((s_{j,i}(W))_{j \neq i}) \right) + \frac{\lambda}{2d^2} \|W\|_F^2 \right\}.$$

Section 3 gives the precise convex formulation of this objective. The main goal of the TAM theory is to understand the asymptotic behavior of this ERM in the regime

$$n, d \rightarrow \infty, \quad \frac{n}{d^2} \rightarrow \alpha \in (0, \infty), \quad r \in (0, 1) \text{ fixed}.$$

Conditional on the six postulates stated in Section 4, our main asymptotic conclusions are as follows:

1. *Scalar reduction.* The high-dimensional ERM admits a scalar variational description involving only two parameters, stated in Result 2 and justified in Section 4.4;
2. *Score and margin laws.* The scalar description gives limiting predictions for the fitting loss, the laws of the true and competing scores, and the induced margin and percentile profiles, developed in Section 4 and Section 5;
3. *Ridgeless phase transition.* In the ridgeless limit, the scalar theory yields a sharp phase transition at a critical load

$$\alpha_c(r) = \frac{1}{(1 + \kappa_r^2)\Phi(\kappa_r) + \kappa_r\varphi(\kappa_r)}, \quad \kappa_r = \frac{\varphi(\Phi^{-1}(1 - r))}{r}, \quad (5)$$

where φ and Φ denote the standard normal density and distribution function. When the load $\alpha < \alpha_c(r)$, the scalar TAM loss vanishes in the ridgeless limit; for $\alpha > \alpha_c(r)$, the limiting loss remains positive.

4. *Small-tail top-1 conjecture.* Since TAM with $r = 1/(n - 1)$ is exactly top-1 retrieval, the fixed-tail formula also suggests a tantalizing small-tail extrapolation back to the top-1 problem. Formally sending r to $1/(n - 1)$ in the critical-load formula in (5) predicts the exact leading constant in the top-1 retrieval threshold:

$$\frac{d^2}{n \log n} = 2,$$

meaning that top-1 retrieval should be feasible when $d^2 \geq (2 + \epsilon)n \log n$ and infeasible when $d^2 \leq (2 - \epsilon)n \log n$. We state this prediction separately as a conjecture in Section 5.4 and give numerical support there. The same constant is also consistent with a conjecture made in the concurrent work of Giorlandino et al. [2026] using a simplified decoupled model; see the discussion in Section 1.2.

The main technical approach in our work is a leave-one-out analysis. These methods are central to precise asymptotics for high-dimensional M -estimators [El Karoui et al., 2013, Lei et al., 2018], but the present problem has a different geometry. The decision variable W is a high-dimensional matrix rather than a vector; the effective features are rank-one tensors $u_j v_i^\top$ rather than independent covariate vectors; and each stored pair appears in two coupled roles, as its own signal and as a competitor for many other queries. Consequently, removing one pair changes not only one summand of the loss, but also the competitor tails, threshold variables, and Hessian coefficients seen throughout the objective. The leave-one-out analysis developed in Section 4 isolates the resulting rank-one perturbation structure, a mechanism that may be useful for other matrix-valued ERM with shared random features and endogenous listwise weights. Establishing the postulates in Sections 4.2 and 4.3 rigorously is the main remaining step toward a fully unconditional theorem for the TAM optimizer.

1.2 Related Work

Neural associative memory. Associative memory dates back to early neural models that store patterns through correlations in the weights [Amari, 1972, Anderson, 1972, Nakano, 1972, Kohonen, 1972, Hopfield, 1982, 1984]. The Hopfield network [Hopfield, 1982] is the symmetric case, where the goal is to memorize and retrieve the patterns themselves. We focus instead on associative recall: a key should retrieve its paired target, and the two need not coincide [Amari, 1972]. In particular, Kohonen [1972] define correlation matrix memory as a sum of rank-one key-target outer products, the construction analyzed in Section 2.1. Bidirectional associative memory is another paired-pattern model based on outer-product updates [Kosko, 1988].

Modern versions of neural networks with built-in associative memory capabilities include the neural Turing machine [Graves et al., 2014], memory networks [Weston et al., 2015], dense associative memory [Krotov and Hopfield, 2016, Lucibello and Mézard, 2024], MCHN [Ramsauer et al., 2021], MCSDM [Kanerva, 1988, Bricken and Pehlevan, 2021], and universal

Hopfield networks [Millidge et al., 2022]. Binary variants of matrix memory have also been studied in the literature [Willshaw et al., 1969, Gardner, 1987, Hobson, 2011].

Transformers and factual recall. Transformer models trained on factual recall tasks have been empirically observed to implement associative memory through their weight matrices. Feedforward as well as attention layers have been shown to behave like key-value memories [Geva et al., 2021, Bietti et al., 2023], and linear-memory viewpoints have been used to study factual editing, induction head tasks, and latent space retrieval [Meng et al., 2022, Bietti et al., 2023, Jiang et al., 2024]. On the theory side, Nichani et al. [2024] study a sequence-based factual recall task where one token functions as the key, and show that a one-layer transformer solves the task with capacity scaling linearly with parameter count, up to logarithmic factors. This storage scaling has also been empirically observed in large language models [Allen-Zhu and Li, 2024]. Related theoretical analyses include optimization dynamics for a single-layer transformer on a token recall task [Vural et al., 2026], and a measure-theoretic formulation of transformers with implicit associative memory mechanisms [Kawata and Suzuki, 2026].

Storage capacity. Nichani et al. [2024] study the capacity of linear and MLP associative memories with random spherical embeddings, showing in particular that correlation matrix memory succeeds when $d^2 \gtrsim n(\log n)^4$. Cabannes et al. [2024a] propose the exact retrieval criterion and study correlation matrix memory under a weighted data distribution. From an optimization perspective, Cabannes et al. [2024b] study gradient descent under a multiclass logistic formulation, and Kim et al. [2026] compare the capacity of memory matrices obtained by different optimizers under Gaussian embeddings. Compared with these works, our top-1 result identifies the exact statistical logarithmic barrier $d^2 \asymp n \log n$ for arbitrary linear memories. We also introduce the TAM criterion and, to our knowledge, give the first capacity theory for linear associative memory under a listwise retrieval rule.

We also briefly mention that storage capacity has been studied in detail for the Hopfield network and its variants. The capacity $d \asymp n$ of Hopfield networks is classical, see e.g., McEliece et al. [1987], Folli et al. [2017] and the works cited within, while modern extensions such as dense associative memory have been shown to achieve exponential capacity [Demircigil et al., 2017, Ramsauer et al., 2021, Lucibello and Mézard, 2024]. The scaling in these works differ from our results since we decode to a fixed vocabulary of size n , while the classical Hopfield formulation decodes to the hypercube.

Concurrent work. Concurrently with this work, Giorlandino et al. [2026] independently studied the capacity of linear associative memory under the top-1 retrieval criterion through a decoupled model. In this simplified formulation, the output embeddings $\{u_j\}_{j \leq n}$ are re-sampled across constraints, thus removing the statistical dependence structure among the retrieval inequalities in (1). Based on numerical evidence comparing training and test errors, and score distributions, they conjecture that the original and decoupled problems have the same critical capacity. For the decoupled model, they carry out a Gardner-type statistical physics calculation and predict the critical threshold $\frac{n \log n}{d^2} = \frac{1}{2}$ for top-1 retrieval. This provides independent support for the small-tail extrapolation of our TAM theory in Sec-

tion 5. They further extend both the decoupling conjecture and the capacity calculation to rank-constrained memories, corresponding to two-layer linear MLPs.

Classification and leave-one-out methods. The TAM learning problem is also connected to high-dimensional classification through its logistic loss. The classical work of Cover [1965] computed the separability capacity of random patterns, and the perceptron capacity was later studied through statistical physics methods, including margin-constrained variants [Gardner, 1988]. In ordinary unregularized logistic regression, the existence of the maximum likelihood estimator is equivalent to nonseparability of the labeled data; equivalently, the failure of existence is a linear separability, or capacity, event. Sur and Candès [2019] established a high-dimensional phase transition for this event under Gaussian covariates. A broader line of work in statistical physics studies optimal errors and phase transitions in high-dimensional generalized linear models via the approximate message passing algorithm [Barbier et al., 2019, Aubin et al., 2020, Mignacco et al., 2020]. From a technical point of view, our TAM analysis uses a leave-one-out approach, which has played an important role in precise asymptotics for high-dimensional M -estimators [El Karoui et al., 2013, El Karoui, 2018, Lei et al., 2018]. At a conceptual level, this is also the cavity method from spin glass theory [Mézard et al., 1987].

The present problem differs from standard high-dimensional regression and classification in several ways. The unknown is a matrix rather than a vector; the effective covariates are rank-one tensors $u_j v_i^\top$ with strong dependencies; each sample contributes both a signal score and a full column of competitor scores; and the TAM threshold is endogenous, selected by the learned scores themselves. These couplings prevent existing leave-one-out theories from applying as black boxes; the resulting cavity calculation is matrix-valued, listwise, and self-consistent in a way that does not arise in standard high-dimensional regression.

Notation

We write

$$U = [u_1 \cdots u_n] \in \mathbb{R}^{d \times n} \quad \text{and} \quad V = [v_1 \cdots v_n] \in \mathbb{R}^{d \times n}$$

for the target and key matrices, respectively. Under the Gaussian model (3), $\sqrt{d}U$ and $\sqrt{d}V$ are independent standard Gaussian matrices in $\mathbb{R}^{d \times n}$. $\text{vec}(\cdot)$ denotes column-wise vectorization. We write $X \stackrel{(d)}{=} Y$ when two random variables, vectors, or matrices have the same distribution.

We use stochastic domination [Erdős and Yau, 2017] to record high-probability bounds uniformly over polynomial-size index sets. Let $\{X_\alpha^{(n)}\}_{\alpha \in \mathcal{I}_n}$ be nonnegative random variables, where $\mathcal{I}_n$ may depend on n and has polynomial cardinality, and let $Y_n \geq 0$ be deterministic. We write

$$X_\alpha^{(n)} = \mathcal{O}_{\prec}(Y_n) \quad \text{uniformly in } \alpha \in \mathcal{I}_n$$

if, for every $\varepsilon > 0$ and $D > 0$,

$$\mathbb{P} \left(\max_{\alpha \in \mathcal{I}_n} X_\alpha^{(n)} > n^\varepsilon Y_n \right) \leq n^{-D}$$

for all sufficiently large n . When $\mathcal{I}_n$ is a singleton, we suppress the uniformity qualifier.

2 Top-1 retrieval and the logarithmic barrier

Throughout this section, we work under the Gaussian model (3) and the top-1 retrieval criterion (1) of Section 1. For a weight matrix $W \in \mathbb{R}^{d \times d}$, define the top-1 retrieval event

$$\text{PR}_n(W) := \bigcap_{i=1}^n \left\{ u_i^\top W v_i > u_j^\top W v_i \text{ for all } j \neq i \right\},$$

and write

$$\text{PR}_n := \left\{ (U, V) : \exists W \in \mathbb{R}^{d \times d} \text{ such that } \text{PR}_n(W) \text{ holds} \right\}.$$

The event PR_n asks whether the observed random instance admits some matrix W satisfying all top-1 retrieval inequalities simultaneously. This is a pointwise feasibility statement for the realized data: the matrix W may depend arbitrarily on the full instance (U, V) . The main result of this section shows that this event has a logarithmic model-size threshold.

Theorem 1 (Top-1 retrieval threshold). *There exist absolute constants $\varrho_1 > \varrho_2 > 0$ such that, along any sequence $n, d \rightarrow \infty$:*

- (i) (*Achievability.*) *If $d^2 \geq \varrho_1 n \log n$, then $\mathbb{P}(\text{PR}_n) \rightarrow 1$.*
- (ii) (*Impossibility.*) *If $d^2 \leq \varrho_2 n \log n$, then $\mathbb{P}(\text{PR}_n) \rightarrow 0$.*

In particular, we may take any $\varrho_1 > 8$ and $\varrho_2 < 2/\pi$.

The two halves of Theorem 1 are established by essentially independent arguments. In Section 2.1, we prove the achievability direction by a direct analysis of the correlation matrix memory construction, and show that this construction exhibits a sharp phase transition around $\varrho_1 = 8$. In Section 2.2, we prove the impossibility direction for all memories $W \in \mathbb{R}^{d \times d}$, by contrasting the average scale of the target scores $\text{tr}(S)$ against the lower bound implied by a uniform anticoncentration of the off-diagonal elements in (1).

2.1 Achievability via correlation matrix memory

Recall the Gaussian model (3), and write $U = [u_1 \cdots u_n]$ and $V = [v_1 \cdots v_n]$. Consider the memory

$$W^\times := UV^\top = \sum_{i=1}^n u_i v_i^\top, \tag{6}$$

obtained by summing the rank-one outer products of the stored pairs. This solution is known as *correlation matrix memory* [Kohonen, 1972] and is the direct analogue of the classical self-organizing network [Amari, 1972] for asymmetric key and target pairs. It is also easy to update, requiring only a local (rank-one) correction upon the addition or deletion of an association, and thus can be seen as a form of Hebbian learning [Hebb, 1949].

Clearly, $W^\times$ enables top-1 retrieval if the (un)embeddings are mutually orthogonal, which requires dimensionality to scale as $d \geq n$. However, $W^\times$ can also store associations much more efficiently by storing them non-orthogonally, i.e., in *superposition* [Elhage et al., 2022].

Indeed, [Nichani et al. \[2024, Theorem 1\]](#) showed that $W^\times$ achieves top-1 retrieval with high probability when $d^2 \geq Cn(\log n)^4$ for a sufficiently large constant C .

Here we sharpen this polylogarithmic sufficient condition by determining the exact threshold for correlation matrix memory. In the regime $d^2 \asymp n \log n$, the retrieval probability $\mathbb{P}(\text{PR}_n(W^\times))$ has a sharp phase transition at the constant 8.

Proposition 1 (Retrieval threshold for correlation matrix memory). *Consider the limiting regime*

$$n, d \rightarrow \infty, \quad \frac{d^2}{n \log n} \rightarrow \varrho \quad (7)$$

for some constant $\varrho \in [0, \infty]$. Then it holds that

$$\lim_{d \rightarrow \infty} \mathbb{P}(\text{PR}_n(W^\times)) = \begin{cases} 1 & \text{if } \varrho > 8, \\ 0 & \text{if } \varrho < 8. \end{cases}$$

Proof. See Appendix [A.3](#). □

Since $\text{PR}_n(W^\times) \subseteq \text{PR}_n$, the case $\varrho > 8$ immediately establishes Theorem [1\(i\)](#). We give an overview of the proof strategy below.

Reduction to a single coordinate. Under [\(6\)](#), the score matrix becomes $S = U^\top U V^\top V$ with $s_{j,i} = u_j^\top W^\times v_i$. Define the per-coordinate *retrieval failure event*

$$F_i = \left\{ s_{i,i} \leq \max_{j \neq i} s_{j,i} \right\}, \quad i \in [n]$$

so that $\text{PR}_n(W^\times) = \bigcap_{i=1}^n F_i^c$. Because the joint distribution of (U, V) is invariant under simultaneous permutations of the columns of U and V , the events $F_1, \dots, F_n$ are *exchangeable*; in particular, $\mathbb{P}(F_i) = \mathbb{P}(F_1)$ for every i . A union bound yields the lower estimate

$$\mathbb{P}(\text{PR}_n(W^\times)) \geq 1 - n\mathbb{P}(F_1), \quad (8)$$

and in fact this is nearly tight as the events F_i are only weakly dependent ([Lemma 8](#)). This reduces the behavior of $\mathbb{P}(\text{PR}_n(W^\times))$ to that of the single failure probability $\mathbb{P}(F_1)$.

A single-coordinate law and extreme values. The probability $\mathbb{P}(F_1)$ is determined by the first column of the score matrix. Write

$$\theta := s_{1,1}, \quad \xi_j := s_{j,1}, \quad 2 \leq j \leq n.$$

[Appendix A.1](#) gives an exact distributional representation for $(\theta, \xi_2, \dots, \xi_n)$. Its content is simple: up to chi-distributed normalization factors that concentrate at one (as $n, d \rightarrow \infty$), the signal θ has mean one and Gaussian fluctuations at scale $\sqrt{n/d^2}$, while the competitors $\{\xi_j : 2 \leq j \leq n\}$ are approximately independent centered Gaussians at the same scale. Thus, if $\alpha_n := n/d^2$, the leading-order picture is

$$\theta \approx 1 + \sqrt{\alpha_n} Z_0, \quad \xi_j \approx \sqrt{\alpha_n} Y_j, \quad \text{where } Z_0, Y_1, \dots, Y_{n-1} \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, 1).$$

The signal θ and each competitor ξ_j therefore have the same fluctuation scale, while the signal has deterministic offset 1. Thus, the single-coordinate failure event $F_1 = \{\theta \leq \max_j \xi_j\}$ becomes, at leading order,

$$F_1 \approx \left\{ \max_{j \leq n-1} Y_j - Z_0 \geq \alpha_n^{-1/2} \right\}. \quad (9)$$

The quantity on the left is the maximum of $n - 1$ i.i.d. standard normals minus an independent standard normal, both of which fluctuate. For n large, $\max_{j \leq n-1} Y_j$ concentrates at $\sqrt{2 \log n}$ with Gumbel fluctuations of order $1/\sqrt{2 \log n}$, while Z_0 retains $O(1)$ Gaussian fluctuations. Hence (9) has probability converging to zero precisely when the right-hand threshold $\alpha_n^{-1/2}$ exceeds $\sqrt{2 \log n}$. If $d^2 = \varrho n \log n$, this threshold is $\alpha_n^{-1/2} = \sqrt{\varrho \log n}$, giving the naive constant $\varrho = 2$.

This heuristic argument is almost, but not quite, the right one: applying the union bound in (8) requires the stronger decay $\mathbb{P}(F_1) = o(n^{-1})$ (instead of merely $o(1)$). Meeting this more stringent threshold requires the fluctuations of Z_0 in (9) to enter the tail analysis, and analyzing this refinement yields the correct threshold $\varrho = 8$ in Proposition 1. The details are given in Appendix A.3.

2.2 A universal upper bound for top-1 retrieval

The lower side of Proposition 1 rules out top-1 retrieval for the specific construction $W^\times$ below its sharp logarithmic threshold. It leaves open whether a more clever choice of weight matrix—possibly obtained through a training algorithm—can succeed at the same scale. In this subsection, we show that once n exceeds a constant multiple of $d^2/\log d$, with high probability, *no* matrix $W \in \mathbb{R}^{d \times d}$ can satisfy all top-1 retrieval inequalities. Equivalently, on the scale $d^2 \asymp n \log n$, this gives the model-size lower bound $d^2 \geq (2/\pi - o(1)) n \log n$, which establishes part (ii) of Theorem 1. Combined with Section 2.1, this result pins down the logarithmic barrier for top-1 retrieval. We note that the existence of this logarithmic factor is also suggested in Nichani et al. [2024, Theorem 7] via an information-theoretic argument, however their result only applies to a finite-bit learner.

Theorem 2 (Universal top-1 impossibility bound). *Fix any $\epsilon > 0$. The following holds for all sufficiently large d : if*

$$n \geq \left(\frac{\pi}{4} + \epsilon \right) \frac{d^2}{\log d},$$

and $\{u_i, v_i\}_{i=1}^n$ are i.i.d. $\mathcal{N}(0, \frac{1}{d} I_d)$, then with probability at least $1 - e^{-\Omega(d)}$, there cannot exist a matrix $W \in \mathbb{R}^{d \times d}$ such that

$$u_i^\top W v_i > u_j^\top W v_i \quad \forall i \in [n], j \neq i. \quad (10)$$

More generally, the statement holds (with a potentially different proportionality constant) if

$\{u_i\}_{i=1}^n$ are i.i.d. $\mathcal{N}(0, \frac{1}{d}I_d)$ and $\{v_i\}_{i=1}^n \subseteq \mathbb{R}^d \setminus \{0\}$ satisfy the following:¹

$$\|v_i\| \lesssim 1, \quad \frac{1}{n} \sum_{i=1}^n v_i v_i^\top \approx \frac{1}{d} I_d, \quad \left\| \frac{1}{n} \sum_{i=1}^n v_i^{\otimes 4} \right\| \lesssim \frac{1}{d^2}. \quad (11)$$

This implies Theorem 1(ii). Indeed, if $\varrho < 2/\pi$ in (7), then $d^2 \leq (2/\pi - \delta)n \log n$ for some fixed $\delta > 0$ and all large n . Since $\log n \sim 2 \log d$ on this scale, the hypothesis of Theorem 2 holds for some $\epsilon > 0$, and hence $\mathbb{P}(\text{PR}_n) \rightarrow 0$.

As the proof is short but nontrivial, we present the full argument below. The key point is that, under top-1 retrieval, the trace of the score matrix $S(W) = U^\top W V$ can be rewritten in two ways:

$$\text{tr}(S(W)) = \sum_i u_i^\top W v_i = \sum_i \max_{j \in [n]} u_j^\top W v_i, \quad \text{tr}(S(W)) = \langle W, U V^\top \rangle_{\text{F}}.$$

The first representation forces $\text{tr}(S(W))$ to be large by a uniform anticoncentration principle for the u_i 's (Lemma 1), which is where the logarithmic factor enters. The second representation forces the same quantity to be small. Both bounds depend on W only through $\|W\|_{\text{F}}$, and comparing them yields a uniform obstruction over all possible choices of W .

We first establish some anticoncentration and concentration lemmas. Without loss of generality, we may suppose $n \gtrsim d^2 / \log d$. When needed, we also assume $\frac{1}{2} \leq \|u_i\|, \|v_i\| \leq 2$ which holds w.p. $1 - e^{-\Omega(d)}$.

Lemma 1. *For any fixed $\epsilon > 0$, it holds w.p. $1 - e^{-\Omega(d)}$ that*

$$\inf_{x \in S^{d-1}} \max_{j \in [n]} u_j^\top x \geq \sqrt{\frac{(2 - \epsilon) \log d}{d}}.$$

Proof. See Appendix B. □

Lemma 2. *If $\{v_i\}_{i=1}^n$ are Gaussian, for any fixed $\epsilon > 0$, it holds w.p. $1 - e^{-\Omega(d)}$ that*

$$\inf_{x \in S^{d-1}} \frac{1}{n} \sum_{i=1}^n |x^\top v_i| \geq \sqrt{\frac{2 - \epsilon}{\pi d}}.$$

Moreover, the statement also holds deterministically under (11) (with a potentially different constant factor).

Proof. See Appendix B. □

We proceed to prove Theorem 2. Suppose (10) is true for some weight matrix $W \in \mathbb{R}^{d \times d}$. Setting $x_i := W v_i$, (10) implies that $u_i^\top x_i = \max_{j \in [n]} u_j^\top x_i$ for all i . Applying Lemma 1 to each of the unit vectors $x_i / \|x_i\|$, we obtain

$$u_i^\top W v_i = u_i^\top x_i = \max_{j \in [n]} u_j^\top x_i \geq \sqrt{\frac{(2 - \epsilon) \log d}{d}} \|x_i\|$$

¹The fourth moment bound is an incoherence condition equivalent to $\frac{1}{n} \sum_i \langle v_i, x \rangle^4 \lesssim \frac{\|x\|^4}{d^2}$ by Banach's theorem for symmetric tensors. The necessity of this condition can be seen by taking $\{v_i\}_{i=1}^n$ to be repetitions of the coordinate vectors $\{e_j\}_{j=1}^d$ for which Lemma 2 fails.

simultaneously for all i . Summing over i gives

$$\sum_{i=1}^n u_i^\top W v_i \geq \sqrt{\frac{(2-\epsilon) \log d}{d}} \sum_{i=1}^n \|W v_i\|. \quad (12)$$

We further lower bound the right-hand side in terms of $\|W\|_F$. Write the SVD of W as $W = ASB^\top = \sum_{k=1}^d s_k a_k b_k^\top$ where $A, B \in O(d)$. Then $\|W\|_F^2 = \sum_{k=1}^d s_k^2$ and

$$W v_i = \sum_{k=1}^d s_k (b_k^\top v_i) a_k, \quad \|W v_i\| = \left(\sum_{k=1}^d s_k^2 (b_k^\top v_i)^2 \right)^{1/2}.$$

By Cauchy–Schwarz,

$$\sum_{k=1}^d s_k^2 |b_k^\top v_i| \leq \left(\sum_{k=1}^d s_k^2 \right)^{1/2} \left(\sum_{k=1}^d s_k^2 (b_k^\top v_i)^2 \right)^{1/2} = \|W\|_F \|W v_i\|.$$

Dividing both sides by $\|W\|_F$ and summing over i , by Lemma 2,

$$\begin{aligned} \sum_{i=1}^n \|W v_i\| &\geq \frac{1}{\|W\|_F} \sum_{k=1}^d s_k^2 \sum_{i=1}^n |b_k^\top v_i| \\ &\geq \frac{1}{\|W\|_F} \sum_{k=1}^d s_k^2 \cdot \sqrt{\frac{2-\epsilon}{\pi d}} n = \sqrt{\frac{2-\epsilon}{\pi d}} n \|W\|_F. \end{aligned}$$

Combining this with (12), we obtain

$$\sum_{i=1}^n u_i^\top W v_i \geq \frac{2-\epsilon}{\sqrt{\pi}} \frac{n \sqrt{\log d}}{d} \|W\|_F.$$

On the other hand, by the standard cross-term concentration estimate recorded as Lemma 13 in Appendix B,

$$\sum_{i=1}^n u_i^\top W v_i = \langle W, UV^\top \rangle_F \leq (1+\epsilon) \sqrt{n} \|W\|_F.$$

Comparing the two bounds, we have thus proved that

$$(1+\epsilon) \sqrt{n} \geq \frac{2-\epsilon}{\sqrt{\pi}} \frac{n \sqrt{\log d}}{d} \implies n \leq \left(\frac{1+\epsilon}{2-\epsilon} \right)^2 \frac{\pi d^2}{\log d},$$

and the statement follows by taking sufficiently small ϵ . $\square$

3 Listwise retrieval and the Tail-Average Margin

Top-1 retrieval is a winner-take-all criterion: each query must identify a unique target in a single step. This is natural for final decoding, but many memory modules are used as intermediate stages. They surface a set of plausible associations, or form a mixture of high-scoring memories, which later computation can refine. In this setting, successful retrieval need not mean that the correct target is already ranked first. It is enough that the correct association remains salient among the strongest competitors. This leads to a listwise notion of retrieval, which is the focus of this section.

A direct listwise requirement would ask the signal score to beat a high competitor quantile, thereby ensuring that the correct target belongs to a controlled candidate set. This rank condition is natural, but it is non-convex and combinatorial. We therefore use a convex certificate: we ask the signal score to beat the *average* of the strongest competitors. We call the resulting functional the *Tail-Average Margin* (TAM). TAM is a listwise retrieval criterion in its own right: it controls the mass of strong distractors rather than only the single largest one, while retaining enough convex structure for optimization and asymptotic analysis.

In this section we define the TAM functional, record its basic properties, and formulate the corresponding learning problem and performance criteria for linear associative memory. In the next section, we derive an asymptotic theory of TAM learning in the quadratic regime $d^2 \asymp n$, with a fixed tail fraction $k/(n-1) \rightarrow r \in (0, 1)$.

3.1 The Tail-Average Margin functional

Let $x = (x_1, \dots, x_m) \in \mathbb{R}^m$ and fix a tail fraction $r \in (0, 1]$. Set $k := \lceil rm \rceil$.

Definition 1 (Tail-Average Margin). *The Tail-Average Margin (TAM) at level r is*

$$\text{TAM}_r(x) := \frac{1}{k} \sum_{j=1}^k x_{(j)}, \quad (13)$$

where $x_{(1)} \geq x_{(2)} \geq \dots \geq x_{(m)}$ are the order statistics of x .

We note two boundary cases: $\text{TAM}_{1/m}(x) = \max_j x_j$ (take $k = 1$), and $\text{TAM}_1(x) = \frac{1}{m} \sum_j x_j$ (take $k = m$). For fixed $r \in (0, 1)$, TAM then interpolates strictly between the maximum and the mean by averaging over the top r -fraction of entries.

The TAM criterion is a conservative certificate for top- k list retrieval. Indeed, since

$$\text{TAM}_r(x) = \frac{1}{k} \sum_{j=1}^k x_{(j)} \geq x_{(k)},$$

the inequality $s_{i,i} > \text{TAM}_r((s_{j,i})_{j \neq i})$ implies $s_{i,i} > s_{(k),i}$ for the ordered competitor scores in column i . Hence at most $k-1$ competitors can outrank the signal. In this sense, exact top-1 retrieval implies TAM retrieval, and TAM retrieval implies inclusion of the correct item in the top- k candidate list.

The TAM functional is essentially a reparametrization of the empirical *Conditional Value-at-Risk* (CVaR) studied extensively in the quantitative risk literature; see e.g., [Rockafellar](#)

and Uryasev [2000]. A classical observation is that TAM admits a convex variational representation,

$$\text{TAM}_r(x) = \min_{\mu \in \mathbb{R}} \left\{ \mu + \frac{1}{k} \sum_{j=1}^m (x_j - \mu)_+ \right\}, \quad (14)$$

where $(x)_+ = \max\{x, 0\}$ is the hinge function. The minimizer $\mu^* = x_{(k)}$ of (14) is the empirical $(1-r)$ -quantile of x , also known as the *Value-at-Risk*. The optimal value of (14) is then the mean of the exceedances of x above this quantile, which recovers the tail-average interpretation of (13). As an immediate consequence of (14), we see that $x \mapsto \text{TAM}_r(x)$ is convex on $\mathbb{R}^m$, as it is the pointwise minimum over μ of functions jointly convex in (x, μ) .

For optimization it is convenient to replace the piecewise-linear hinge $(\cdot)_+$ in (14) with a differentiable surrogate. For a smoothing parameter $\beta > 0$, let

$$\phi_\beta(t) := \frac{1}{\beta} \log(1 + e^{\beta t})$$

denote the softplus function, and define the *smoothed* TAM

$$\text{TAM}_{r,\beta}(x) := \min_{\mu \in \mathbb{R}} \left\{ \mu + \frac{1}{k} \sum_{j=1}^m \phi_\beta(x_j - \mu) \right\}. \quad (15)$$

Since ϕ_β is convex, $\text{TAM}_{r,\beta}$ is convex in x ; moreover $\phi_\beta(t) \rightarrow (t)_+$ pointwise as $\beta \rightarrow \infty$, and hence $\text{TAM}_{r,\beta}(x) \rightarrow \text{TAM}_r(x)$ in the same limit. The exact TAM is recovered in the limiting notation $\text{TAM}_{r,\infty} = \text{TAM}_r$.

Throughout the remainder of the paper we specialize TAM to the linear associative memory setting: the collection $x_1, \dots, x_m$ is taken to be the competing scores $(s_{j,i})_{j \neq i}$ associated with a query v_i , so that $m = n - 1$ and TAM_r quantifies the strength of the top r -fraction of competitors. The next subsection turns this into a convex learning objective.

3.2 Learning under the TAM criterion

With the TAM functional in hand, we now formulate retrieval and learning for the linear associative memory as a convex optimization problem.

Given stored pairs $\{(u_i, v_i)\}_{i=1}^n \subset \mathbb{R}^d \times \mathbb{R}^d$ and a memory or weight matrix $W \in \mathbb{R}^{d \times d}$, recall that the score matrix is $s_{j,i} = u_j^\top W v_i$. Fix a tail fraction $r \in (0, 1)$ and set $k = \lceil r(n-1) \rceil$. The *TAM retrieval criterion* asks that each signal score exceed the TAM-aggregate of its competitors:

$$s_{i,i} > \text{TAM}_r((s_{j,i})_{j \neq i}), \quad \forall i \in [n]. \quad (16)$$

For $r = 1/(n-1)$, (16) reduces to the exact top-1 criterion (1) studied in Section 2; for fixed $r \in (0, 1)$, it imposes an upper-tail margin requirement that controls the mass of strong competitors rather than a single extreme-value outlier.

Rather than demanding strict satisfaction of (16), we minimize a convex surrogate that penalizes violations softly. Let

$$\ell(t) := \log(1 + e^{-t})$$

denote the logistic loss, and let $\beta > 0$ be the smoothing parameter of $\text{TAM}_{r,\beta}$. We consider the regularized objective

$$\min_{W \in \mathbb{R}^{d \times d}} \left\{ \sum_{i=1}^n \ell(s_{i,i} - \text{TAM}_{r,\beta}((s_{j,i})_{j \neq i})) + \frac{\lambda n}{2d^2} \|W\|_{\text{F}}^2 \right\}, \quad (17)$$

where $\lambda \geq 0$ is a ridge penalty. Since each score $s_{j,i}$ is linear in W and $\text{TAM}_{r,\beta}$ is convex, the margin $s_{i,i} - \text{TAM}_{r,\beta}((s_{j,i})_{j \neq i})$ is concave in W ; composing with the decreasing convex ℓ yields a convex function of W , so (17) is a convex program (strongly convex whenever $\lambda > 0$).

The variational representation (15) of $\text{TAM}_{r,\beta}$ suggests lifting (17) to a joint optimization over W and the auxiliary quantile variables $\mu = (\mu_1, \dots, \mu_n) \in \mathbb{R}^n$. Substituting (15) into (17) yields

$$\min_{W, \mu} \left\{ \sum_{i=1}^n \ell \left(s_{i,i} - \mu_i - \frac{1}{k} \sum_{j \neq i} \phi_{\beta}(s_{j,i} - \mu_i) \right) + \frac{\lambda n}{2d^2} \|W\|_{\text{F}}^2 \right\}, \quad (18)$$

Minimizing (18) over μ at fixed W recovers (17). This is the workhorse formulation for the leave-one-out analysis of Section 4.

Remark 1 (Gaussian heuristic for the quadratic scale). *The asymptotic distinction between the exact maximum and TAM_r can be made precise under a Gaussian toy model. Suppose the competitor scores $(s_{j,i})_{j \neq i}$ were i.i.d. $\mathcal{N}(0, \sigma^2)$ with $\sigma = \Theta(\sqrt{n}/d)$. Then the maximum would concentrate at $\sigma\sqrt{2 \log n}$, while TAM_r at fixed $r \in (0, 1)$ would concentrate at $\sigma \cdot \kappa_r$ with*

$$\kappa_r = \mathbb{E}[Z \mid Z \geq \Phi^{-1}(1-r)] = \frac{\varphi(\Phi^{-1}(1-r))}{r}, \quad Z \sim \mathcal{N}(0, 1),$$

depending only on r . Comparing either scale with the $O(1)$ signal strength locates the heuristic retrieval threshold at $d^2 \asymp n \log n$ in the first case—consistent with Theorem 1—and $d^2 \asymp n$ in the second.

In this model, the Gaussianity and scale of the competitor scores are not assumed a priori; they are asymptotically exact under the correlation matrix memory solution of Section 2.1. The proof of Lemma 9 shows that, in the regime $n/d^2 \rightarrow \alpha$, the competitor scores are (approximately) i.i.d. $\mathcal{N}(0, \alpha)$. For the TAM-optimal weight matrix $\widehat{W}$ defined below, the same Gaussian structure emerges asymptotically as an output of the leave-one-out analysis of Section 4; in that case the Gaussian variance is no longer explicit a priori, but is determined self-consistently by the scalar fixed point equations. In both cases the heuristic above is borne out by the theory.

It is useful to compare (17) with the canonical convex surrogate for the top-1 retrieval criterion (1), namely the regularized multiclass softmax cross-entropy:

$$\min_{W \in \mathbb{R}^{d \times d}} \sum_{i=1}^n \left[-s_{i,i} + \log \sum_{j=1}^n e^{s_{j,i}} \right] + \frac{\lambda n}{2d^2} \|W\|_{\text{F}}^2. \quad (19)$$

Denote the log-sum-exp aggregate of the competitor scores by $\text{LSE}((s_{j,i})_{j \neq i}) := \log \sum_{j \neq i} e^{s_{j,i}}$. A short calculation rewrites each summand in (19) as

$$-s_{i,i} + \log \sum_{j=1}^n e^{s_{j,i}} = \ell \left(s_{i,i} - \text{LSE}((s_{j,i})_{j \neq i}) \right),$$

with the same logistic loss ℓ used in (17). In this form, (19) has precisely the same structure as the TAM objective (17): a logistic loss applied to the margin between the signal score $s_{i,i}$ and a convex aggregate of the competitor scores $(s_{j,i})_{j \neq i}$. They differ only in the choice of aggregate. The log-sum-exp is a smoothed maximum, with $\text{LSE}((s_{j,i})_{j \neq i}) = \max_{j \neq i} s_{j,i} + O(\log n)$, and therefore inherits the $\sqrt{2 \log n}$ extreme-value statistics of the top-1 criterion. Replacing it with the tail average $\text{TAM}_{r,\beta}$, whose fluctuations are $O(\sigma)$ (Remark 1), enables retrieval at the quadratic model-size scale $d^2 \asymp n$ instead of $d^2 \asymp n \log n$.

3.3 Performance metrics

With the TAM learning problem in place, we consider three asymptotic quantities that together characterize retrieval performance: the empirical average fitting loss, the per-sample TAM margin, and the distribution of signal percentiles. Throughout this subsection, $(\widehat{W}, \widehat{\mu})$ denotes the unique minimizer of (18), with corresponding scores $\widehat{s}_{j,i} = u_j^\top \widehat{W} v_i$.

Empirical average fitting loss. Let

$$L_n(W, \mu) := \frac{1}{n} \sum_{i=1}^n \ell \left(s_{i,i} - \mu_i - \frac{1}{k} \sum_{j \neq i} \phi_\beta(s_{j,i} - \mu_i) \right)$$

denote the normalized data-fitting part of (18), and write $\widehat{L}_n := L_n(\widehat{W}, \widehat{\mu})$ for its value at the optimizer. We will say that the TAM problem is *asymptotically fit at load α* if $\widehat{L}_n \rightarrow 0$ in probability as $n, d \rightarrow \infty$ with $n/d^2 \rightarrow \alpha$ and $\lambda \rightarrow 0$. Since ℓ is nonnegative and vanishes only in the large-margin limit, asymptotic fit is equivalent to the TAM retrieval criterion (16) holding with a positive margin for all but a vanishing fraction of keys.

TAM margin distribution. A finer summary than $\widehat{L}_n$ is the empirical distribution of the per-sample TAM margins

$$\widehat{m}_i := \widehat{s}_{i,i} - \widehat{\mu}_i - \frac{1}{k} \sum_{j \neq i} \phi_\beta(\widehat{s}_{j,i} - \widehat{\mu}_i), \quad i \in [n],$$

which are precisely the arguments of the logistic loss appearing in $\widehat{L}_n$. A positively-supported limiting distribution for $\widehat{m}_i$ indicates asymptotic fit; its shape shows how comfortably the signal beats the tail average.

Signal percentile distribution. As a complementary, rank-based summary, we also ask, for each sample, what fraction of competitors have score below the signal. Formally, define the *signal percentile*

$$\Omega_i := \frac{1}{n-1} \sum_{j \neq i} \mathbf{1}\{\widehat{s}_{j,i} \leq \widehat{s}_{i,i}\}, \quad i \in [n],$$

so that Ω_i close to 1 signals that the i th signal dominates nearly all of its competitors and $\Omega_i = 0$ that every competitor beats it. The random probability measure

$$\nu_n := \frac{1}{n} \sum_{i=1}^n \delta_{\Omega_i}$$

on $[0, 1]$ is our object of study: when does ν_n converge in probability (say, in Wasserstein-1) to a deterministic limit?

The next section develops a leave-one-out theory for these three observables under the TAM optimizer in the quadratic regime $n/d^2 \rightarrow \alpha$ with fixed $r \in (0, 1)$.

Benchmark: correlation matrix memory. We close this section with a benchmark calculation. For the non-optimized correlation matrix memory $W^\times$ of Section 2.1, the percentile law can be computed explicitly. In the regime $n, d \rightarrow \infty$ with $n/d^2 \rightarrow \alpha \in (0, \infty)$, the single-coordinate law, Lemma 3 in Appendix A.1, says that the signal score is asymptotically Gaussian with mean 1 and variance α , while a typical competitor score is asymptotically Gaussian with mean 0 and the same variance. Thus the signal is shifted above the competitor cloud by signal-to-noise ratio $\alpha^{-1/2}$. Writing Φ for the standard normal CDF, the limiting percentile has the form $\Phi(G + \alpha^{-1/2})$, where $G \sim \mathcal{N}(0, 1)$. Equivalently, its CDF is

$$F_\alpha(\omega) := \Phi\left(\Phi^{-1}(\omega) - \alpha^{-1/2}\right), \quad \omega \in (0, 1), \quad (20)$$

with density $f_\alpha(\omega) = \exp(\Phi^{-1}(\omega)/\sqrt{\alpha} - 1/(2\alpha))$. The parameter $\alpha^{-1/2}$ plays the role of a signal-to-noise ratio: the law concentrates at 1 as $\alpha \rightarrow 0$ and tends to $\text{Unif}(0, 1)$ as $\alpha \rightarrow \infty$. Appendix A.4 upgrades this one-coordinate heuristic to a law of large numbers for the empirical percentile distribution.

4 Asymptotic theory of TAM

We now study the TAM optimizer in the quadratic regime $n, d \rightarrow \infty$ with $n/d^2 \rightarrow \alpha$ and $k/n \rightarrow r \in (0, 1)$. Throughout this section the smoothing parameter $\beta < \infty$ and ridge parameter $\lambda > 0$ are fixed. The goal is to predict the empirical laws of the true scores $s_{i,i}(\widehat{W})$ and competitor scores $s_{j,i}(\widehat{W})$, and hence the limiting TAM loss, margin distribution, and signal-percentile profile introduced in Section 3.3.

The main technique is a leave-one-out (LOO) analysis. We compare the TAM optimizer with the optimizer obtained after removing a held-out pair (u_h, v_h) , and use this comparison to separate the fresh randomness of the held-out sample from the remaining problem. The results in this section are conditional on six postulates stated below. The first three are leave-one-out stability assumptions, used to justify the rank-one LOO reduction. The last three are spectral and self-averaging inputs, used to close the scalar equations. Under these postulates, the high-dimensional TAM optimizer is described by a two-parameter scalar variational problem, stated in Result 2.

The section is organized as follows. We first compute the gradient and Hessian of the smoothed TAM loss. We then formulate the LOO rank-one reduction, state the scalar

variational prediction, and finally derive the finite-dimensional equations that lead to this prediction. The ridgeless limit $\lambda \downarrow 0$ and the associated phase transition are deferred to Section 5.

4.1 Gradient and Hessian of the TAM loss

Throughout this section the threshold variables are unregularized, and we work with the joint TAM objective of (18), written with the reparameterized ridge:

$$\Psi(W, \mu) = \sum_{i=1}^n \ell \left(s_{i,i}(W) - \mu_i - \frac{1}{k} \sum_{j \neq i} \phi_\beta(s_{j,i}(W) - \mu_i) \right) + \frac{\lambda n}{2d^2} \|W\|_F^2,$$

where $s(W) = U^\top W V \in \mathbb{R}^{n \times n}$, with entries $s_{j,i}(W) = u_j^\top W v_i$, and $U = [u_1 \cdots u_n]$, $V = [v_1 \cdots v_n]$ are the data matrices from the Gaussian model (3). We use lowercase $s_{j,i}$ for finite-dimensional scores, reserving uppercase S for scalar limiting random variables. When the underlying matrix is clear from context, we write $s_{j,i}$ instead of $s_{j,i}(W)$; in particular, at the full optimizer this means $s_{j,i} = s_{j,i}(\widehat{W})$. Since Ψ depends on W only through $s(W)$, the chain rule reduces the computation of $\nabla_W \Psi$ to an $n \times n$ matrix of score derivatives,

$$\nabla_W \Psi = U G V^\top + \frac{\lambda n}{d^2} W, \quad G_{j,i} := \frac{\partial \Psi}{\partial s_{j,i}}. \quad (21)$$

The remainder of this subsection evaluates the entries of G and rewrites $\nabla_W \Psi$ in a form convenient for the leave-one-out analysis in Section 4.2.

The i -th summand of Ψ equals $\ell(m_i)$, where the *TAM aggregate* and *margin* associated with sample i are

$$c_i := \mu_i + \frac{1}{k} \sum_{j \neq i} \phi_\beta(s_{j,i} - \mu_i), \quad m_i := s_{i,i} - c_i.$$

The margin m_i is the only part of the i -th summand that depends on column i of the score matrix, and the i -th summand is the only summand that does. Differentiating $\ell(m_i)$ in $s_{j,i}$ therefore yields

$$G_{j,i} = \ell'(m_i) \frac{\partial m_i}{\partial s_{j,i}} = \begin{cases} \ell'(m_i), & j = i, \\ -\frac{1}{k} \ell'(m_i) \phi'_\beta(s_{j,i} - \mu_i), & j \neq i. \end{cases} \quad (22)$$

Both ℓ' and ϕ'_β are conveniently expressed through the sigmoid $\sigma(t) := (1 + e^{-t})^{-1}$:

$$\ell'(t) = \sigma(t) - 1 = -\sigma(-t), \quad \phi'_\beta(t) = \sigma(\beta t).$$

Substituting into (22) gives the two scalar weights

$$a_i := \sigma(-m_i) = -\ell'(m_i) \in (0, 1),$$

and, for $j \neq i$,

$$q_{j,i} := \sigma(\beta(s_{j,i} - \mu_i)) = \phi'_\beta(s_{j,i} - \mu_i) \in (0, 1), \quad j \neq i.$$

In these variables (22) becomes $G_{j,i} = -a_i$ for $j = i$ and $G_{j,i} = \frac{1}{k}q_{j,i}a_i$ otherwise. Equivalently $G = Q\Lambda_a$, where $\Lambda_a := \text{diag}(a_1, \dots, a_n)$ and

$$Q_{j,i} := \begin{cases} -1, & j = i, \\ \frac{1}{k}q_{j,i}, & j \neq i. \end{cases}$$

Substituting $G = Q\Lambda_a$ into (21) and performing the analogous calculation for μ — where $\partial m_i / \partial \mu_i = -1 + \frac{1}{k} \sum_{j \neq i} q_{j,i} = -(Q^\top \mathbf{1}_n)_i$ — yields

$$\nabla_W \Psi(W, \mu) = U Q \Lambda_a V^\top + \frac{\lambda n}{d^2} W, \quad (23)$$

$$\nabla_\mu \Psi(W, \mu) = -\Lambda_a Q^\top \mathbf{1}_n,$$

where $\mathbf{1}_n \in \mathbb{R}^n$ is the all-ones vector. Expanding (23) by rows of Q separates the diagonal and off-diagonal contributions,

$$\nabla_W \Psi = - \sum_{i=1}^n a_i u_i v_i^\top + \frac{1}{k} \sum_{i=1}^n a_i \sum_{j \neq i} q_{j,i} u_j v_i^\top + \frac{\lambda n}{d^2} W. \quad (24)$$

For the Taylor expansion in Section 4.2 we will also need the second derivative of the outer loss at the margin, which is governed by a companion diagonal:

$$b_i := \ell''(m_i) = a_i(1 - a_i), \quad \Lambda_b := \text{diag}(b_1, \dots, b_n). \quad (25)$$

Definition 2 (Full Hessian). *Let $H : \mathbb{R}^{d \times d} \rightarrow \mathbb{R}^{d \times d}$ denote the Hessian operator of Ψ in the W variable, evaluated at a fixed point (W, μ) . Applied to a perturbation Δ , it is*

$$H[\Delta] := \frac{\lambda n}{d^2} \Delta + U M (U^\top \Delta V) V^\top.$$

Here $M : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$ acts columnwise,

$$M(X)_{:,i} = M_i X_{:,i},$$

where

$$M_i := b_i Q_{:,i} Q_{:,i}^\top + a_i \text{diag}(\tilde{R}_{:,i}),$$

$$\tilde{R}_{j,i} := \begin{cases} \frac{\beta}{k} q_{j,i} (1 - q_{j,i}), & j \neq i, \\ 0, & j = i. \end{cases}$$

The operator H is self-adjoint and strictly positive-definite, with smallest eigenvalue at least $\lambda n / d^2$. Under vectorization, writing $\mathcal{D} := \text{diag}(M_1, \dots, M_n) \in \mathbb{R}^{n^2 \times n^2}$, the same operator has matrix representation

$$H = \frac{\lambda n}{d^2} I_{d^2} + (V \otimes U) \mathcal{D} (V \otimes U)^\top.$$

4.2 Leave-one-out reduction

We relate the joint minimizer $(\widehat{W}, \widehat{\mu})$ of (18) to a leave-one-out reference point by a first-order expansion. Fix a held-out index $h \in [n]$, and let $(\widehat{W}^{\setminus h}, \widehat{\mu}^{\setminus h})$ denote the minimizer of the TAM objective with sample h removed from both the outer sum and from all inner summations over interferers:

$$\Psi^{\setminus h}(W, \mu) := \sum_{i \neq h} \ell(s_{i,i}(W) - \mu_i - \frac{1}{k} \sum_{\substack{j \neq i \\ j \neq h}} \phi_\beta(s_{j,i}(W) - \mu_i)) + \frac{\lambda n}{2d^2} \|W\|_F^2.$$

Write $\Delta W := \widehat{W} - \widehat{W}^{\setminus h}$, $\Delta \mu := \widehat{\mu} - \widehat{\mu}^{\setminus h}$, and $\Delta S := U^\top (\Delta W) V \in \mathbb{R}^{n \times n}$ for the corresponding perturbations. A hat decoration (e.g., $\hat{a}_i, \hat{q}_{j,i}, \hat{b}_i, \hat{m}_i$) denotes a quantity from Section 4.1 evaluated at $(\widehat{W}, \widehat{\mu})$; a superscript $\setminus h$ denotes the same quantity evaluated at $(\widehat{W}^{\setminus h}, \widehat{\mu}^{\setminus h})$. We adopt the conventions $a_h^{\setminus h} := 0$ and $q_{j,h}^{\setminus h} := q_{h,i}^{\setminus h} := 0$, so that $Q^{\setminus h}, \Lambda_{a^{\setminus h}}$ extend naturally to $n \times n$ arrays.

4.2.1 LOO Hessian and rank-one reduction

The LOO Hessian has the same form as the full Hessian in Definition 2, with the h -th sample removed and all coefficients evaluated at $(\widehat{W}^{\setminus h}, \widehat{\mu}^{\setminus h})$. Explicitly, $H^{\setminus h} : \mathbb{R}^{d \times d} \rightarrow \mathbb{R}^{d \times d}$ is defined by

$$H^{\setminus h}[\Delta] := \frac{\lambda n}{d^2} \Delta + U M^{\setminus h} (U^\top \Delta V) V^\top, \quad (26)$$

where $M^{\setminus h} : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n \times n}$ acts columnwise as

$$M^{\setminus h}(X)_{:,i} = M_i^{\setminus h} X_{:,i} \quad (i \neq h), \quad M^{\setminus h}(X)_{:,h} = 0. \quad (27)$$

In the above display, $M_i^{\setminus h} \in \mathbb{R}^{n \times n}$ is a symmetric matrix with a rank-one-plus-diagonal form

$$\begin{aligned} M_i^{\setminus h} &:= b_i^{\setminus h} Q_{:,i}^{\setminus h} (Q_{:,i}^{\setminus h})^\top + a_i^{\setminus h} \text{diag}(\widetilde{R}_{:,i}^{\setminus h}) \\ \widetilde{R}_{j,i}^{\setminus h} &:= \begin{cases} \frac{\beta}{k} q_{j,i}^{\setminus h} (1 - q_{j,i}^{\setminus h}) & j \neq i, j \neq h \\ 0, & \text{otherwise} \end{cases}. \end{aligned} \quad (28)$$

Here $b_i^{\setminus h}$ is the quantity in (25) evaluated at $(\widehat{W}^{\setminus h}, \widehat{\mu}^{\setminus h})$. The operator $H^{\setminus h}$ is self-adjoint and strictly positive-definite on $\mathbb{R}^{d \times d}$: its smallest eigenvalue is at least $\lambda n / d^2 = \Theta(1)$ in the regime $d^2 \asymp n$, so that $\|(H^{\setminus h})^{-1}\|_{\text{op}} = \mathcal{O}_{\prec}(1)$.

Main result. The leave-one-out perturbation is expected to be governed, to first order, by the held-out feature pair (u_h, v_h) .

Result 1 (LOO rank-one reduction). *Under the postulates (A1)–(A3) stated below, with high probability,*

$$\widehat{W} - \widehat{W}^{\setminus h} = \sigma(-\hat{m}_h) (H^{\setminus h})^{-1} [u_h v_h^\top] + R_h, \quad \|R_h\|_F = \mathcal{O}_{\prec}(d^{-1/2}). \quad (29)$$

Here $\hat{m}_h$ is the TAM margin at the held-out sample evaluated at the full minimizer $(\widehat{W}, \widehat{\mu})$:

$$\hat{m}_h = s_{h,h}(\widehat{W}) - \hat{\mu}_h - \frac{1}{k} \sum_{j \neq h} \phi_\beta(s_{j,h}(\widehat{W}) - \hat{\mu}_h).$$

Thus the leading change in the weight matrix is the inverse LOO Hessian applied to $\sigma(-\hat{m}_h) u_h v_h^\top$, with a Frobenius-norm correction of lower order. In Section 4.3, this formula is combined with trace concentration for $(H^{\setminus h})^{-1}$ to obtain the susceptibility equation.

4.2.2 Justification of Result 1

We derive (29) from the following LOO postulates.

LOO postulates. We assume:

$$(A1) \max_{i \neq h} |(\Delta \mu)_i| = \mathcal{O}_{\prec}(n^{-1});$$

$$(A2) \max_{j, i \neq h} |(\Delta S)_{j,i}| = \mathcal{O}_{\prec}(d^{-1});$$

$$(A3) \max_{j, i \in [n]} |s_{j,i}(\widehat{W})| = \mathcal{O}_{\prec}(1).$$

in the sense of stochastic domination, absorbing d^ε factors. We use the same postulates after removing any fixed finite set of samples, uniformly over such sets; in particular, (A3) is imposed on the optimizer of the problem under consideration. These assumptions control the Taylor remainder and the coefficient perturbations in the LOO expansion. The spectral and self-averaging postulates needed to close the scalar equations are stated separately in Section 4.3.

Step 1: Subtract the stationarity conditions. The W -components of the stationarity conditions (23) at the full and LOO optimizers, subtracted, yield

$$\frac{\lambda n}{d^2} \Delta W + U (\hat{Q} \Lambda_{\hat{a}} - Q^{\setminus h} \Lambda_{a^{\setminus h}}) V^\top = 0. \quad (30)$$

Step 2: Separate the terms involving the held-out sample. With the extension conventions on $a^{\setminus h}$ and $q^{\setminus h}$, the matrix $\hat{Q} \Lambda_{\hat{a}} - Q^{\setminus h} \Lambda_{a^{\setminus h}}$ decomposes entrywise as

$$(\hat{Q} \Lambda_{\hat{a}} - Q^{\setminus h} \Lambda_{a^{\setminus h}})_{j,i} = \begin{cases} -\hat{a}_h & (j, i) = (h, h), \\ \hat{a}_h \hat{q}_{j,h}/k & j \neq h, i = h, \\ \hat{a}_i \hat{q}_{h,i}/k & j = h, i \neq h, \\ \mathcal{R}_{j,i}^{\setminus h} & j, i \neq h, \end{cases}$$

where the first three cases are the entries involving the held-out sample: the held-out outer term ($i = h$) and the held-out interferer row ($j = h$, across interior columns) are present in

the full objective but absent in the LOO objective. The remaining entries are collected in $\mathcal{R}^{\setminus h} \in \mathbb{R}^{n \times n}$, supported on $\{j, i \neq h\}$:

$$\mathcal{R}_{j,i}^{\setminus h} := \begin{cases} \frac{\hat{a}_i \hat{q}_{j,i} - a_i^{\setminus h} q_{j,i}^{\setminus h}}{k} & j \neq i, j, i \neq h, \\ a_i^{\setminus h} - \hat{a}_i & j = i \neq h. \end{cases} \quad (31)$$

Separating these entries inside (30),

$$\frac{\lambda n}{d^2} \Delta W + U \mathcal{R}^{\setminus h} V^\top = \hat{a}_h u_h v_h^\top - \hat{a}_h \eta_h v_h^\top - u_h \xi_h^\top, \quad (32)$$

where

$$\eta_h := \frac{1}{k} \sum_{j \neq h} \hat{q}_{j,h} u_j, \quad \xi_h := \frac{1}{k} \sum_{i \neq h} \hat{a}_i \hat{q}_{h,i} v_i.$$

Step 3: Linearize the common part. In Appendix C, we carry out a first-order Taylor expansion of $\mathcal{R}^{\setminus h}$ in the perturbations ΔS and $\Delta \mu$, with all higher-order terms collected into a residual matrix $\hat{E}^{\setminus h}$ supported on $\{j, i \neq h\}$. This yields

$$\mathcal{R}_{:,i}^{\setminus h} = M_i^{\setminus h} \Delta S_{:,i} + \hat{E}_{:,i}^{\setminus h}, \quad i \neq h, \quad (33)$$

with $M_i^{\setminus h}$ as in (28). Combining (33) with (32) and the definition of $H^{\setminus h}$ yields

$$H^{\setminus h}[\Delta W] = \hat{a}_h u_h v_h^\top - \hat{a}_h \eta_h v_h^\top - u_h \xi_h^\top - U \hat{E}^{\setminus h} V^\top. \quad (34)$$

Moreover, we show in Appendix C that, under (A1)–(A3),

$$|\hat{E}_{i,i}^{\setminus h}| = \mathcal{O}_{\prec}(n^{-1}) \text{ for } i \neq h, \quad |\hat{E}_{j,i}^{\setminus h}| = \mathcal{O}_{\prec}(n^{-2}) \text{ for } j \neq i, j, i \neq h. \quad (35)$$

Step 4: Invert and bound the correction. Applying $(H^{\setminus h})^{-1}$ to (34) produces (29) with

$$R_h = (H^{\setminus h})^{-1} \left[-\hat{a}_h \eta_h v_h^\top - u_h \xi_h^\top - U \hat{E}^{\setminus h} V^\top \right]. \quad (36)$$

Since $\|(H^{\setminus h})^{-1}\|_{\text{op}} = \mathcal{O}_{\prec}(1)$, it suffices to bound each of the three remaining terms in (36) in Frobenius norm at scale $\mathcal{O}_{\prec}(d^{-1/2})$.

Writing $k \eta_h = U \hat{q}_{:,h}$ (with the h -th entry of $\hat{q}_{:,h}$ understood to be zero), the Marchenko–Pastur bound $\|U\|_{\text{op}} = \mathcal{O}_{\prec}(\sqrt{d})$ in the regime $d^2 \asymp n$, together with the deterministic entry-wise bound $\|\hat{q}_{:,h}\|_2 \leq \sqrt{n-1}$, gives $\|\eta_h\|_2 \leq \|U\|_{\text{op}} \|\hat{q}_{:,h}\|_2 / k = \mathcal{O}_{\prec}(\sqrt{nd}/n) = \mathcal{O}_{\prec}(d^{-1/2})$; the same argument applies to ξ_h . Consequently

$$\left\| \hat{a}_h \eta_h v_h^\top \right\|_{\text{F}} = |\hat{a}_h| \|\eta_h\|_2 \|v_h\|_2 = \mathcal{O}_{\prec}(d^{-1/2}), \quad \left\| u_h \xi_h^\top \right\|_{\text{F}} = \mathcal{O}_{\prec}(d^{-1/2}). \quad (37)$$

Next, we consider the term $U \hat{E}^{\setminus h} V^\top$ in (36). Let $\hat{E}^{\text{d}}$ and $\hat{E}^{\text{o}}$ denote the diagonal and off-diagonal parts of $\hat{E}^{\setminus h}$, respectively. The diagonal part is a diagonal matrix, so $\left\| \hat{E}^{\text{d}} \right\|_{\text{op}} =$

$\max_{i \neq h} |\hat{E}_{i,i}^{\setminus h}| = \mathcal{O}_{\prec}(n^{-1})$ by (35). The off-diagonal part is controlled through its Frobenius norm,

$$\left\| \hat{E}^{\circ} \right\|_{\text{op}} \leq \left\| \hat{E}^{\circ} \right\|_{\text{F}} = \left(\sum_{\substack{j \neq i \\ j, i \neq h}} |\hat{E}_{j,i}^{\setminus h}|^2 \right)^{1/2} \leq \sqrt{n^2 \cdot \mathcal{O}_{\prec}(n^{-4})} = \mathcal{O}_{\prec}(n^{-1}),$$

again by (35). Combining, $\left\| \hat{E}^{\setminus h} \right\|_{\text{op}} = \mathcal{O}_{\prec}(n^{-1}) = \mathcal{O}_{\prec}(d^{-2})$. Submultiplicativity with $\|U\|_{\text{op}}, \|V\|_{\text{op}} = \mathcal{O}_{\prec}(\sqrt{d})$ yields $\left\| U \hat{E}^{\setminus h} V^{\top} \right\|_{\text{op}} \leq \mathcal{O}_{\prec}(d \cdot d^{-2}) = \mathcal{O}_{\prec}(d^{-1})$. Since $U \hat{E}^{\setminus h} V^{\top} \in \mathbb{R}^{d \times d}$ has rank at most d , $\|\cdot\|_{\text{F}} \leq \sqrt{d} \|\cdot\|_{\text{op}}$, and

$$\left\| U \hat{E}^{\setminus h} V^{\top} \right\|_{\text{F}} \leq \sqrt{d} \left\| U \hat{E}^{\setminus h} V^{\top} \right\|_{\text{op}} = \mathcal{O}_{\prec}(d^{-1/2}). \quad (38)$$

Combining (37) and (38) and applying $(H^{\setminus h})^{-1}$ establishes (29). $\square$

4.3 Scalar variational prediction

We now state the asymptotic characterization of the TAM minimizer. In addition to the LOO postulates (A1)–(A3), we assume the following spectral and self-averaging postulates (A4)–(A6). The derivation is given in Section 4.4.

Spectral and self-averaging postulates. We assume:

$$(A4) \quad \left\| \widehat{W} \right\|_{\text{op}} = \mathcal{O}_{\prec}(\sqrt{d});$$

(A5) (Empirical-mean concentration.) For the bounded or pseudo-Lipschitz score observables used in the sequel, including the observables generated by the scalar reductions below,

$$\frac{1}{n} \sum_{i=1}^n \psi(s_{i,i}(\widehat{W})) - \mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n \psi(s_{i,i}(\widehat{W})) \right] = \mathcal{O}_{\prec}(d^{-1/2}),$$

and

$$\frac{1}{n(n-1)} \sum_{i \neq j} \psi(s_{j,i}(\widehat{W})) - \mathbb{E} \left[\frac{1}{n(n-1)} \sum_{i \neq j} \psi(s_{j,i}(\widehat{W})) \right] = \mathcal{O}_{\prec}(d^{-1/2}).$$

(A6) (Trace concentration.)

$$\frac{1}{d^2} \text{tr}(H^{-1}) - \mathbb{E} \left[\frac{1}{d^2} \text{tr}(H^{-1}) \right] = \mathcal{O}_{\prec}(d^{-1/2}).$$

Here (A4) is the spectral input, while (A5) and (A6) are self-averaging inputs at the common scale $\mathcal{O}_{\prec}(d^{-1/2})$ for empirical means and normalized inverse traces, respectively. The

corresponding trace estimate for a fixed finite LOO Hessian follows from (A6) by the same finite-removal stability used in Section 4.2.

For $\nu > 0$, define the scalar TAM functional

$$c_r(\nu) := \min_{\mu \in \mathbb{R}} \left\{ \mu + \frac{1}{r} \mathbb{E}_{Z \sim \mathcal{N}(0,1)} [\phi_\beta(\nu Z - \mu)] \right\}. \quad (39)$$

Let $\mu_r(\nu)$ denote the minimizer in (39). It is characterized by

$$r = \mathbb{E}_{Z \sim \mathcal{N}(0,1)} \left[\sigma(\beta(\nu Z - \mu_r(\nu))) \right]. \quad (40)$$

Definition 3 (Scalar saddle and reduced potentials). *For $\nu, \chi > 0$, define the scalar Moreau envelope*

$$M_r(\nu, \chi) := \mathbb{E}_{G \sim \mathcal{N}(0,1)} \left[\min_{s \in \mathbb{R}} \left[\frac{(s - \nu G)^2}{2\chi} + \ell(s - c_r(\nu)) \right] \right]. \quad (41)$$

The scalar saddle potential is

$$\mathcal{P}_{\alpha,r}(\nu, \chi) := \frac{\lambda}{2} \nu^2 - \frac{\nu^2}{2\alpha\chi} + M_r(\nu, \chi),$$

and the reduced scalar potential is

$$\mathcal{V}_{\alpha,r}(\nu) := \sup_{\chi > 0} \mathcal{P}_{\alpha,r}(\nu, \chi). \quad (42)$$

Proposition 6, stated in Section 4.4, shows that, for each fixed $\nu > 0$, the maximization over χ is well posed and has a unique maximizer, and that the reduced potential $\mathcal{V}_{\alpha,r}$ is strongly convex in ν .

For $\chi > 0$, define

$$\text{Prox}_\chi(x) := \arg \min_{y \in \mathbb{R}} \left\{ \frac{(y - x)^2}{2\chi} + \ell(y) \right\}. \quad (43)$$

Equivalently, $y = \text{Prox}_\chi(x)$ if and only if

$$y - \chi \sigma(-y) = x.$$

If $G \sim \mathcal{N}(0, 1)$, define

$$S_{\nu,\chi} := c_r(\nu) + \text{Prox}_\chi(\nu G - c_r(\nu)). \quad (44)$$

Let ν_* be the minimizer of $\mathcal{V}_{\alpha,r}$, and let χ_* be the maximizer of $\mathcal{P}_{\alpha,r}(\nu_*, \cdot)$. Define μ_* as the minimizer in (39) at $\nu = \nu_*$, and set

$$c_* := c_r(\nu_*).$$

The variational facts used here, including uniqueness and the equivalence with the finite-dimensional equations derived below, are recorded in Proposition 6.

Result 2 (Scalar description of the joint TAM minimizer). *Fix $(\alpha, r, \beta, \lambda)$ with $\beta, \lambda > 0$, and assume (A1)–(A6). Assume also that the minimizer ν_* of $\mathcal{V}_{\alpha, r}$ is positive. In the joint limit $n, d \rightarrow \infty$ with $n/d^2 \rightarrow \alpha$ and $k/n \rightarrow r$, the following hold in probability:*

(i) (Order-parameter concentration.)

$$\frac{1}{d} \|\widehat{W}\|_{\mathbb{F}} \longrightarrow \nu_*, \quad \frac{1}{d^2} \operatorname{tr}(H^{-1}) \longrightarrow \chi_*. \quad (45)$$

(ii) (Gaussian law for competitor scores.) *For each $i \in [n]$, the empirical distribution $\frac{1}{n-1} \sum_{j \neq i} \delta_{s_{j,i}(\widehat{W})}$ converges weakly to $\mathcal{N}(0, \nu_*^2)$.*

(iii) (Proximal law for diagonal scores.) *The empirical distribution $\frac{1}{n} \sum_{i=1}^n \delta_{s_{i,i}(\widehat{W})}$ converges weakly to the law of S_{ν_*, χ_*} in (44).*

Remark 2 (Reading off the retrieval metrics). *Result 2 directly answers the three performance questions posed in Section 3.3. Since the TAM aggregate converges to c_* , the limiting margin is*

$$M_* := S_{\nu_*, \chi_*} - c_*,$$

and hence the average fitting loss converges to

$$\mathbb{E}[\ell(M_*)]. \quad (46)$$

The empirical margin distribution converges to the law of M_ . Finally, writing Φ for the standard normal CDF, the signal percentile distribution converges to the law of*

$$\Phi\left(\frac{S_{\nu_*, \chi_*}}{\nu_*}\right),$$

because the competitor scores have limiting law $\mathcal{N}(0, \nu_^2)$.*

4.4 Justification of Result 2

Define

$$\nu_d^2 := \mathbb{E}\left[\frac{\|\widehat{W}\|_{\mathbb{F}}^2}{d^2}\right]. \quad (47)$$

With H denoting the full Hessian operator from Definition 2, evaluated at $(\widehat{W}, \widehat{\mu})$, define

$$\chi_d := \mathbb{E}\left[\frac{1}{d^2} \operatorname{tr}(H^{-1})\right]. \quad (48)$$

In this section, we derive the finite-dimensional equations for ν_d and χ_d that lead to Result 2. The derivation proceeds in four steps. First, using (A4) and the bilinear coupling estimate in Appendix E, we identify the off-diagonal competitor scores with Gaussians of

variance ν_d^2 . Second, this Gaussian approximation and the trace input (A6) give the proximal characterization of the diagonal score $s_{i,i}(\widehat{W})$. Third, W -stationarity, together with empirical concentration (A5), yields a finite-dimensional equation for ν_d . Fourth, a random matrix calculation for a simplified Hessian gives the companion equation for χ_d . Combining the two equations gives an approximate residual system, whose zero-residual limit is the saddle-point condition for the scalar potential in Definition 3.

4.4.1 Gaussian law for competitor scores

We begin with the off-diagonal scores. The statement below gives a finite-dimensional joint Gaussian coupling for any fixed number of competitor scores in a single column, using an iterated leave-one-out reduction and the bilinear coupling estimate of Corollary 1.

Proposition 2 (Joint Gaussian limit for competitor scores). *Fix $i \in [n]$, $m \geq 1$, and distinct $j_1, \dots, j_m \in [n] \setminus \{i\}$. Assume (A1)–(A6). Then, in the joint regime $n, d \rightarrow \infty$ with $n/d^2 \rightarrow \alpha$ and $k/n \rightarrow r$: on a common probability space, there exist i.i.d. Gaussians $Y_1, \dots, Y_m \sim \mathcal{N}(0, \nu_d^2)$ such that*

$$\|(s_{j_1,i}(\widehat{W}), \dots, s_{j_m,i}(\widehat{W})) - (Y_1, \dots, Y_m)\|_2 = \mathcal{O}_{\prec}(d^{-1/2}). \quad (49)$$

Remark 3 (Connection with the limiting statement). *Proposition 2 is the finite-dimensional form of the off-diagonal Gaussian law in Result 2. Empirical-measure consequences, such as weak convergence of $\frac{1}{n-1} \sum_{j \neq i} \delta_{s_{j,i}(\widehat{W})}$ to $\mathcal{N}(0, \nu_d^2)$, follow by combining exchangeability with concentration of the optimizer. The expectation-level replacements used below are obtained from this coupling for the finite list of bounded or pseudo-Lipschitz observables appearing in the stationarity identities, together with the moment bounds implicit in (A5).*

Proof outline. Throughout, write $X_k := s_{j_k,i}(\widehat{W})$ for $k = 1, \dots, m$. Set $\widehat{W}_{\text{fresh}} := \widehat{W} \setminus \{i, j_1, \dots, j_m\}$ and $V := \|\widehat{W}_{\text{fresh}} v_i\|^2/d$.

Step 1: Iterated leave-one-out. Apply (29) successively after removing the samples $i, j_1, \dots, j_m$. The finite-removal version of the LOO postulates, together with the spectral and self-averaging postulates (A4)–(A5), gives the same rank-one expansion at each level. The resulting iterated identity expresses

$$\widehat{W} = \widehat{W}_{\text{fresh}} + \sum_{\ell=0}^m \left[\hat{a}_{i_\ell}^{I_\ell} (H^{I_\ell})^{-1} [u_{i_\ell} v_{i_\ell}^\top] + R_\ell \right], \quad (50)$$

where $i_0 = i$, $i_\ell = j_\ell$ for $\ell \geq 1$, $I_\ell := \{i, j_1, \dots, j_\ell\}$, and each $\|R_\ell\|_F = \mathcal{O}_{\prec}(d^{-1/2})$ by Section 4.2 applied at the corresponding LOO level.

Step 2: Score decomposition. For each $k \in [m]$, sandwiching (50) with $u_{j_k}^\top \cdot v_i$ yields

$$X_k = u_{j_k}^\top \widehat{W}_{\text{fresh}} v_i + \mathcal{E}_k, \quad (51)$$

where $\mathcal{E}_k$ is the sum of $m+1$ rank-one bilinear contributions plus residual terms.

Step 3: Bound the corrections. Each rank-one contribution at level ℓ

$$\hat{a}_{i_\ell}^{\setminus I_\ell} u_{j_k}^\top (H^{\setminus I_\ell})^{-1} [u_{i_\ell} v_{i_\ell}^\top] v_i$$

admits a self-adjoint rewrite that places one of the held-out vectors — u_i for $\ell = 0$ and v_i for $\ell \geq 1$ — on the outer test side, independent of the LOO operator $(H^{\setminus I_\ell})^{-1}$ (which sees no sample in I_ℓ). Combined with the operator-norm bound $\|(H^{\setminus I_\ell})^{-1}\|_{\text{op}} \leq d^2/(\lambda n) = O(1)$ from the ridge floor and $\|u_a v_b^\top\|_{\text{F}} = O(1)$, the unmatched-Gaussian inner product gives an $\mathcal{O}_{\prec}(d^{-1/2})$ bound on each rank-one contribution. The residuals $u_{j_k}^\top R_\ell v_i$ are bounded by $\|R_\ell\|_{\text{F}} \|u_{j_k}\| \|v_i\| = \mathcal{O}_{\prec}(d^{-1/2})$. Summing $m+1$ such terms,

$$|\mathcal{E}_k| = \mathcal{O}_{\prec}(d^{-1/2}) \quad \text{uniformly in } k. \quad (52)$$

Step 4: Apply the bilinear coupling. Since $\widehat{W}_{\text{fresh}}$ is a function of $\{(u_l, v_l) : l \notin \{i, j_1, \dots, j_m\}\}$, it is independent of $u_{j_1}, \dots, u_{j_m}, v_i$. The bilinear coupling Corollary 1 requires a Frobenius self-averaging estimate and an operator-norm input for $\widehat{W}_{\text{fresh}}$; both are transferred from $\widehat{W}$ by the same finite-removal stability used above. In particular, the stochastic-dominance deviation $\|\widehat{W}_{\text{fresh}}\|_{\text{F}}^2/d^2 - \nu_d^2 = \mathcal{O}_{\prec}(d^{-1/2})$ follows from the Frobenius consequence of (A5), while the operator-norm input $\|(\widehat{W}_{\text{fresh}})^\top \widehat{W}_{\text{fresh}}\|_{\text{F}}/d^2 = \mathcal{O}_{\prec}(d^{-1/2})$ follows from (A4). Apply Corollary 1 with $W = \widehat{W}_{\text{fresh}}$, $v_0 = v_i$, and $\rho_f = \rho_{\text{op}} = O(d^{-1/2})$: there exist i.i.d. $Y_1, \dots, Y_m \sim \mathcal{N}(0, \nu_d^2)$ with

$$\|(u_{j_1}^\top \widehat{W}_{\text{fresh}} v_i, \dots, u_{j_m}^\top \widehat{W}_{\text{fresh}} v_i) - (Y_1, \dots, Y_m)\|_2 = \mathcal{O}_{\prec}(d^{-1/2}), \quad (53)$$

with $Y_1, \dots, Y_m$ i.i.d. $\mathcal{N}(0, \nu_d^2)$.

Step 5: Combine. Triangle inequality applied to (51), (52), and (53) yields (49). $\square$

4.4.2 Proximal law for diagonal scores

We next record the finite-dimensional proximal law for the diagonal scores. Set

$$\mu_d := \mu_r(\nu_d), \quad c_d := c_r(\nu_d),$$

where μ_r and c_r are defined in (39)–(40).

Proposition 3 (Coupled proximal law for s_{ii}). *Assume (A1)–(A6). Fix $i \in [n]$. On a common probability space, there is a standard Gaussian random variable $G_i \sim \mathcal{N}(0, 1)$ such that*

$$s_{i,i}(\widehat{W}) = c_d + \text{Prox}_{\chi_d}(\nu_d G_i - c_d) + \mathcal{O}_{\prec}(d^{-1/2}), \quad (54)$$

where Prox_{χ} is defined in (43).

Proof. Apply the LOO rank-one expansion (29) with $h = i$ and contract with $u_i^\top(\cdot)v_i$. This gives

$$s_{i,i}(\widehat{W}) = g_i + \hat{a}_i \langle u_i v_i^\top, (H^{\setminus i})^{-1} [u_i v_i^\top] \rangle + \mathcal{O}_{\prec}(d^{-1/2}), \quad (55)$$

where

$$g_i := u_i^\top \widehat{W}^{\setminus i} v_i, \quad \hat{a}_i = \sigma(-\hat{m}_i).$$

The LOO Hessian $H^{\setminus i}$ is independent of (u_i, v_i) . Lemma 17, (A6), and the finite-removal trace stability therefore give

$$\langle u_i v_i^\top, (H^{\setminus i})^{-1} [u_i v_i^\top] \rangle = \chi_d + \mathcal{O}_{\prec}(d^{-1/2}).$$

Since $\hat{a}_i$ is bounded, (55) becomes

$$s_{i,i}(\widehat{W}) = g_i + \chi_d \sigma(-s_{i,i}(\widehat{W}) + \hat{c}_i) + \mathcal{O}_{\prec}(d^{-1/2}), \quad (56)$$

where

$$\hat{c}_i := \hat{\mu}_i + \frac{1}{k} \sum_{j \neq i} \phi_\beta(s_{j,i}(\widehat{W}) - \hat{\mu}_i).$$

We now replace g_i and $\hat{c}_i$ by their finite-dimensional deterministic counterparts. The bilinear coupling estimate, applied to the leave-one-out matrix $\widehat{W}^{\setminus i}$ and the fresh pair (u_i, v_i) , gives a coupling

$$g_i = \nu_d G_i + \mathcal{O}_{\prec}(d^{-1/2}), \quad G_i \sim \mathcal{N}(0, 1). \quad (57)$$

Next, μ -stationarity gives

$$\frac{1}{n} \sum_{j \neq i} \sigma(\beta(s_{j,i}(\widehat{W}) - \hat{\mu}_i)) = r + \mathcal{O}_{\prec}(d^{-1/2}).$$

The off-diagonal Gaussian law from Proposition 2, together with the empirical-mean concentration postulate, implies that the left-hand side is

$$\mathbb{E}_{Z \sim \mathcal{N}(0,1)} \left[\sigma(\beta(\nu_d Z - \hat{\mu}_i)) \right] + \mathcal{O}_{\prec}(d^{-1/2}).$$

For fixed $r \in (0, 1)$ and fixed β , the bounded-score event keeps $\hat{\mu}_i$ and μ_d in a compact set; on this set the derivative in μ of the preceding expectation is bounded away from zero. Hence (40) yields

$$\hat{\mu}_i = \mu_d + \mathcal{O}_{\prec}(d^{-1/2}).$$

Applying the same Gaussian approximation to the pseudo-Lipschitz test function $\phi_\beta(\cdot - \mu)$, and using the Lipschitz dependence on μ , gives

$$\hat{c}_i = c_d + \mathcal{O}_{\prec}(d^{-1/2}). \quad (58)$$

Combining (56), (57), and (58), we obtain

$$s_{i,i}(\widehat{W}) = \nu_d G_i + \chi_d \sigma(-s_{i,i}(\widehat{W}) + c_d) + \mathcal{O}_{\prec}(d^{-1/2}).$$

Let $y_i := s_{i,i}(\widehat{W}) - c_d$. The last display is equivalent to

$$y_i - \chi_d \sigma(-y_i) = \nu_d G_i - c_d + \mathcal{O}_{\prec}(d^{-1/2}).$$

The map $y \mapsto y - \chi_d \sigma(-y)$ is strictly increasing with derivative at least one, so its inverse Prox_{χ_d} is 1-Lipschitz. This proves (54). $\square$

4.4.3 Finite-dimensional scalar equations and variational consistency

We now derive the two finite-dimensional scalar equations for ν_d and χ_d , and then identify their zero-residual limit with the saddle-point equations of the scalar variational problem. The variance equation is obtained directly from W -stationarity. Let G and Z be independent standard Gaussian variables, and define

$$S_d := c_d + \text{Prox}_{\chi_d}(\nu_d G - c_d), \quad A_d := \sigma(-S_d + c_d). \quad (59)$$

Proposition 4 (Finite-dimensional variance equation). *Assume (A1)–(A6), and use the coupled diagonal score law of Proposition 3. Then*

$$\nu_d^2 \left[\lambda + \frac{1}{r} \mathbb{E}[A_d] \mathbb{E} \left[\phi''_{\beta}(\nu_d Z - \mu_d) \right] \right] = \mathbb{E}[A_d S_d] + O(d^{-1/2}). \quad (60)$$

Proof. At the full minimizer, the W -stationarity condition (23) gives

$$UQ\Lambda_a V^\top + \frac{\lambda n}{d^2} \widehat{W} = 0.$$

Taking the Frobenius inner product with $\widehat{W}$ and using $s_{j,i}(\widehat{W}) = u_j^\top \widehat{W} v_i$ gives the exact identity

$$\lambda \frac{\|\widehat{W}\|_F^2}{d^2} = \frac{1}{n} \sum_{i=1}^n a_i \left(s_{i,i}(\widehat{W}) - \frac{1}{k} \sum_{j \neq i} q_{j,i} s_{j,i}(\widehat{W}) \right). \quad (61)$$

Taking expectations turns the left-hand side into $\lambda \nu_d^2$.

The first empirical average in (61) is a smooth test function of the diagonal score, after replacing c_i by c_d at the $d^{-1/2}$ scale. Proposition 3 and (A5) therefore yield

$$\mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n a_i s_{i,i}(\widehat{W}) \right] = \mathbb{E}[A_d S_d] + O(d^{-1/2}).$$

For the off-diagonal contribution, the competitor Gaussian law Proposition 2, the same empirical-mean concentration, and the leave-one-out independence between the diagonal factor and the competitor score give

$$Q_d := \sigma(\beta(\nu_d Z - \mu_d)).$$

$$\mathbb{E} \left[\frac{1}{n} \sum_{i=1}^n a_i \frac{1}{k} \sum_{j \neq i} q_{j,i} s_{j,i}(\widehat{W}) \right] = \frac{1}{r} \mathbb{E}[A_d] \mathbb{E}[\nu_d Z Q_d] + O(d^{-1/2}). \quad (62)$$

Here the distinction between k/n and $k/(n-1)$ contributes only $O(n^{-1})$, which is absorbed into $O(d^{-1/2})$. Combining (61)–(62) gives

$$\lambda \nu_d^2 = \mathbb{E}[A_d S_d] - \frac{1}{r} \mathbb{E}[A_d] \mathbb{E}[\nu_d Z Q_d] + O(d^{-1/2}).$$

Finally, Stein's identity for the standard Gaussian Z gives

$$\mathbb{E}[\nu_d Z Q_d] = \nu_d^2 \mathbb{E} \left[\phi''_{\beta}(\nu_d Z - \mu_d) \right],$$

which gives (60). □

We next derive the companion equation for the scalar χ_d defined in (48). It is the expectation of the normalized inverse trace of the full Hessian, and is often referred to as the susceptibility in the statistical physics literature. Set

$$B_d := A_d(1 - A_d).$$

Proposition 5 (Finite-dimensional susceptibility equation). *Assume (A1)–(A6). Then the finite-dimensional susceptibility χ_d satisfies*

$$1 = \alpha \lambda \chi_d + \alpha \mathbb{E} \left[\frac{\chi_d B_d}{1 + \chi_d B_d} \right] + \frac{\alpha \chi_d}{r} \mathbb{E}[A_d] \mathbb{E} \left[\phi''_{\beta}(\nu_d Z - \mu_d) \right] + O(d^{-1/2}). \quad (63)$$

In the above display, A_d is the random variable defined in (59).

Proof. For $j, i \in [n]$, write

$$f_{ji} := \text{vec}(u_j v_i^{\top}) \in \mathbb{R}^{d^2}.$$

Also let

$$r_{j,i} := \phi''_{\beta}(s_{j,i} - \mu_i) = \beta q_{j,i}(1 - q_{j,i}), \quad q_{j,i} = \sigma(\beta(s_{j,i} - \mu_i)),$$

so that $\tilde{R}_{j,i} = r_{j,i}/k$. Define the simplified Hessian

$$\tilde{H} = \frac{\lambda n}{d^2} I + \sum_{i=1}^n b_i f_{ii} f_{ii}^{\top} + \sum_{i=1}^n \sum_{j \neq i} \frac{a_i r_{j,i}}{k} f_{ji} f_{ji}^{\top}. \quad (64)$$

By Lemmas 18 and 19 in Appendix F, the normalized inverse trace of the original Hessian may be replaced, up to $\mathcal{O}_{\prec}(d^{-1/2})$, by the normalized inverse trace of $\tilde{H}$. Thus, by the trace concentration postulate (A6),

$$\frac{1}{d^2} \text{tr}(\tilde{H}^{-1}) = \chi_d + \mathcal{O}_{\prec}(d^{-1/2}). \quad (65)$$

We now use the identity

$$1 = \frac{1}{d^2} \text{tr}(\tilde{H} \tilde{H}^{-1}).$$

Expanding the three terms in (64) gives

$$\begin{aligned} 1 &= \frac{\lambda n}{d^2} \frac{1}{d^2} \text{tr}(\tilde{H}^{-1}) + \frac{1}{d^2} \sum_{i=1}^n b_i f_{ii}^{\top} \tilde{H}^{-1} f_{ii} \\ &\quad + \frac{1}{d^2} \sum_{i=1}^n \sum_{j \neq i} \frac{a_i r_{j,i}}{k} f_{ji}^{\top} \tilde{H}^{-1} f_{ji}. \end{aligned}$$

The first term is evaluated by (65). Lemmas 21 and 22 in Appendix F evaluate the two families of quadratic forms:

$$f_{ii}^{\top} \tilde{H}^{-1} f_{ii} = \frac{\chi_d}{1 + b_i \chi_d} + \mathcal{O}_{\prec}(d^{-1/2}), \quad f_{ji}^{\top} \tilde{H}^{-1} f_{ji} = \chi_d + \mathcal{O}_{\prec}(d^{-1/2}).$$

Substitution gives the finite-dimensional trace identity

$$1 = \frac{\lambda n}{d^2} \chi_d + \frac{1}{d^2} \sum_{i=1}^n b_i \frac{\chi_d}{1 + b_i \chi_d} + \frac{\chi_d}{d^2} \sum_{i=1}^n \sum_{j \neq i} \frac{a_i r_{j,i}}{k} + \mathcal{O}_{\prec}(d^{-1/2}). \quad (66)$$

Taking expectations under the moment bounds implicit in the postulates, and then applying the diagonal proximal law, the off-diagonal Gaussian law, and (A5), gives

$$\mathbb{E} \left[b_i \frac{\chi_d}{1 + b_i \chi_d} \right] = \mathbb{E} \left[\frac{\chi_d B_d}{1 + \chi_d B_d} \right] + O(d^{-1/2}),$$

and

$$\mathbb{E} \left[a_i \frac{1}{k} \sum_{j \neq i} r_{j,i} \right] = \frac{1}{r} \mathbb{E}[A_d] \mathbb{E} \left[\phi''_{\beta}(\nu_d Z - \mu_d) \right] + O(d^{-1/2}).$$

Substituting these two replacements into (66), and replacing n/d^2 by α at the same error scale, yields (63). $\square$

Combining Propositions 4 and 5 gives the finite-dimensional residual system

$$\begin{aligned} \nu_d^2 \left[\lambda + \frac{1}{r} \mathbb{E}[A_d] \mathbb{E} \left[\phi''_{\beta}(\nu_d Z - \mu_d) \right] \right] - \mathbb{E}[A_d S_d] &= O(d^{-1/2}), \\ 1 - \alpha \lambda \chi_d - \alpha \mathbb{E} \left[\frac{\chi_d B_d}{1 + \chi_d B_d} \right] - \frac{\alpha}{r} \mathbb{E}[A_d] \mathbb{E} \left[\phi''_{\beta}(\nu_d Z - \mu_d) \right] \chi_d &= O(d^{-1/2}). \end{aligned} \quad (67)$$

Here $\mu_d = \mu_r(\nu_d)$ is the deterministic threshold defined in (40), evaluated at the finite-dimensional variance parameter ν_d . Thus (ν_d, χ_d) is an *approximate* solution of the nonlinear scalar equations whose zero-residual limit turns out to be exactly equal to the saddle-point equations for the potential $\mathcal{P}_{\alpha,r}$ in Definition 3. This variational connection is established in the next proposition.

Proposition 6 (Variational form of the scalar equations). *For every $\nu > 0$, $\chi \mapsto \mathcal{P}_{\alpha,r}(\nu, \chi)$ has a unique maximizer. Moreover, the reduced potential $\mathcal{V}_{\alpha,r}$ is λ -strongly convex in ν . If ν_* is the unique minimizer of $\mathcal{V}_{\alpha,r}$ and χ_* is the unique maximizer of $\mathcal{P}_{\alpha,r}(\nu_*, \cdot)$, then the saddle-point equations of*

$$\inf_{\nu > 0} \sup_{\chi > 0} \mathcal{P}_{\alpha,r}(\nu, \chi)$$

are exactly the zero-residual limit of (67), with the finite-dimensional proximal channel replaced by (44) and with c_r given by (39).

Proof. See Appendix G. $\square$

5 Ridgeless limit and phase transition

The scalar theory in Section 4.3 gives a limiting variational problem for each fixed ridge parameter $\lambda > 0$ and fixed smoothing parameter $\beta < \infty$. In this section we take the

sequential ridgeless limit: first $n, d \rightarrow \infty$ with $n/d^2 \rightarrow \alpha$ and $k/n \rightarrow r$, and then $\lambda \downarrow 0$ while keeping $\beta \in (0, \infty)$ fixed. The limit separates two regimes. Below the critical load $\alpha_c(r)$ defined in (68) below, the scalar norm ν escapes to infinity as $\lambda \downarrow 0$ and the limiting fitting loss vanishes; we call this the satisfiable, or SAT, phase. Above this critical load the scalar norm remains finite and the limiting fitting loss is positive; we call this the unsatisfiable, or UNSAT, phase. After deriving the transition from the variational principle, we describe the score, margin, and percentile profiles on both sides.

All results in this section concern the scalar saddle problem of Definition 3. They translate into statements about the original TAM minimizer $(\widehat{W}, \widehat{\mu})$ through the conditional scalar identification in Result 2.

5.1 Phase transition

Definition 4 (Ridgeless threshold). *Let $G \sim N(0, 1)$, and define*

$$\kappa_r := \frac{\varphi(\Phi^{-1}(1-r))}{r},$$

where Φ and φ are the CDF and density of a standard normal random variable. The critical load is

$$\alpha_c(r) = \frac{1}{\mathbb{E}[(\kappa_r - G)_+^2]} = \frac{1}{(1 + \kappa_r^2)\Phi(\kappa_r) + \kappa_r\varphi(\kappa_r)}. \quad (68)$$

In what follows, we first explain the origin and underlying ideas behind the transition at $\alpha_c(r)$. The formal proof is then given in Proposition 7.

We start from the dual form of the reduced potential in (185), writing the ridge parameter explicitly:

$$\begin{aligned} \mathcal{V}_{\alpha,r,\lambda}(\nu) = \frac{\lambda}{2}\nu^2 + \sup_{0 \leq a(\cdot) \leq 1} \left\{ \mathbb{E} \left[a(G)(c_r(\nu) - \nu G) - I(a(G)) \right] \right. \\ \left. - \frac{\nu}{\sqrt{\alpha}} (\mathbb{E}[a(G)^2])^{1/2} \right\}, \end{aligned} \quad (69)$$

where $G \sim N(0, 1)$ and $I(a) = a \log a + (1-a) \log(1-a)$. The supremum is over measurable functions $a : \mathbb{R} \rightarrow [0, 1]$. We write $\mathcal{V}_{\alpha,r,0}$ for the ridgeless potential obtained by setting $\lambda = 0$; explicitly,

$$\mathcal{V}_{\alpha,r,0}(\nu) = \sup_{0 \leq a(\cdot) \leq 1} \left\{ \mathbb{E} \left[a(G)(c_r(\nu) - \nu G) - I(a(G)) \right] - \frac{\nu}{\sqrt{\alpha}} (\mathbb{E}[a(G)^2])^{1/2} \right\}. \quad (70)$$

The phase transition is determined by the large- ν behavior of the ridgeless potential $\mathcal{V}_{\alpha,r,0}$. The only input from the TAM threshold $c_r(\nu)$ needed for the leading-order calculation is the asymptotic relation

$$\frac{c_r(\nu)}{\nu} \longrightarrow \kappa_r. \quad (71)$$

We defer the verification of (71) to the proof of Proposition 7. Now set $m_r(\nu) := c_r(\nu)/\nu$. For fixed $a(\cdot)$, the linear-in- ν part of (69) is

$$\nu \left[\mathbb{E} \left[a(G)(m_r(\nu) - G) \right] - \frac{1}{\sqrt{\alpha}} \|a\|_2 \right], \quad \|a\|_2 := (\mathbb{E}[a(G)^2])^{1/2}.$$

The entropy term $I(a)$ is not multiplied by ν . Thus the large-norm behavior is governed by a simple question: can we choose a nonzero dual profile a so that the coefficient of the leading linear term in ν is positive?

Letting $\nu \rightarrow \infty$ in this coefficient and using (71), the relevant expression becomes

$$\mathbb{E}[a(G)(\kappa_r - G)] - \frac{1}{\sqrt{\alpha}} \|a\|_2.$$

Since $a \geq 0$, the negative part of $\kappa_r - G$ can only hurt, so the best profile should live on the event $\{G < \kappa_r\}$. More precisely, by Cauchy–Schwarz, for any $0 \leq a \leq 1$,

$$\mathbb{E}[a(G)(\kappa_r - G)] \leq \mathbb{E}[a(G)(\kappa_r - G)_+] \leq \|a\|_2 \left(\mathbb{E}[(\kappa_r - G)_+^2] \right)^{1/2}.$$

The upper bound is sharp despite the pointwise constraint $a \leq 1$. Indeed, equality in Cauchy–Schwarz would ask for a proportional to $(\kappa_r - G)_+$, which is unbounded. We can approximate this direction by the admissible truncations

$$a_M(G) := \min \left\{ \frac{(\kappa_r - G)_+}{M}, 1 \right\}, \quad M \rightarrow \infty.$$

The ratio $\mathbb{E}[a_M(G)(\kappa_r - G)]/\|a_M\|_2$ converges to $(\mathbb{E}[(\kappa_r - G)_+^2])^{1/2}$. Hence a positive large- ν direction exists exactly when $\alpha > \alpha_c(r)$, with $\alpha_c(r)$ as in (68).

Proposition 7 (Ridgeless scalar phase transition). *Fix $r \in (0, 1)$ and $\beta \in (0, \infty)$. For each $\lambda > 0$, let ν_λ be the unique minimizer of $\mathcal{V}_{\alpha, r, \lambda}$.*

(i) *If $\alpha < \alpha_c(r)$, then*

$$\nu_\lambda \rightarrow \infty \quad \text{and} \quad \mathcal{V}_{\alpha, r, 0}(\nu_\lambda) \rightarrow 0$$

as $\lambda \downarrow 0$.

(ii) *If $\alpha > \alpha_c(r)$, then ν_λ remains bounded as $\lambda \downarrow 0$ and converges to the unique minimizer $\nu_0 < \infty$ of $\mathcal{V}_{\alpha, r, 0}$. The ridgeless fitting loss $\mathcal{V}_{\alpha, r, 0}(\nu_0)$ is strictly positive.*

Proof. We first justify (71). Rescaling the definition of c_r by writing $\mu = \nu t$ gives

$$\frac{c_r(\nu)}{\nu} = \min_{t \in \mathbb{R}} \left\{ t + \frac{1}{r} \mathbb{E} \left[\frac{\phi_\beta(\nu(G - t))}{\nu} \right] \right\}.$$

For each fixed y , the function $\nu \mapsto \phi_\beta(\nu y)/\nu$ is nonincreasing and converges to y_+ . Thus the limiting problem is

$$\min_{t \in \mathbb{R}} \left\{ t + \frac{1}{r} \mathbb{E}[(G - t)_+] \right\}.$$

Its minimizer is $t = \Phi^{-1}(1 - r)$ and its value is κ_r , which proves (71). In fact the convergence is monotone decreasing.

SAT case: $\alpha < \alpha_c(r)$. By (71) and the definition (68), there are $\delta > 0$ and $\nu_0 < \infty$ such that, for all $\nu \geq \nu_0$,

$$(\mathbb{E}[(m_r(\nu) - G)_+^2])^{1/2} \leq \frac{1}{\sqrt{\alpha}} - \delta.$$

Therefore, uniformly over $0 \leq a \leq 1$,

$$\mathbb{E}[a(G)(m_r(\nu) - G)] - \frac{1}{\sqrt{\alpha}} \|a\|_2 \leq -\delta \|a\|_2.$$

The remaining terms in the dual representation are lower order. More precisely, for any $p \in (0, 1)$ there is $C_p < \infty$ such that $-I(x) \leq C_p x^p$ for $0 \leq x \leq 1$, while $\mathbb{E}[a^p] \leq \|a\|_2^p$. Hence

$$\mathcal{V}_{\alpha,r,0}(\nu) \leq \sup_{0 \leq x \leq 1} \{-\delta \nu x + C_p x^p\} \longrightarrow 0 \quad \text{as } \nu \rightarrow \infty.$$

Since $a = 0$ is admissible, $\mathcal{V}_{\alpha,r,0}(\nu) \geq 0$, and therefore $\mathcal{V}_{\alpha,r,0}(\nu) \rightarrow 0$ as $\nu \rightarrow \infty$.

To identify the variational value with the scalar fitting loss, fix ν and let $\chi(\nu)$ be the unique maximizer of $\mathcal{P}_{\alpha,r,\lambda}(\nu, \cdot)$, whose existence and uniqueness are given by Proposition 6. With the scalar proximal channel $S_{\nu,\chi}$ defined in (44), the identity

$$\mathcal{V}_{\alpha,r,\lambda}(\nu) = \frac{\lambda}{2} \nu^2 + \mathbb{E}[\ell(S_{\nu,\chi} - c_r(\nu))]$$

follows from a short cancellation. Writing $A := \sigma(-S_{\nu,\chi} + c_r(\nu))$, the proximal equation gives $S_{\nu,\chi} - \nu G = \chi A$, and hence

$$M_r(\nu, \chi) = \frac{\chi}{2} \mathbb{E}[A^2] + \mathbb{E}[\ell(S_{\nu,\chi} - c_r(\nu))] .$$

At the maximizing χ , the saddle equation $\nu^2 = \alpha \chi^2 \mathbb{E}[A^2]$ gives $\nu^2/(2\alpha\chi) = (\chi/2) \mathbb{E}[A^2]$, so the two quadratic terms in $\mathcal{P}_{\alpha,r,\lambda}$ cancel. Thus the ridgeless potential is the scalar fitting loss and is strictly positive at every finite ν . For any $\varepsilon > 0$, choose N so large that $\mathcal{V}_{\alpha,r,0}(N) \leq \varepsilon$, and then choose λ so small that $\lambda N^2/2 \leq \varepsilon$. This gives

$$\mathcal{V}_{\alpha,r,\lambda}(\nu_\lambda) \leq 2\varepsilon.$$

Hence the optimal value tends to zero. If ν_λ remained bounded along a subsequence, lower semicontinuity would force a finite point with zero ridgeless fitting loss, contradicting strict positivity. Therefore $\nu_\lambda \rightarrow \infty$, and the displayed identity implies that the per-sample fitting loss tends to zero.

UNSAT case: $\alpha > \alpha_c(r)$. By the sharpness of the Cauchy–Schwarz bound, there is a bounded measurable $a : \mathbb{R} \rightarrow [0, 1]$ and a number $\delta > 0$ such that

$$\mathbb{E}[a(G)(\kappa_r - G)] - \frac{1}{\sqrt{\alpha}} \|a\|_2 \geq 2\delta.$$

Using $m_r(\nu) \rightarrow \kappa_r$ and bounded convergence, the same a satisfies, for all sufficiently large ν ,

$$\mathbb{E}[a(G)(m_r(\nu) - G)] - \frac{1}{\sqrt{\alpha}} \|a\|_2 \geq \delta.$$

Substituting this fixed test function into (69) gives

$$\mathcal{V}_{\alpha,r,0}(\nu) \geq \delta \nu - O(1),$$

so the ridgeless potential is coercive. The minimizers of $\mathcal{V}_{\alpha,r,\lambda} = \mathcal{V}_{\alpha,r,0} + \lambda\nu^2/2$ therefore remain in a compact set as $\lambda \downarrow 0$. Any limit point minimizes $\mathcal{V}_{\alpha,r,0}$ by the standard variational argument: compare ν_λ against an arbitrary fixed ν and let $\lambda \downarrow 0$.

We now show that the ridgeless minimizer is unique. For finite β , the TAM threshold c_r is strictly convex. Indeed, with

$$w_\nu(Z) := \phi''_\beta(\nu Z - \mu_r(\nu)),$$

the envelope theorem and the defining equation for $\mu_r(\nu)$ give

$$c_r''(\nu) = \frac{1}{r} \left\{ \mathbb{E}[w_\nu(Z)Z^2] - \frac{\mathbb{E}[w_\nu(Z)Z]^2}{\mathbb{E}[w_\nu(Z)]} \right\} > 0.$$

For each finite ν , the dual maximizer in (70) has positive expectation. In the primal variables this maximizer is $a(G) = \sigma(-S_{\nu,\chi} + c_r(\nu))$, and hence $0 < a(G) < 1$ almost surely. Therefore $\mathcal{V}_{\alpha,r,0}$ is strictly convex: if $\nu_\theta = \theta\nu_1 + (1-\theta)\nu_2$ with $0 < \theta < 1$ and $\nu_1 \neq \nu_2$, evaluating the dual formula at the maximizer for ν_θ and using the strict convexity of c_r gives

$$\mathcal{V}_{\alpha,r,0}(\nu_\theta) < \theta\mathcal{V}_{\alpha,r,0}(\nu_1) + (1-\theta)\mathcal{V}_{\alpha,r,0}(\nu_2).$$

Thus the ridgeless minimizer is unique; call it ν_0 . Hence $\nu_\lambda \rightarrow \nu_0$. Since ν_0 is finite, the corresponding scalar logistic loss is strictly positive. $\square$

Remark 4 (No dependence on the smoothing parameter). *The critical load $\alpha_c(r)$ does not depend on the smoothing parameter $\beta \in (0, \infty)$. The parameter β affects the finite- λ scalar channel and the finite ridgeless minimizer in the UNSAT phase, but it disappears from the large- ν limit (71), where the smoothed logistic tail $\phi_\beta(\nu y)/\nu$ converges to y_+ .*

5.2 Score, margin, and percentile profiles

We next translate the ridgeless phase transition into predictions for the true scores, the competing scores, the margins, and the percentile rank of the true score among its competitors. The two phases have different scaling behavior. In the SAT phase, Proposition 7 shows that $\nu_\lambda \rightarrow \infty$. Thus the true scores and the competitor scores both diverge on the original scale. The meaningful objects are the normalized scores, obtained by dividing by ν_λ , and these have a non-degenerate limit. In the UNSAT phase, by contrast, ν_λ remains finite, so the ridgeless profiles live on the original scale.

5.2.1 The SAT phase

We begin with the SAT phase, where the limiting normalized profiles are explicit. The single parameter controlling them is the number $\rho_\alpha > \kappa_r$, defined as the unique solution of

$$\mathbb{E}[(\rho_\alpha - G)_+^2] = \frac{1}{\alpha}, \quad G \sim N(0, 1). \quad (72)$$

The left-hand side is strictly increasing in ρ_α , and the boundary value $\rho_\alpha = \kappa_r$ is exactly (68). Hence ρ_α is monotone decreasing in α and satisfies $\rho_\alpha \downarrow \kappa_r$ as $\alpha \uparrow \alpha_c(r)$.

Proposition 8 (SAT score, margin, and percentile profiles). *Assume $\alpha < \alpha_c(r)$, and let $\rho_\alpha > \kappa_r$ be the unique solution of (72). In the sequential ridgeless limit $\lambda \downarrow 0$, the normalized true score satisfies*

$$\frac{S_{\nu_\lambda, \chi_\lambda}}{\nu_\lambda} \Rightarrow \max\{G, \rho_\alpha\}, \quad G \sim N(0, 1). \quad (73)$$

Consequently, the normalized TAM margin satisfies

$$\frac{S_{\nu_\lambda, \chi_\lambda} - c_r(\nu_\lambda)}{\nu_\lambda} \Rightarrow \max\{G, \rho_\alpha\} - \kappa_r, \quad G \sim N(0, 1). \quad (74)$$

Here and below, $\Rightarrow$ denotes weak convergence of laws as $\lambda \downarrow 0$. The associated percentile variable satisfies

$$\Omega \Rightarrow \Phi(\max\{G, \rho_\alpha\}). \quad (75)$$

Equivalently, if $p_\alpha := \Phi(\rho_\alpha)$, then

$$\mathbb{P}(\Omega \leq \omega) = \begin{cases} 0, & \omega < p_\alpha, \\ \omega, & p_\alpha \leq \omega \leq 1. \end{cases} \quad (76)$$

Proof. By Proposition 7, in the SAT phase $\nu_\lambda \rightarrow \infty$ and the scalar fitting loss tends to zero. The proof has three steps. First, we rewrite the proximal channel in terms of the normalized score and the correction term created by the loss. Second, we introduce the scalar quantity ρ_λ and use the saddle-point equation to show that it is bounded. Third, we identify every subsequential limit of ρ_λ ; uniqueness of the equation defining ρ_α then gives the stated weak convergence.

Set $c_\lambda := c_r(\nu_\lambda)$. For each deterministic scalar input $g \in \mathbb{R}$, define

$$S_\lambda(g) := c_\lambda + \text{Prox}_{\chi_\lambda}(\nu_\lambda g - c_\lambda), \quad X_\lambda(g) := \frac{S_\lambda(g)}{\nu_\lambda}.$$

The proximal first-order condition is the pointwise identity

$$S_\lambda(g) = \nu_\lambda g + \chi_\lambda \sigma(-S_\lambda(g) + c_\lambda).$$

We also define the pointwise correction

$$B_\lambda(g) := X_\lambda(g) - g = \frac{\chi_\lambda}{\nu_\lambda} \sigma(-S_\lambda(g) + c_\lambda) \geq 0. \quad (77)$$

Only now do we insert the Gaussian variable: when $G \sim N(0, 1)$, the random variables in the scalar channel are $S_\lambda(G)$, $X_\lambda(G)$, and $B_\lambda(G)$. The χ saddle-point equation for the scalar potential in Definition 3, recorded in (181) of Appendix G, gives

$$\mathbb{E}[B_\lambda(G)^2] = \frac{1}{\alpha}. \quad (78)$$

Moreover, the zero-loss statement implies that the normalized score does not fall below the TAM threshold. Indeed, with $M_\lambda(g) := S_\lambda(g) - c_\lambda$, we have pointwise

$$\left(\frac{c_\lambda}{\nu_\lambda} - X_\lambda(g) \right)_+ = \frac{(-M_\lambda(g))_+}{\nu_\lambda}.$$

For every $m \in \mathbb{R}$,

$$(-m)_+ \leq \ell(m).$$

Therefore, since the fitting loss $\mathbb{E}[\ell(M_\lambda)]$ tends to zero and $\nu_\lambda \rightarrow \infty$,

$$\mathbb{E} \left[\left(\frac{c_\lambda}{\nu_\lambda} - X_\lambda(G) \right)_+ \right] \rightarrow 0.$$

Since $c_\lambda/\nu_\lambda \rightarrow \kappa_r$ by (71), the 1-Lipschitz property of $x \mapsto x_+$ gives

$$\mathbb{E} \left[(\kappa_r - X_\lambda(G))_+ \right] \rightarrow 0. \quad (79)$$

We now identify the limit of the channel. The useful normalization is the level at which the sigmoid correction in (77) changes from negligible to order one. Define

$$\rho_\lambda := \frac{c_\lambda + \log(\chi_\lambda/\nu_\lambda)}{\nu_\lambda}.$$

Combining this definition with (77) and the identity $\sigma(x) = (1 + \exp\{-x\})^{-1}$ gives the exact formula

$$B_\lambda(g) = \frac{\exp\{\nu_\lambda(\rho_\lambda - X_\lambda(g))\}}{1 + \exp\{c_\lambda - \nu_\lambda X_\lambda(g)\}}. \quad (80)$$

The sequence ρ_λ is bounded. Indeed, if $\rho_\lambda \rightarrow -\infty$ along a subsequence, then $\chi_\lambda/\nu_\lambda = \exp\{\nu_\lambda \rho_\lambda - c_\lambda\} \rightarrow 0$, and hence $B_\lambda(g) \leq \chi_\lambda/\nu_\lambda \rightarrow 0$ uniformly in g , contradicting (78). Next suppose that $\rho_\lambda \rightarrow +\infty$ along a subsequence. We claim that, for all sufficiently small λ ,

$$B_\lambda(g) \geq \frac{\rho_\lambda}{2} \quad \text{for every } g \in [-1, 0].$$

Indeed, if the reverse inequality held at some $g \in [-1, 0]$, then $X_\lambda(g) = g + B_\lambda(g) \leq \rho_\lambda/2$. Using (80) and the boundedness of c_λ/ν_λ , we would get

$$B_\lambda(g) \geq \exp\{\nu_\lambda \rho_\lambda/3\}$$

for all sufficiently small λ , contradicting $B_\lambda(g) < \rho_\lambda/2$. Hence the claim holds. Since $\mathbb{P}(-1 \leq G \leq 0) = \Phi(0) - \Phi(-1) > 0$, the claim implies

$$\mathbb{E}[B_\lambda(G)^2] \geq (\Phi(0) - \Phi(-1)) \frac{\rho_\lambda^2}{4} \rightarrow \infty,$$

contradicting (78).

It remains to identify the possible limits of ρ_λ . Take a subsequence along which $\rho_\lambda \rightarrow \rho$. Passing to a further subsequence, we may assume from (79) that

$$\liminf_{\lambda \downarrow 0} X_\lambda(G) \geq \kappa_r \quad \text{almost surely.} \quad (81)$$

The rest of the argument is pointwise in g . First, if $g > \rho$, then $B_\lambda(g) \rightarrow 0$ and $X_\lambda(g) \rightarrow g$. Indeed, any positive subsequential limit of $B_\lambda(g)$ would force $X_\lambda(g) = g + B_\lambda(g) > \rho_\lambda$

eventually, and the right-hand side of (80) would then converge to zero. This excludes $\rho < \kappa_r$, because on the positive-probability event $\{\rho < G < \kappa_r\}$ it would imply $X_\lambda(G) \rightarrow G < \kappa_r$, contradicting (81). Hence $\rho \geq \kappa_r$.

We now identify the channel limit for any subsequential limit $\rho \geq \kappa_r$. Along any bounded subsequence of ρ_λ , (80) gives the domination

$$B_\lambda(g)^2 \leq 1 + \left(\sup_\lambda \rho_\lambda - g \right)_+^2. \quad (82)$$

Indeed, if $B_\lambda(g) \leq 1$ there is nothing to prove; if $B_\lambda(g) > 1$, then (80) implies $X_\lambda(g) \leq \rho_\lambda$, and so $B_\lambda(g) = X_\lambda(g) - g \leq (\rho_\lambda - g)_+$.

For $g > \rho$, we already proved $X_\lambda(g) \rightarrow g$. Now fix $g < \rho$. For almost every such g , every subsequential limit x of $X_\lambda(g)$ satisfies $x \geq \kappa_r$ by (81). If $x < \rho$, then the exponential numerator in (80) grows like $\exp\{\nu_\lambda(\rho - x + o(1))\}$, while the denominator grows at most like $\exp\{\nu_\lambda(\kappa_r - x)_+ + o(\nu_\lambda)\}$. Since $\rho \geq \kappa_r$ and $x \geq \kappa_r$, this is impossible. If $x > \rho$, the correction term vanishes by (80), forcing $x = g$, also impossible because $g < \rho$. Hence $x = \rho$. Thus, with $G \sim N(0, 1)$,

$$X_\lambda(G) \longrightarrow \max\{G, \rho\}, \quad B_\lambda(G) \longrightarrow (\rho - G)_+.$$

The domination bound (82) allows us to pass to the second moment in (78), so

$$\mathbb{E}[(\rho - G)_+]^2 = \frac{1}{\alpha}.$$

By uniqueness of the solution to (72), every subsequential limit is ρ_α . Hence the whole sequence satisfies $X_\lambda(G) \Rightarrow \max\{G, \rho_\alpha\}$. Subtracting $c_\lambda/\nu_\lambda \rightarrow \kappa_r$ proves (73) and (74).

The percentile distribution is obtained by comparing the diagonal score with the limiting competitor field $N(0, \nu_\lambda^2)$. Hence (75). The CDF formula (76) follows by noting that $\Phi(\max\{G, \rho_\alpha\})$ has an atom of mass $p_\alpha = \Phi(\rho_\alpha)$ at p_α , followed by a uniform tail on $(p_\alpha, 1)$. $\square$

5.2.2 The UNSAT phase

Assume $\alpha > \alpha_c(r)$. In this regime the ridgeless limit does not escape to infinite norm. By Proposition 7, the minimizers ν_λ converge to the unique finite minimizer ν_0 of the ridgeless reduced potential. The next proposition records the corresponding finite-scale profiles. We omit the proof, since the statement is obtained by specializing the scalar channel from Section 4.3 at the ridgeless minimizer.

Proposition 9 (UNSAT score, margin, and percentile profiles). *Let χ_0 be the maximizer in the scalar saddle problem corresponding to ν_0 , and define*

$$c_0 := c_r(\nu_0).$$

Then the limiting true score is the finite-scale scalar random variable

$$S_0 := S_{\nu_0, \chi_0} = c_0 + \text{Prox}_{\chi_0}(\nu_0 G - c_0), \quad G \sim N(0, 1).$$




Figure 1: Finite-dimensional validation of the predicted phase boundary. The solid curve is the theoretical threshold $\alpha_c(r)$ from (68), and open circles show empirical boundary estimates from the numerical sweep. **a.** Optimized TAM fitting loss per sample, after subtracting the ridge contribution; colors are clipped at one. **b.** Norm scale $\nu = \|\widehat{W}\|_F/d$ on a logarithmic color scale; colors are clipped at 100. Both panels use $d = 400$ and small ridge regularization $\lambda = 1e - 6$. The observed transition is sharply aligned with the scalar prediction.

The competing scores remain on the same finite scale, with limiting Gaussian scale ν_0 . The limiting TAM margin and fitting loss are

$$M_0 := S_0 - c_0, \quad L_0 := \mathbb{E}[\ell(M_0)] > 0.$$

Finally, the limiting percentile rank of the true score is

$$\Omega_0 := \Phi\left(\frac{S_0}{\nu_0}\right).$$

Remark 5. Unlike the phase boundary and the SAT normalized profiles, these UNSAT quantities generally retain a dependence on the fixed value of β through the scalar functions $c_r(\nu)$ and $S_{\nu,\chi}$.

5.3 Numerical validation

We now give finite-dimensional numerical checks of the scalar predictions. The most direct prediction is the phase boundary of Proposition 7: as (α, r) crosses the curve $\alpha = \alpha_c(r)$, the optimized fitting loss should separate from zero and the norm scale $\nu = \|\widehat{W}\|_F/d$ should change sharply. Figure 1 tests this prediction in a two-dimensional sweep over the load $\alpha = n/d^2$ and the tail scale $r = k/(n-1)$. The empirical transition extracted from the simulations is closely aligned with the theoretical curve (68), both in the loss landscape and in the growth of the norm scale.

The phase diagram gives a global view of the transition. We next look more locally, along a one-dimensional slice in α (with $r = 0.15$), and compare the finite-dimensional optimizer

with the scalar predictions on both sides of the transition. This second comparison provides more information: it tests not only the location of the phase boundary, but also the predicted loss, norm scale, margin profile, and percentile distribution.

Figure 2 shows good agreement between the scalar theory and the finite-dimensional simulations. The upper panels compare two scalar observables: the fitting loss and the inverse norm scale $1/\nu$. The loss remains close to zero throughout the SAT phase and becomes positive beyond the predicted critical load. At the same transition, $1/\nu$ collapses toward zero on the SAT side, while it remains positive in the UNSAT phase, as predicted by Proposition 7.

The lower panels compare distributional predictions in the SAT phase. The empirical normalized margin distribution follows the limiting law (74), with the visible rounding caused by finite dimension. The percentile panel is particularly informative. The TAM prediction (76) has a hard lower edge at $p_\alpha = \Phi(\rho_\alpha)$, meaning that in the ridgeless SAT limit the true item is never predicted to fall below this percentile among its competitors. The finite-dimensional curve tracks this edge closely. The same panel also shows the correlation-memory baseline (20); compared with this non-optimized baseline, the TAM optimizer shifts the signal percentiles sharply toward the upper tail, reflecting the improvement obtained by learning the memory matrix for the listwise objective.




Figure 2: Scalar theory compared with finite-dimensional simulations along a one-dimensional slice at $r = 0.15$. **a.** Optimized TAM fitting loss per sample as a function of the load α , after subtracting the ridge contribution. **b.** Inverse norm scale $1/\nu$, showing the collapse of $1/\nu$ in the SAT phase and its positive value in the UNSAT phase. In both upper panels, the solid curve is obtained from the scalar prediction (42), with the norm and loss read off through (45) and (46), at $\beta = 30$ and ridge $\lambda = 10^{-7}$. Open circles are finite-dimensional simulations at $d = 600$, and the dashed vertical line marks the ridgeless threshold $\alpha_c(r)$. **c.** Empirical CDF of the normalized TAM margin $(S_{ii} - c_r(\nu))/\nu$ in the SAT phase, compared with (74). **d.** Empirical CDF of the signal percentile Ω_i , compared with (76) and the correlation-memory baseline (20). The lower panels use the representative SAT load $\alpha = 0.28$.

5.4 A small-tail conjecture for top-1 retrieval

The scalar theory above is derived for a fixed tail fraction $r \in (0, 1)$. It therefore does not directly cover top-1 retrieval, which corresponds to the extreme-tail choice $r = 1/(n - 1)$. Still, formally extrapolating the fixed- r phase boundary to this vanishing tail fraction gives an intriguing prediction for the sharp top-1 constant.

As $r \downarrow 0$,

$$\kappa_r = \frac{\varphi(\Phi^{-1}(1 - r))}{r} \sim \sqrt{2 \log(1/r)}$$

and therefore

$$\alpha_c(r) = \frac{1}{\mathbb{E}[(\kappa_r - G)_+^2]} \sim \frac{1}{2 \log(1/r)}.$$

Setting $r = 1/(n - 1)$ makes TAM coincide with the top-1 criterion, at least formally. Since the TAM load is $\alpha = n/d^2$, the condition $\alpha < \alpha_c(r)$ then becomes

$$\frac{n}{d^2} \lesssim \frac{1}{2 \log n}, \quad \text{or equivalently} \quad d^2 \gtrsim 2n \log n.$$

Conjecture 1 (Small-tail extrapolation to top-1 retrieval). *Along sequences with $n, d \rightarrow \infty$, the optimal top-1 retrieval threshold is*

$$\frac{d^2}{n \log n} = 2. \tag{83}$$

In other words, for every fixed $\epsilon > 0$, top-1 retrieval should be feasible with high probability when $d^2 > (2 + \epsilon)n \log n$, and infeasible with high probability when $d^2 < (2 - \epsilon)n \log n$.

This conjecture is consistent with the logarithmic scale proved in Section 2, but predicts a sharper constant than what is currently proved. The same sharp constant was proposed concurrently by Giorlandino et al. [2026] based on a simplified decoupled model of linear associative memory. It is also supported by numerical experiments in the top-1 regime. We empirically estimate the threshold by optimizing the regularized multiclass cross-entropy objective (19) with a small ridge parameter, while sweeping the rescaled load

$$c = \frac{n \log n}{d^2}.$$

For each dimension, the norm scale $\|\widehat{W}\|_{\mathbb{F}}/d$ develops a pronounced peak near the transition. We use the location of this peak as a finite-dimensional estimate $\widehat{c}(d)$ of the critical load.

Figure 3 shows strong finite-dimensional effects, as expected in the extreme-tail regime: the observed peaks occur at values of c above $1/2$ for the simulated dimensions. However, the drift with d is systematic. A linear fit in $1/\log d$ gives an intercept close to $1/2$, matching the formal small-tail extrapolation above. We therefore view Conjecture 1 as a tantalizing consequence of the scalar theory of TAM retrieval.




Figure 3: Numerical evidence for the small-tail top-1 conjecture. **a.** Norm scale $\|\widehat{W}\|_F/d$ for the cross-entropy top-1 objective (19), plotted against $c = n \log n / d^2$. Stars mark the largest norm value for each dimension, which we use as the empirical threshold $\widehat{c}(d)$. **b.** Extrapolation of these empirical thresholds against $1/\log d$. The linear fit gives $\widehat{c}(d) \approx 0.507 + 1.63/\log d$, close to the conjectural limit $c_\infty = 1/2$, equivalently the threshold (83), $d^2/(n \log n) \rightarrow 2$.

6 Conclusion

We have shown that the capacity of linear associative memory is governed not only by the number of degrees of freedom in the memory matrix, but also by the retrieval statistic used to read it out. For exact top-1 retrieval, the relevant statistic is the largest distractor. This extreme-value requirement forces the logarithmic model-size scale

$$d^2 \asymp n \log n.$$

The correlation matrix memory achieves this scale, and our universal impossibility theorem shows that the logarithmic factor cannot be removed by choosing a more sophisticated linear memory. It is therefore not an artifact of the specific construction of correlation matrix memory, but the intrinsic price of winner-take-all decoding.

The picture changes for listwise retrieval. We propose the Tail-Average Margin criterion, which replaces the maximum competitor by the average strength of a fixed upper tail. This still certifies that the correct association remains in a controlled candidate list, but it avoids making success depend on a single extreme distractor. At fixed tail fraction, the resulting theory operates at the quadratic scale

$$d^2 \asymp n,$$

the scale suggested by the number of parameters. Conditional on six precise postulates stated in Section 4, we show that the TAM optimizer admits a scalar variational description that predicts the limiting loss, score laws, margin distribution, percentile profile, and ridgeless phase transition. This scalar theory reveals a rich phenomenology: a satisfiable/unsatisfiable

transition, explicit limiting score profiles, and percentile laws for where the true association sits among its competitors. Proving the six postulates would turn this conditional prediction into a fully rigorous capacity theory for listwise associative memory.

The fixed-tail theory also points back toward exact retrieval. Formally sending the tail fraction r to the extreme value $1/(n-1)$ gives a tantalizing small-tail conjecture: the sharp top-1 threshold should be

$$d^2 \sim 2n \log n,$$

consistent with the concurrent prediction of Giorlandino et al. [2026], based on a simplified decoupled model, and with the finite-dimensional simulations in Section 5. Making this prediction rigorous requires understanding a singular extreme-tail limit, where the fixed-tail asymptotics and the top-1 extreme-value regime meet.

References

- Zeyuan Allen-Zhu and Yuanzhi Li. Physics of language models: part 3.3, knowledge capacity scaling laws, 2024. URL <https://arxiv.org/abs/2404.05405>.
- S.-I. Amari. Learning patterns and pattern sequences by self-organizing nets of threshold elements. *IEEE Transactions on Computers*, C-21(11):1197–1206, 1972. doi: 10.1109/T-C.1972.223477.
- James A. Anderson. A simple neural network generating an interactive memory. *Mathematical Biosciences*, 14(3):197–220, 1972. ISSN 0025-5564. doi: 10.1016/0025-5564(72)90075-2.
- Benjamin Aubin, Florent Krzakala, Yue M. Lu, and Lenka Zdeborová. Generalization error in high-dimensional perceptrons: Approaching Bayes error with convex optimization. In *Advances in Neural Information Processing Systems*, 2020.
- Jean Barbier, Florent Krzakala, Nicolas Macris, Léo Miolane, and Lenka Zdeborová. Optimal errors and phase transitions in high-dimensional generalized linear models. *Proceedings of the National Academy of Sciences*, 116(12):5451–5460, 2019. doi: 10.1073/pnas.1802705116.
- Alberto Bietti, Vivien Cabannes, Diane Bouchacourt, Herve Jegou, and Leon Bottou. Birth of a transformer: a memory viewpoint, 2023. URL <https://arxiv.org/abs/2306.00802>.
- Trenton Bricken and Cengiz Pehlevan. Attention approximates sparse distributed memory. In *Advances in Neural Information Processing Systems*, 2021.
- Vivien Cabannes, Elvis Dohmatob, and Alberto Bietti. Scaling laws for associative memories, 2024a. URL <https://arxiv.org/abs/2310.02984>.
- Vivien Cabannes, Berfin Simsek, and Alberto Bietti. Learning associative memories with gradient descent, 2024b. URL <https://arxiv.org/abs/2402.18724>.

- Zhe Cao, Tao Qin, Tie-Yan Liu, Ming-Feng Tsai, and Hang Li. Learning to rank: from pairwise approach to listwise approach. In *Proceedings of the 24th International Conference on Machine Learning*, pages 129–136. ACM, 2007. doi: 10.1145/1273496.1273513.
- Thomas M. Cover. Geometrical and statistical properties of systems of linear inequalities with applications in pattern recognition. *IEEE Transactions on Electronic Computers*, EC-14(3):326–334, 1965. doi: 10.1109/PGEC.1965.264137.
- Mete Demircigil, Judith Heusel, Matthias Löwe, Sven Uppang, and Franck Vermet. On a model of associative memory with huge storage capacity. *Journal of Statistical Physics*, 168(2):288–299, 2017.
- Noureddine El Karoui. On the impact of predictor geometry on the performance on high-dimensional ridge-regularized generalized robust regression estimators. *Probability Theory and Related Fields*, 170(1–2):95–175, 2018. doi: 10.1007/s00440-016-0754-9.
- Noureddine El Karoui, Derek Bean, Peter J. Bickel, Chinghway Lim, and Bin Yu. On robust regression with high-dimensional predictors. *Proceedings of the National Academy of Sciences*, 110(36):14557–14562, 2013. doi: 10.1073/pnas.1307842110.
- Nelson Elhage, Tristan Hume, Catherine Olsson, Nicholas Schiefer, Tom Henighan, Shauna Kravec, Zac Hatfield-Dodds, Robert Lasenby, Dawn Drain, Carol Chen, Roger Grosse, Sam McCandlish, Jared Kaplan, Dario Amodei, Martin Wattenberg, and Christopher Olah. Toy models of superposition, 2022. URL <https://arxiv.org/abs/2209.10652>.
- Peter Elias. List decoding for noisy channels. Technical Report 335, Research Laboratory of Electronics, Massachusetts Institute of Technology, Cambridge, MA, 1957.
- László Erdős and Horng-Tzer Yau. *A dynamical approach to random matrix theory*, volume 28 of *Courant Lecture Notes in Mathematics*. American Mathematical Society, 2017.
- Viola Folli, Marco Leonetti, and Giancarlo Ruocco. On the maximum storage capacity of the Hopfield model. *Frontiers in Computational Neuroscience*, 10:144, 2017. ISSN 1662-5188. doi: 10.3389/fncom.2016.00144.
- E. Gardner. Maximum storage capacity in neural networks. *Europhysics Letters*, 4(4):481–485, aug 1987. doi: 10.1209/0295-5075/4/4/016.
- E. Gardner. The space of interactions in neural network models. *Journal of Physics A: Mathematical and General*, 21(1):257–270, 1988. doi: 10.1088/0305-4470/21/1/030.
- Samuel J. Gershman, Ila Fiete, and Kazuki Irie. Key-value memory in the brain, 2025. URL <https://arxiv.org/abs/2501.02950>.
- Mor Geva, Roei Schuster, Jonathan Berant, and Omer Levy. Transformer feed-forward layers are key-value memories, 2021. URL <https://arxiv.org/abs/2012.14913>.
- Alessio Giorlandino, Sebastian Goldt, and Antoine Maillard. Factual recall in linear associative memories: sharp asymptotics and mechanistic insights, 2026. arXiv preprint.

- Alex Graves, Greg Wayne, and Ivo Danihelka. Neural Turing machines, 2014. URL <https://arxiv.org/abs/1410.5401>.
- Venkatesan Guruswami. Algorithmic results in list decoding. *Foundations and Trends in Theoretical Computer Science*, 2(2):107–195, 2007. doi: 10.1561/0400000007.
- D. O. Hebb. *The organization of behavior: a neuropsychological theory*. John Wiley & Sons, New York, 1949.
- Stephen Hobson. *Correlation matrix memories: improving performance for capacity and generalisation*. PhD thesis, University of York, September 2011. URL <https://etheses.whiterose.ac.uk/id/eprint/2211/>.
- J.J. Hopfield. Neural networks and physical systems with emergent collective computational abilities. *Proceedings of the National Academy of Sciences*, 79:2554–8, April 1982. doi: 10.1073/pnas.79.8.2554.
- J.J. Hopfield. Neurons with graded response have collective computational properties like those of two-state neurons. *Proceedings of the National Academy of Sciences*, 81:3088–92, May 1984. doi: 10.1073/pnas.81.10.3088.
- Yibo Jiang, Goutham Rajendran, Pradeep Ravikumar, and Bryon Aragam. Do LLMs dream of elephants (when told not to)? Latent concept association and associative memory in transformers, 2024. URL <https://arxiv.org/abs/2406.18400>.
- Pentti Kanerva. *Sparse distributed memory*. MIT Press, Cambridge, MA, 1988.
- Ryotaro Kawata and Taiji Suzuki. Transformers as measure-theoretic associative memory: a statistical perspective and minimax optimality, 2026. URL <https://arxiv.org/abs/2602.01863>.
- Juno Kim, Eshaan Nichani, Denny Wu, Alberto Bietti, and Jason D. Lee. Sharp capacity scaling of spectral optimizers in learning associative memory, 2026. URL <https://arxiv.org/abs/2603.26554>.
- Teuvo Kohonen. Correlation matrix memories. *IEEE Transactions on Computers*, C-21: 353–359, 1972. doi: 10.1109/TC.1972.5008975.
- B. Kosko. Bidirectional associative memories. *IEEE Transactions on Systems, Man, and Cybernetics*, 18(1):49–60, 1988. doi: 10.1109/21.87054.
- Dmitry Krotov and John J Hopfield. Dense associative memory for pattern recognition, 2016. URL <https://arxiv.org/abs/1606.01164>.
- Lihua Lei, Peter J. Bickel, and Nouredine El Karoui. Asymptotics for high dimensional regression M-estimates: fixed design results. *Probability Theory and Related Fields*, 172(3–4):983–1079, 2018. doi: 10.1007/s00440-017-0824-7.

- Carlo Lucibello and Marc Mézard. Exponential capacity of dense associative memories. *Physical Review Letters*, 132(7):077301, February 2024. doi: 10.1103/PhysRevLett.132.077301.
- Robert J. McEliece, Edward C. Posner, Eugene R. Rodemich, and Santosh S. Venkatesh. The capacity of the hopfield associative memory. *IEEE Transactions on Information Theory*, 33:461–482, 1987.
- Kevin Meng, David Bau, Alex Andonian, and Yonatan Belinkov. Locating and editing factual associations in GPT. In *Advances in Neural Information Processing Systems*, 2022.
- Marc Mézard, Giorgio Parisi, and Miguel Angel Virasoro. *Spin Glass Theory and Beyond*. World Scientific, 1987. doi: 10.1142/0271.
- Francesca Mignacco, Florent Krzakala, Yue Lu, Pierfrancesco Urbani, and Lenka Zdeborova. The role of regularization in classification of high-dimensional noisy gaussian mixture. In *International conference on machine learning*, pages 6874–6883, 2020.
- Beren Millidge, Tommaso Salvatori, Yuhang Song, Thomas Lukasiewicz, and Rafal Bogacz. Universal Hopfield networks: a general framework for single-shot associative memory models, 2022. URL <https://arxiv.org/abs/2202.04557>.
- Kaoru Nakano. Associatron-A model of associative memory. *IEEE Transactions on Systems, Man Cybernetics*, SMC-2:380–388, July 1972. doi: 10.1109/TSMC.1972.4309133.
- Eshaan Nichani, Jason D. Lee, and Alberto Bietti. Understanding factual recall in Transformers via associative memories, 2024. URL <https://arxiv.org/abs/2412.06538>.
- Hubert Ramsauer, Bernhard Schäfl, Johannes Lehner, Philipp Seidl, Michael Widrich, Thomas Adler, Lukas Gruber, Markus Holzleitner, Milena Pavlović, Geir Kjetil Sandve, Victor Greiff, David Kreil, Michael Kopp, Günter Klambauer, Johannes Brandstetter, and Sepp Hochreiter. Hopfield networks is all you need, 2021. URL <https://arxiv.org/abs/2008.02217>.
- R. Tyrrell Rockafellar and Stanislav Uryasev. Optimization of conditional value-at-risk. *The Journal of Risk*, 2(3):21–41, 2000. doi: 10.21314/JOR.2000.038.
- R. Tyrrell Rockafellar and Stanislav Uryasev. Conditional value-at-risk for general loss distributions. *Journal of Banking & Finance*, 26(7):1443–1471, 2002. doi: 10.1016/S0378-4266(02)00271-6.
- Pragya Sur and Emmanuel J. Candès. A modern maximum-likelihood theory for high-dimensional logistic regression. *Proceedings of the National Academy of Sciences*, 116(29):14516–14525, 2019. doi: 10.1073/pnas.1810420116.
- Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N. Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. In *Proceedings of the 31st International Conference on Neural Information Processing Systems*, NIPS’17, page 6000–6010, 2017.

Nuri Mert Vural, Alberto Bietti, Mahdi Soltanolkotabi, and Denny Wu. Learning to recall with Transformers beyond orthogonal embeddings, 2026. URL <https://arxiv.org/abs/2603.15923>.

Jason Weston, Sumit Chopra, and Antoine Bordes. Memory networks, 2015. URL <https://arxiv.org/abs/1410.3916>.

D. J. Willshaw, O. P. Buneman, and H. C. Longuet-Higgins. Non-holographic associative memory. *Nature*, 222:960–962, 1969. doi: 10.1038/222960a0.

A Correlation matrix memory

Throughout this appendix, $\chi(k)$ denotes the chi distribution with k degrees of freedom.

A.1 A single-coordinate law

Lemma 3 (Law of a single coordinate). *Let $[\theta \ \xi^\top]^\top := S e_1 \in \mathbb{R} \times \mathbb{R}^{n-1}$. Let $a \sim \mathcal{N}(0, 1)$, $g \sim \mathcal{N}(0, I_{n-1})$, and*

$$\rho_1, \rho_2 \sim \frac{\chi(d)}{\sqrt{d}}, \quad \rho_3 \sim \frac{\chi(n-1)}{\sqrt{n}}, \quad \rho_4 \sim \frac{\chi(d-1)}{\sqrt{d}}, \quad \rho_5 \sim \frac{\chi(n-2)}{\sqrt{n}},$$

all mutually independent. Then

$$\begin{bmatrix} \theta \\ \xi \end{bmatrix} \stackrel{(d)}{=} \rho_1 \begin{bmatrix} \rho_2 s \\ \frac{r}{\|g\|/\sqrt{n}} g \end{bmatrix},$$

where

$$s = \rho_1 \rho_2 + \sqrt{\frac{n}{d^2}} \rho_3 a, \quad r = \left[\frac{(as/\sqrt{n} + \rho_3 \rho_4^2)^2}{d} + \frac{s^2 \rho_5^2}{d} + \frac{n}{d^2} \rho_3^2 \rho_4^2 \rho_5^2 \right]^{1/2}. \quad (84)$$

Proof. We derive the distributional representation (84) of the score vector for a single coordinate. It is easy to see that, for any deterministic orthogonal matrix $O \in O(n-1)$,

$$\begin{bmatrix} \theta \\ \xi \end{bmatrix} \stackrel{(d)}{=} \begin{bmatrix} \theta \\ O \xi \end{bmatrix}. \quad (85)$$

To see this, writing

$$Q := \begin{bmatrix} 1 & 0 \\ 0 & O \end{bmatrix} \in O(n),$$

we have $(UQ^\top, VQ^\top) \stackrel{(d)}{=} (U, V)$. Defining

$$w := U^\top U V^\top V e_1, \quad w' := (UQ^\top)^\top (UQ^\top) (VQ^\top)^\top (VQ^\top) e_1,$$

it holds that $w' = Qw$ since $Q^\top e_1 = e_1$. Since $w' \stackrel{(d)}{=} w$, we have $w \stackrel{(d)}{=} Qw$ and so (85) holds. From rotational invariance, it follows that

$$\begin{bmatrix} \theta \\ \xi \end{bmatrix} \stackrel{(d)}{=} \begin{bmatrix} \theta \\ \|\xi\| \frac{g}{\|g\|} \end{bmatrix}, \quad (86)$$

where $g \sim \mathcal{N}(0, I_{n-1})$ is a Gaussian vector independent of θ and $\|\xi\|$.

Let $v_1 \in \mathbb{R}^d$ denote the first column of V . It is straightforward to check that

$$V^\top V e_1 \stackrel{(d)}{=} \|v_1\| \begin{bmatrix} \|v_1\| \\ z \end{bmatrix},$$

where $z \sim \mathcal{N}(0, I_{n-1}/d)$ is independent of v_1 . We may also write

$$U = \begin{bmatrix} u_1 & H_{u_1} M H_z^\top \end{bmatrix} = H_{u_1} \begin{bmatrix} \|u_1\| e_1 & M H_z^\top \end{bmatrix},$$

where $H_{u_1} \in \mathbb{R}^{d \times d}$ and $H_z \in \mathbb{R}^{(n-1) \times (n-1)}$ denote orthogonal matrices whose first columns are aligned with u_1 and z respectively, and

$$M = \begin{bmatrix} \tilde{a} & q^\top \\ p & G \end{bmatrix} \in \mathbb{R}^{d \times (n-1)}, \quad \tilde{a} \in \mathbb{R}, \quad q \in \mathbb{R}^{n-2}, \quad p \in \mathbb{R}^{d-1}, \quad G \in \mathbb{R}^{(d-1) \times (n-2)}$$

is a Gaussian matrix with i.i.d. $\mathcal{N}(0, 1/d)$ entries. Substituting the above representations in

$$U^\top U V^\top V e_1 = \begin{bmatrix} \theta \\ \xi \end{bmatrix}$$

yields

$$\theta = \|v_1\| \|u_1\| s, \quad \text{where} \quad s := \|u_1\| \|v_1\| + \tilde{a} \|z\|.$$

Moreover,

$$\xi = \|v_1\| H_z \begin{bmatrix} \tilde{a} s + \|z\| \|p\|^2 \\ s q + \|z\| G^\top p \end{bmatrix} \stackrel{(d)}{=} \|v_1\| H_z \begin{bmatrix} \tilde{a} s + \|z\| \|p\|^2 \\ \sqrt{s^2 + \|z\|^2 \|p\|^2} \tilde{z} \end{bmatrix}, \quad (87)$$

where $\tilde{z} \sim \mathcal{N}(0, I_{n-2}/d)$ is independent of (s, z, p) . To reach the claim of the lemma, set

$$\rho_1 = \|v_1\|, \quad \rho_2 = \|u_1\|, \quad \rho_3 = \sqrt{\frac{d}{n}} \|z\|, \quad \rho_4 = \|p\|, \quad \rho_5 = \sqrt{\frac{d}{n}} \|\tilde{z}\|.$$

Moreover, the standard Gaussian variable a in the statement of the lemma is $\tilde{a} \sqrt{d}$. The claim of the lemma then follows readily by applying (86) with (87). $\square$

A.2 Joint distribution of two columns

To characterize the joint distribution of two columns, we will make use of the following lemma.

Lemma 4. *For $k \leq d$, let $\xi_1, \dots, \xi_k \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_d/d)$. Define the orthogonal matrix $H \in O(d)$ recursively by*

$$h_1 := \frac{\xi_1}{\|\xi_1\|},$$

$$h_j := \frac{r_j}{\|r_j\|}, \quad r_j := (I - h_1 h_1^\top - \dots - h_{j-1} h_{j-1}^\top) \xi_j$$

for $j \leq k$ and complete the remaining $h_{k+1}, \dots, h_d$ columns to an orthonormal basis of $\mathbb{R}^d$. Then, the entries of

$$R = H^\top \begin{bmatrix} \xi_1 & \dots & \xi_k \end{bmatrix}$$

are mutually independent and have marginal distributions

$$R_{ij} \sim \begin{cases} \mathcal{N}(0, 1/d), & i < j, \\ \frac{1}{\sqrt{d}}\chi(d - j + 1), & i = j, \\ 0, & i > j. \end{cases} \quad (88)$$

Moreover,

$$\xi_i^\top \xi_i = \sum_{t=1}^i R_{ti}^2 \sim \frac{1}{d}\chi^2(d) \quad (89)$$

and for $i < j$

$$\xi_i^\top \xi_j = \sum_{t=1}^i R_{ti} R_{tj} \stackrel{(d)}{=} \frac{1}{d} \sum_{t=1}^{i-1} \zeta_t \varepsilon_t + \frac{1}{d} \chi(d - i + 1) Z \quad (90)$$

where $\zeta_t, \varepsilon_t, Z \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$ are independent of $\chi(d - i + 1)$.

Proof. Since $\xi_j \in \text{span}(h_1, \dots, h_j)$ by construction, we have

$$\xi_j = \sum_{i=1}^j (h_i^\top \xi_j) h_i = \sum_{i=1}^j R_{ij} h_i.$$

Now, fix $j \in \{1, \dots, k\}$ and condition on $\mathcal{F}_{j-1} = \sigma(\xi_1, \dots, \xi_{j-1})$, equivalently on $(h_1, \dots, h_{j-1})$. Since $\xi_j \sim \mathcal{N}(0, I_d/d)$ is independent of $\mathcal{F}_{j-1}$, the vector $(h_1^\top \xi_j, \dots, h_{j-1}^\top \xi_j, r_j) \in \mathbb{R}^{j-1+d}$ is jointly Gaussian. In particular, one finds

$$(h_1^\top \xi_j, \dots, h_{j-1}^\top \xi_j, P_{j-1}^\perp \xi_j) \mid \mathcal{F}_{j-1} \sim \mathcal{N}(0, \Sigma) \quad \text{with} \quad \Sigma = \begin{bmatrix} \frac{1}{d} I_{j-1} & 0_{(j-1) \times d} \\ 0_{d \times (j-1)} & \frac{1}{d} P_{j-1}^\perp \end{bmatrix}$$

$$\text{where} \quad P_{j-1}^\perp := I - \sum_{i=1}^{j-1} h_i h_i^\top.$$

Hence, conditional on $\mathcal{F}_{j-1}$, for $i < j$, the scalars

$$R_{ij} = h_i^\top \xi_j \sim \mathcal{N}(0, 1/d)$$

are independent of one another and independent of r_j . Moreover, since $r_j = P_{j-1}^\perp \xi_j$ is an isotropic Gaussian in a $(d - j + 1)$ -dimensional subspace, we have

$$R_{jj} = \frac{r_j^\top \xi_j}{\|r_j\|} = \|r_j\| \sim \frac{1}{\sqrt{d}} \chi(d - j + 1).$$

As the conditional laws above do not depend on $\mathcal{F}_{j-1}$, they also hold unconditionally, and since $\xi_1, \dots, \xi_k$ are independent, the random variables $\{R_{ij} : 1 \leq i \leq j\}$ are independent across $1 \leq j \leq k$. By construction $R_{ij} = 0$ for $i > j$, and thus we arrive at the representation (88). The characterizations (89) and (90) follow from

$$R^\top R = \begin{bmatrix} \xi_1 & \cdots & \xi_k \end{bmatrix}^\top \begin{bmatrix} \xi_1 & \cdots & \xi_k \end{bmatrix}$$

and the distributional representation (88). $\square$

Recalling that the score matrix $S = U^\top UV^\top V$, write $S_{:,1:2}$ for the first two columns:

$$S_{:,1:2} = U^\top UV^\top V[e_1 \ e_2] =: \begin{bmatrix} T \\ \Xi \end{bmatrix}$$

where $T \in \mathbb{R}^{2 \times 2}$ and $\Xi \in \mathbb{R}^{(n-2) \times 2}$. For an orthogonal matrix $O \in \mathbb{R}^{(n-2) \times (n-2)}$, it is not hard to see that

$$\begin{bmatrix} T \\ \Xi \end{bmatrix} \stackrel{(d)}{=} \begin{bmatrix} T \\ O\Xi \end{bmatrix}.$$

Let $V = [V_{12} \ \tilde{V}']$ where $V_{12} = [v_1 \ v_2] \in \mathbb{R}^{d \times 2}$ and $\tilde{V}' = [v_3 \dots v_n] \in \mathbb{R}^{d \times (n-2)}$. By rotational invariance, $\tilde{V}' \stackrel{(d)}{=} H_{V_{12}} \tilde{V}$ where $\tilde{V}$ is an independent copy of $\tilde{V}'$ and $H_{V_{12}}$ is an orthogonal matrix constructed via Lemma 4 using $(\xi_1, \xi_2) = (v_1, v_2)$. Letting $Z = [z_1 \ z_2] \in \mathbb{R}^{(n-2) \times 2}$ denote the first two columns of $\tilde{V}^\top$ where $z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_{n-2}/d)$, we have

$$V^\top V[e_1 \ e_2] = \begin{bmatrix} \|v_1\|^2 & v_1^\top v_2 \\ v_1^\top v_2 & \|v_2\|^2 \\ \tilde{V}^\top H_{V_{12}}^\top v_1 & \tilde{V}^\top H_{V_{12}}^\top v_2 \end{bmatrix} \stackrel{(d)}{=} \begin{bmatrix} A_1 \\ Z A_2 \end{bmatrix}$$

where

$$A_1 = \begin{bmatrix} c_1^2 & c_1 c_2 \\ c_1 c_2 & c_2^2 + c_3^2 \end{bmatrix}, \quad A_2 = \begin{bmatrix} c_1 & c_2 \\ 0 & c_3 \end{bmatrix}$$

and c_1, c_2, c_3 are mutually independent, independent of z_1, z_2 , and distributed as

$$c_1 \sim \frac{1}{\sqrt{d}} \chi(d), \quad c_2 \sim \mathcal{N}\left(0, \frac{1}{d}\right), \quad c_3 \sim \frac{1}{\sqrt{d}} \chi(d-1).$$

Continuing, we have

$$UV^\top V[e_1 \ e_2] = U_{12} A_1 + \tilde{U}' Z A_2$$

where $U_{12} = [u_1 \ u_2]$ and $\tilde{U}' = [u_3 \dots u_n] \in \mathbb{R}^{d \times (n-2)}$. In the same way as before, we write $\tilde{U}' \stackrel{(d)}{=} \tilde{U} H_Z^\top$ where $\tilde{U}$ is an independent copy of $\tilde{U}'$ and H_Z is an orthogonal matrix constructed via Lemma 4 using $(\xi_1, \xi_2) = (z_1, z_2)$. Letting $Y = [y_1 \ y_2] \in \mathbb{R}^{d \times 2}$ denote the first two columns of $\tilde{U}$ where $y_1, y_2 \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_d/d)$, we have

$$UV^\top V[e_1 \ e_2] = U_{12} A_1 + Y A_3 A_2 \quad \text{where} \quad A_3 = \begin{bmatrix} c_4 & c_5 \\ 0 & c_6 \end{bmatrix}.$$

and c_4, c_5, c_6 are mutually independent and independent of y_1, y_2 and U_{12} , and have distributions:

$$c_4 \sim \frac{1}{\sqrt{d}} \chi(n-2), \quad c_5 \sim \mathcal{N}\left(0, \frac{1}{d}\right), \quad c_6 \sim \frac{1}{\sqrt{d}} \chi(n-3).$$

We are approaching the quantity of interest:

$$\begin{aligned} S_{:,1:2} &= U^\top U V^\top V [e_1 \ e_2] = \begin{bmatrix} U_{12}^\top U_{12} A_1 + U_{12}^\top Y A_3 A_2 \\ H_Z \left(\tilde{U}^\top U_{12} A_1 + \tilde{U}^\top Y A_3 A_2 \right) \end{bmatrix} \\ &= \begin{bmatrix} U_{12}^\top U_{12} A_1 + U_{12}^\top Y A_3 A_2 \\ H_Z \left(\begin{bmatrix} Y^\top U_{12} \\ \bar{U}'^\top U_{12} \end{bmatrix} A_1 + \begin{bmatrix} Y^\top Y \\ \bar{U}'^\top Y \end{bmatrix} A_3 A_2 \right) \end{bmatrix} \end{aligned}$$

where $\tilde{U} = [Y \ \bar{U}']$ with $\bar{U}' \in \mathbb{R}^{d \times (n-4)}$. We seek to determine the joint distribution of

$$(Y^\top U_{12}, \bar{U}'^\top U_{12}, \bar{U}'^\top Y).$$

For this purpose, we write $\bar{U}' \stackrel{(d)}{=} H \bar{U}$ where $\bar{U}$ is an independent copy of $\bar{U}'$ and H is an orthogonal matrix constructed via Lemma 4 using $(\xi_1, \xi_2, \xi_3, \xi_4) = (u_1, u_2, y_1, y_2)$. Let $\Gamma = [\gamma_1 \dots \gamma_4] \in \mathbb{R}^{(n-4) \times 4}$ denote the first four columns of $\bar{U}^\top$ such that $\gamma_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, I_{n-4}/d)$. Observe that

$$\bar{U}'^\top U_{12} = \bar{U}^\top H^\top U_{12} = \Gamma_{:,1:2} A_4 \quad \text{where} \quad A_4 = \begin{bmatrix} c_7 & c_8 \\ 0 & c_9 \end{bmatrix}$$

and

$$\bar{U}'^\top Y = \bar{U}^\top H^\top Y = \Gamma A_5 \quad \text{where} \quad A_5 = \begin{bmatrix} c_{10} & c_{11} \\ c_{12} & c_{13} \\ c_{14} & c_{15} \\ 0 & c_{16} \end{bmatrix}$$

and $c_7, \dots, c_{16}$ are mutually independent and independent of $\gamma_1, \dots, \gamma_4$, and have distributions:

$$\begin{aligned} c_7 &\sim \frac{1}{\sqrt{d}} \chi(d), \quad c_9 \sim \frac{1}{\sqrt{d}} \chi(d-1), \quad c_{14} \sim \frac{1}{\sqrt{d}} \chi(d-2), \quad c_{16} \sim \frac{1}{\sqrt{d}} \chi(d-3), \\ c_8, c_{10}, c_{11}, c_{12}, c_{13}, c_{15} &\stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1/d). \end{aligned}$$

Appealing to Lemma 4, we can write

$$A_6 = U_{12}^\top U_{12} \stackrel{(d)}{=} \begin{bmatrix} c_7^2 & c_7 c_8 \\ c_7 c_8 & c_8^2 + c_9^2 \end{bmatrix}$$

and

$$A_7 = Y^\top Y \stackrel{(d)}{=} \begin{bmatrix} c_{10}^2 + c_{12}^2 + c_{14}^2 & c_{10} c_{11} + c_{12} c_{13} + c_{14} c_{15} \\ c_{10} c_{11} + c_{12} c_{13} + c_{14} c_{15} & c_{11}^2 + c_{13}^2 + c_{15}^2 + c_{16}^2 \end{bmatrix}.$$

Finally, we can write

$$\begin{aligned} A_8 &= Y^\top U_{12} = (H^\top Y)^\top (H^\top U_{12}) = \begin{bmatrix} A_5 \\ 0_{(d-4) \times 2} \end{bmatrix}^\top \begin{bmatrix} A_4 \\ 0_{(d-2) \times 2} \end{bmatrix} \\ &= \begin{bmatrix} c_{10} & c_{12} \\ c_{11} & c_{13} \end{bmatrix} \begin{bmatrix} c_7 & c_8 \\ 0 & c_9 \end{bmatrix} = \begin{bmatrix} c_7 c_{10} & c_8 c_{10} + c_9 c_{12} \\ c_7 c_{11} & c_8 c_{11} + c_9 c_{13} \end{bmatrix}. \end{aligned}$$

Putting things together, we have

$$U^\top UV^\top V[e_1 \ e_2] = \begin{bmatrix} A_6 A_1 + A_8^\top A_3 A_2 \\ H_Z \left(\begin{bmatrix} A_8 \\ \Gamma_{:,1:2} A_4 \end{bmatrix} A_1 + \begin{bmatrix} A_7 \\ \Gamma A_5 \end{bmatrix} A_3 A_2 \right) \end{bmatrix}.$$

Remark 6. As $n, d \rightarrow \infty$ with $\frac{n}{d^2} \rightarrow \alpha \in (0, \infty)$, for fixed $m \in \mathbb{N}$, one finds

$$S_{1:m,1:2} \xrightarrow{(d)} \begin{bmatrix} I_2 + \sqrt{\alpha} G \\ \sqrt{\alpha} G' \end{bmatrix}$$

where G, G' are independent matrices of i.i.d. $\mathcal{N}(0, 1)$ entries.

A.3 Proof of Proposition 1

Denote

$$\alpha_n := \frac{n}{d^2} \asymp \frac{1}{\log n}.$$

We will make use of the following concentration inequalities. The proofs are standard and will be omitted.

Lemma 5. For $Z \sim \chi^2(k)$ and $\delta \in (0, 1)$,

$$\mathbb{P}(Z \leq (1 - \delta)k) \leq \exp\left(-\frac{\delta^2 k}{4}\right), \quad \mathbb{P}(Z \geq (1 + \delta)k) \leq \exp\left(-\frac{\delta^2 k}{4}\right).$$

Moreover, for $R = \sqrt{Z} \sim \chi(k)$,

$$\mathbb{P}\left(\left|\frac{R}{\sqrt{k}} - 1\right| > \delta\right) \leq 2 \exp\left(-\frac{\delta^2 k}{4}\right).$$

Lemma 6. For $a \sim \mathcal{N}(0, 1)$ and $0 < \beta < 2$,

$$\mathbb{P}\left(a \geq \sqrt{(2 - \beta) \log n}\right) \asymp \frac{n^{-1+\beta/2}}{\sqrt{\log n}},$$

where the hidden constants can be made independent of β . Moreover for $g \sim \mathcal{N}(0, I_{n-1})$,

$$\mathbb{P}\left(\max_{1 \leq i \leq n-1} g_i < \sqrt{2 \log n} - 1\right) = n^{-\omega(1)}.$$

We define the per-coordinate failure event

$$F_i = \left\{ s_{i,i}^n \leq \max_{j \neq i} s_{j,i}^n \right\}$$

for each $1 \leq i \leq n$ —hiding the implicit n -dependence of the sets—and first prove two preliminary results. Without loss of generality, we operate under the regime where d^2 and $n \log n$ are asymptotically comparable, that is $\varrho \in (0, \infty)$.

Lemma 7. For $d^2/n \log n \rightarrow \varrho \in (0, \infty)$, it holds that

$$\lim_{n \rightarrow \infty} n\mathbb{P}(F_1) = \begin{cases} 0 & \text{if } \varrho > 8, \\ \infty & \text{if } \varrho < 8. \end{cases}$$

Proof. For simplicity, we adopt the notation $\theta = s_{1,1}^n$ and $\xi_j = s_{j,1}^n$ from Lemma 3 where we recall the distributional representation:

$$\begin{bmatrix} \theta \\ \xi \end{bmatrix} \stackrel{(d)}{=} \rho_1 \begin{bmatrix} \rho_2 s \\ \frac{r}{\|g\|/\sqrt{n}} g \end{bmatrix},$$

where

$$s = \rho_1 \rho_2 + \sqrt{\frac{n}{d^2}} \rho_3 a, \quad r^2 = \frac{1}{d} (as/\sqrt{n} + \rho_3 \rho_4^2)^2 + \frac{1}{d} s^2 \rho_5^2 + \frac{n}{d^2} \rho_3^2 \rho_4^2 \rho_5^2$$

and $a, g, \rho_1, \dots, \rho_5$ are independent random variables such that

$$a \sim \mathcal{N}(0, 1), \quad g \sim \mathcal{N}(0, I_{n-1}),$$

and

$$\rho_1, \rho_2 \sim \frac{\chi(d)}{\sqrt{d}}, \quad \rho_3 \sim \frac{\chi(n-1)}{\sqrt{n}}, \quad \rho_4 \sim \frac{\chi(d-1)}{\sqrt{d}}, \quad \rho_5 \sim \frac{\chi(n-2)}{\sqrt{n}}.$$

Fix $\delta \in (0, 1)$ and define the concentration event

$$\mathcal{E}_\delta := \{\rho_i \in [1 - \delta, 1 + \delta] \ \forall i = 1, \dots, 5\} \cap \{\|g\|^2/n \in [1 - \delta, 1 + \delta]\}.$$

Applying Lemma 5, we have

$$\mathbb{P}(\mathcal{E}_\delta^c) \leq C \exp\left(-\frac{\delta^2 \min\{d-1, n-2\}}{4}\right)$$

for some constant $C > 0$. In particular, under the regime $d^2 \asymp n \log n$, we have

$$n\mathbb{P}(\mathcal{E}_\delta^c) \xrightarrow[n \rightarrow \infty]{} 0. \tag{91}$$

Define the truncation event

$$\mathcal{A}_n := \{|a| \leq 2\sqrt{\log n}\} \cap \left\{ \max_{1 \leq i \leq n-1} g_i \geq \sqrt{2 \log n} - 1 \right\}.$$

Since $a \sim \mathcal{N}(0, 1)$, a standard Gaussian tail bound combined with Lemma 6 yields that $\mathbb{P}(\mathcal{A}_n^c) \leq 2e^{-2 \log n} + n^{-\omega(1)} = O(n^{-2})$ and hence

$$n\mathbb{P}(\mathcal{A}_n^c) \xrightarrow[n \rightarrow \infty]{} 0. \tag{92}$$

Now on $\mathcal{E}_\delta \cap \mathcal{A}_n$, we have

$$|s| \leq (1 + \delta)^2 + 2(1 + \delta)\sqrt{\alpha_n \log n} = O(1),$$

therefore $as/\sqrt{n} = o(1)$ and

$$\begin{aligned} r^2 - \alpha_n &= \frac{1}{d} (as/\sqrt{n} + \rho_3 \rho_4^2)^2 + \frac{1}{d} s^2 \rho_5^2 + \alpha_n (\rho_3^2 \rho_4^2 \rho_5^2 - 1) \\ &= O\left(\frac{1}{d}\right) + O(\delta) \alpha_n = O(\delta) \alpha_n, \end{aligned}$$

which further implies $r/\sqrt{\alpha_n} = 1 + O(\delta)$. Moreover, the event F_1 is equivalent to

$$\theta = \rho_1 \rho_2 s \leq \max_{1 \leq j \leq n-1} \xi_j = \rho_1 r \frac{\sqrt{n}}{\|g\|} \max_{1 \leq j \leq n-1} g_j$$

by using the distributional representation for θ, ξ . Substituting the representation for s , this gives

$$(\rho_1 \rho_2)^2 + (\rho_1 \rho_2) \rho_3 \sqrt{\alpha_n} a \leq \rho_1 r \frac{\sqrt{n}}{\|g\|} \max_{1 \leq j \leq n-1} g_j.$$

Dividing both sides by $(\rho_1 \rho_2) \rho_3 \sqrt{\alpha_n}$, we get

$$F_1 = \left\{ a \leq \frac{r \sqrt{n}}{\rho_2 \rho_3 \|g\| \sqrt{\alpha_n}} \max_{1 \leq j \leq n-1} g_j - \frac{\rho_1 \rho_2}{\rho_3 \sqrt{\alpha_n}} \right\}. \quad (93)$$

We now consider the two regimes separately.

Case I: $\varrho > 8$. On $\mathcal{E}_\delta \cap \mathcal{A}_n$, (93) implies

$$a \leq A_\delta \max_{1 \leq j \leq n-1} g_j - \frac{B_\delta}{\sqrt{\alpha_n}}$$

for δ -dependent constants $A_\delta, B_\delta = 1 + O(\delta)$. Since a is independent of g , each variable $A_\delta g_j - a$ is Gaussian with marginal distribution

$$A_\delta g_j - a \sim \mathcal{N}(0, A_\delta^2 + 1),$$

hence a union bound and a Gaussian tail estimate give

$$\mathbb{P}(F_1 \cap \mathcal{E}_\delta \cap \mathcal{A}_n) \leq \sum_{j=1}^{n-1} \mathbb{P}\left(A_\delta g_j - a \geq \frac{B_\delta}{\sqrt{\alpha_n}}\right) \leq n \exp\left(-\frac{B_\delta^2}{2(A_\delta^2 + 1)\alpha_n}\right).$$

Since $\alpha_n \log n \rightarrow 1/\varrho$ as $n \rightarrow \infty$ and $\varrho > 8$, it holds that

$$\alpha_n \log n \leq \frac{1}{2\varrho} + \frac{1}{16}$$

for all sufficiently large n . Moreover since both $A_\delta, B_\delta = 1 + O(\delta)$, we may choose $\delta > 0$ sufficiently small (depending only on ϱ) so that

$$\frac{B_\delta^2}{2(A_\delta^2 + 1)} \geq \frac{1}{2\varrho} + \frac{3}{16}.$$

It follows that

$$\frac{B_\delta^2}{2(A_\delta^2 + 1)\alpha_n} \geq \frac{\frac{1}{2\varrho} + \frac{3}{16}}{\frac{1}{2\varrho} + \frac{1}{16}} \log n = \left(2 + \frac{\varrho - 8}{\varrho + 8}\right) \log n,$$

hence

$$n\mathbb{P}(F_1 \cap \mathcal{E}_\delta \cap \mathcal{A}_n) \leq n^2 \exp\left(-\frac{B_\delta^2}{2(A_\delta^2 + 1)\alpha_n}\right) \xrightarrow{n \rightarrow \infty} 0.$$

Furthermore,

$$n\mathbb{P}(F_1) \leq n\mathbb{P}(\mathcal{E}_\delta^c) + n\mathbb{P}(\mathcal{A}_n^c) + n\mathbb{P}(F_1 \cap \mathcal{E}_\delta \cap \mathcal{A}_n)$$

and each term on the right-hand side tends to zero by (91), (92), and the previous display. Therefore,

$$\lim_{n \rightarrow \infty} n\mathbb{P}(F_1) = 0.$$

Case II: $\varrho < 8$. Similar to Case I, on $\mathcal{E}_\delta \cap \mathcal{A}_n$, (93) is implied by

$$a \leq \left(\sqrt{2 \log n} - 1\right) A'_\delta - \frac{B'_\delta}{\sqrt{\alpha_n}} \leq A'_\delta \max_{1 \leq j \leq n-1} g_j - \frac{B'_\delta}{\sqrt{\alpha_n}}$$

for δ -dependent constants $A'_\delta, B'_\delta = 1 + O(\delta)$ (note that the second inequality always holds due to the definition of $\mathcal{A}_n$). Define

$$C_{\delta,n} := \frac{1}{\sqrt{\log n}} \left(\left(\sqrt{2 \log n} - 1\right) A'_\delta - \frac{B'_\delta}{\sqrt{\alpha_n}} \right).$$

Since $\sqrt{2} - \sqrt{\varrho} > -\sqrt{2}$, we may choose $\delta > 0$ sufficiently small so that

$$\lim_{n \rightarrow \infty} C_{\delta,n} = \sqrt{2}A'_\delta - \sqrt{\varrho}B'_\delta > -\sqrt{2},$$

thus there exists $\beta > 0$ such that $C_{\delta,n} > -\sqrt{2 - \beta}$ for all sufficiently large n . It follows that

$$\mathcal{E}_\delta \cap \mathcal{A}_n \cap \left\{a \leq -\sqrt{(2 - \beta) \log n}\right\} \subseteq F_1$$

and so

$$n\mathbb{P}(F_1) \geq n\mathbb{P}\left(a \leq -\sqrt{(2 - \beta) \log n}\right) - n\mathbb{P}(\mathcal{E}_\delta^c) - n\mathbb{P}(\mathcal{A}_n^c).$$

The first term in the right-hand side diverges from Lemma 6, while the remaining terms again tend to zero, therefore

$$\lim_{n \rightarrow \infty} n\mathbb{P}(F_1) = \infty.$$

This completes the proof. □

We now show that the events $F_1, \dots, F_n$ are only weakly dependent.

Lemma 8. *In the regime $d^2 \asymp n \log n$, if*

$$n\mathbb{P}(F_1) \xrightarrow[n \rightarrow \infty]{} \infty,$$

then the events F_1 and F_2 are asymptotically independent in the sense that

$$\mathbb{P}(F_1 \cap F_2) = \mathbb{P}(F_1)\mathbb{P}(F_2)(1 + \omega_n)$$

where $\lim_{n \rightarrow \infty} \omega_n = 0$.

Proof. Adopting the notation of Appendix A.2, we have

$$F_1 = \left\{ T_{11} \leq \max \left(T_{21}, \max_{1 \leq j \leq n-2} \Xi_{j1} \right) \right\}, \quad F_2 = \left\{ T_{22} \leq \max \left(T_{12}, \max_{1 \leq j \leq n-2} \Xi_{j2} \right) \right\}.$$

We first write out the relevant entries of T . From the matrix multiplication

$$T = A_6 A_1 + A_8^\top A_3 A_2,$$

one obtains

$$T_{11} = c_1^2 c_7^2 + c_1 c_4 c_7 c_{10} + c_1 c_2 c_7 c_8, \quad (94)$$

$$T_{12} = c_1 c_2 c_7^2 + c_2 c_4 c_7 c_{10} + c_3 c_5 c_7 c_{10} + c_3 c_6 c_7 c_{11} + (c_2^2 + c_3^2) c_7 c_8, \quad (95)$$

$$T_{21} = c_1^2 c_7 c_8 + c_1 c_4 (c_{10} c_8 + c_{12} c_9) + c_1 c_2 (c_8^2 + c_9^2), \quad (96)$$

$$T_{22} = (c_2^2 + c_3^2)(c_8^2 + c_9^2) + c_1 c_2 c_7 c_8 + c_2 c_4 (c_{10} c_8 + c_{12} c_9) \\ + c_3 c_5 (c_{10} c_8 + c_{12} c_9) + c_3 c_6 (c_{11} c_8 + c_{13} c_9). \quad (97)$$

Next, we isolate the lower block. In the notation of Appendix A.2, the lower $(n-2)$ block is

$$\Xi = H_Z Y$$

with

$$Y = \begin{bmatrix} A_8 \\ \Gamma_{:,1:2} A_4 \end{bmatrix} A_1 + \begin{bmatrix} A_7 \\ \Gamma A_5 \end{bmatrix} A_3 A_2.$$

Writing the last $n-4$ rows of Y as ΓB , direct multiplication gives

$$B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \\ 0 & b_{42} \end{bmatrix},$$

where

$$b_{11} = c_1^2 c_7 + c_1 c_4 c_{10} + c_1 c_2 c_8, \quad (98)$$

$$b_{12} = c_1 c_2 c_7 + c_2 c_4 c_{10} + c_3 (c_{10} c_5 + c_{11} c_6) + c_8 (c_2^2 + c_3^2),$$

$$b_{21} = c_1 c_4 c_{12} + c_1 c_2 c_9,$$

$$b_{22} = c_2 c_4 c_{12} + c_3 (c_{12} c_5 + c_{13} c_6) + c_9 (c_2^2 + c_3^2),$$

$$b_{31} = c_1 c_4 c_{14},$$

$$b_{32} = c_2 c_4 c_{14} + c_3 (c_{14} c_5 + c_{15} c_6),$$

$$b_{42} = c_3 c_6 c_{16}. \quad (99)$$

Since H_Z is orthogonal, one has $\Xi^\top \Xi = Y^\top Y$. Moreover, the entries of the first two rows of Y are $O_{\mathbb{P}}(1)$, hence their contribution to $n^{-1}\Xi^\top \Xi$ is $O_{\mathbb{P}}(n^{-1})$, which is negligible. Thus the main contribution comes from the ΓB block:

$$\frac{1}{n}\Xi^\top \Xi = \frac{1}{n}B^\top \Gamma^\top \Gamma B + O_{\mathbb{P}}\left(\frac{1}{n}\right). \quad (100)$$

Define

$$\delta_n := (\log n)^{-3}, \quad L_n := \sqrt{8 \log n}, \quad \eta_n := \delta_n + \frac{\log n}{\sqrt{d}} + n^{-1/3}.$$

and note that since $d^2 \asymp n \log n$, we have

$$\eta_n = o\left(\frac{\alpha_n}{\sqrt{\log n}}\right), \quad \frac{\eta_n}{\sqrt{\alpha_n}} = o\left(\frac{1}{\sqrt{\log n}}\right).$$

Let $\mathcal{E}_n := \mathcal{E}_n^1 \cap \mathcal{E}_n^2 \cap \mathcal{E}_n^3$ be the intersection of the following three events:

$$\begin{aligned} \mathcal{E}_n^1 &= \left\{ \left| \sqrt{\frac{d}{n}} c_4 - 1 \right| + \left| \sqrt{\frac{d}{n}} c_6 - 1 \right| + \sum_{i \in \{1, 3, 7, 9, 14, 16\}} |c_i - 1| \leq \delta_n \right\}, \\ \mathcal{E}_n^2 &= \left\{ |\sqrt{d} c_j| \leq L_n \quad \text{for } j \in \{2, 5, 8, 11, 12, 15\} \right\}, \\ \mathcal{E}_n^3 &= \left\{ \left\| \Gamma^\top \Gamma - \frac{n-4}{d} I_4 \right\|_{\text{op}} \leq \delta_n \frac{n}{d} \right\}. \end{aligned}$$

By Lemma 5, Gaussian tail bounds, and the fact that the dimension of $\Gamma^\top \Gamma$ is fixed,

$$\mathbb{P}(\mathcal{E}_n^c) = O(n^{-4}) + \exp\left(-\Omega\left(\frac{d}{\log^6 n}\right)\right) = O(n^{-4}). \quad (101)$$

Now define

$$a_1 := \sqrt{d} c_{10}, \quad a_2 := \sqrt{d} c_{13},$$

so that $a_1, a_2 \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$, and consider the event

$$\mathcal{A}_n := \{|a_1| \leq L_n, |a_2| \leq L_n\}.$$

Then

$$\mathbb{P}(\mathcal{A}_n^c) = O(n^{-4}) \quad (102)$$

by standard Gaussian concentration. On $\mathcal{E}_n \cap \mathcal{A}_n$, (94)–(97) imply

$$T_{11} = 1 + \sqrt{\alpha_n} a_1 + r_{1,n}, \quad (103)$$

$$T_{22} = 1 + \sqrt{\alpha_n} a_2 + r_{2,n}, \quad (104)$$

with

$$|r_{1,n}| + |r_{2,n}| \leq C \eta_n \quad (105)$$

for some universal constant $C > 0$. Indeed, the leading random term in T_{11} is $c_1 c_4 c_7 c_{10} = \sqrt{\alpha_n} a_1 (1 + o(1))$, while every other term in (94) is absorbed in $O(\eta_n)$; similarly, the leading random term in T_{22} is $c_3 c_6 c_9 c_{13} = \sqrt{\alpha_n} a_2 (1 + o(1))$, and every remaining term in (97) is again $O(\eta_n)$.

Next, from (98)–(99), on the event $\mathcal{E}_n \cap \mathcal{A}_n$,

$$b_{31} = \sqrt{\frac{n}{d}}(1 + O(\delta_n)), \quad b_{42} = \sqrt{\frac{n}{d}}(1 + O(\delta_n)),$$

while

$$b_{11}, b_{22} = O(1), \quad b_{12}, b_{21}, b_{32} = O\left(\sqrt{\alpha_n \log n}\right) = O(1).$$

Therefore,

$$\frac{1}{d} B^\top B = \begin{bmatrix} \alpha_n & 0 \\ 0 & \alpha_n \end{bmatrix} + E_n^{(0)}, \quad \|E_n^{(0)}\|_{\text{op}} \leq C\eta_n.$$

Combining this with (100) and the concentration of $\Gamma^\top \Gamma$ on $\mathcal{E}_n$, we obtain

$$\frac{1}{n} \Xi^\top \Xi = \begin{bmatrix} \sigma_{1,n}^2 & \rho_n \\ \rho_n & \sigma_{2,n}^2 \end{bmatrix}, \quad |\sigma_{1,n}^2 - \alpha_n| + |\sigma_{2,n}^2 - \alpha_n| + |\rho_n| \leq C\eta_n.$$

In particular,

$$\sigma_{1,n} = \sqrt{\alpha_n}(1 + o(\log^{-1} n)), \quad \sigma_{2,n} = \sqrt{\alpha_n}(1 + o(\log^{-1} n)), \quad |\rho_n| \leq C\eta_n. \quad (106)$$

We now employ the left-orthogonal invariance of Ξ . Conditional on $\Xi^\top \Xi$, there exist independent Gaussian vectors $g, h \sim \mathcal{N}(0, I_{n-2})$ such that

$$\Xi_{:,1} \stackrel{(d)}{=} \sigma_{1,n} \frac{\sqrt{n}}{\|g\|} g, \quad (107)$$

$$\Xi_{:,2} \stackrel{(d)}{=} \frac{\rho_n}{\sigma_{1,n}} \frac{\sqrt{n}}{\|g\|} g + \tau_n \frac{\sqrt{n}}{\|\tilde{h}\|} \tilde{h}, \quad (108)$$

where

$$\tilde{h} := h - \frac{\langle g, h \rangle}{\|g\|^2} g, \quad \tau_n^2 := \sigma_{2,n}^2 - \frac{\rho_n^2}{\sigma_{1,n}^2}.$$

By (106),

$$\tau_n = \sqrt{\alpha_n}(1 + o(\log^{-1} n)). \quad (109)$$

Let $\mathcal{G}_n := \mathcal{G}_n^1 \cap \mathcal{G}_n^2 \cap \mathcal{G}_n^3$ where

$$\begin{aligned} \mathcal{G}_n^1 &= \left\{ \left| \frac{\|g\|}{\sqrt{n}} - 1 \right| + \left| \frac{\|\tilde{h}\|}{\sqrt{n}} - 1 \right| \leq \delta_n \right\}, \\ \mathcal{G}_n^2 &= \left\{ \frac{|\langle g, h \rangle|}{n} \leq n^{-1/3} \right\}, \\ \mathcal{G}_n^3 &= \left\{ \left(\max_{j \leq n-2} |g_j| \right) \vee \left(\max_{j \leq n-2} |h_j| \right) \leq L_n \right\}. \end{aligned}$$

By standard Gaussian concentration, we have

$$\mathbb{P}(\mathcal{G}_n^c) = O(n^{-4}). \quad (110)$$

On $\mathcal{E}_n \cap \mathcal{A}_n \cap \mathcal{G}_n$, using (107)–(108), (106), and (109), we have

$$\max_{j \leq n-2} |\Xi_{j1} - \sqrt{\alpha_n} g_j| \leq \frac{C}{\log^2 n}, \quad (111)$$

$$\max_{j \leq n-2} |\Xi_{j2} - \sqrt{\alpha_n} h_j| \leq \frac{C}{\log^2 n}. \quad (112)$$

Indeed, the projection correction in $\tilde{h}$ contributes at most

$$\tau_n \frac{|\langle g, h \rangle|}{\|g\|^2} \max_{j \leq n-2} |g_j| \lesssim \sqrt{\alpha_n} n^{-1/3} L_n = O(n^{-1/3}),$$

and the shared term $\frac{\rho_n}{\sigma_{1,n}} \frac{\sqrt{n}}{\|g\|} g$ contributes at most

$$\frac{|\rho_n|}{\sigma_{1,n}} L_n \lesssim \frac{\eta_n}{\sqrt{\alpha_n}} L_n = O(\log^{-2} n).$$

We now define

$$w_n := \frac{C}{\log^2 n \sqrt{\alpha_n}} = o\left(\frac{1}{\sqrt{\log n}}\right),$$

$$u_{1,n} := \alpha_n^{-1/2} + a_1, \quad u_{2,n} := \alpha_n^{-1/2} + a_2,$$

and

$$\psi_n(t) := 1 - \Phi(t)^{n-2},$$

where Φ is the standard Gaussian CDF. By (103), (104), (105), (111), and (112), we have on $\mathcal{E}_n \cap \mathcal{A}_n \cap \mathcal{G}_n$ the following set of inclusions

$$\left\{ \max_{j \leq n-2} g_j \geq u_{1,n} + w_n \right\} \subseteq \left\{ \max_{j \leq n-2} \Xi_{j1} \geq T_{11} \right\} \subseteq \left\{ \max_{j \leq n-2} g_j \geq u_{1,n} - w_n \right\}, \quad (113)$$

$$\left\{ \max_{j \leq n-2} h_j \geq u_{2,n} + w_n \right\} \subseteq \left\{ \max_{j \leq n-2} \Xi_{j2} \geq T_{22} \right\} \subseteq \left\{ \max_{j \leq n-2} h_j \geq u_{2,n} - w_n \right\}. \quad (114)$$

We now show that the single top competitors T_{21} and T_{12} are negligible. Conditional on $\mathcal{E}_n \cap \mathcal{A}_n$, (95) and (96) show that

$$T_{21} = \sqrt{\alpha_n} b_{1,n} + O(\eta_n), \quad T_{12} = \sqrt{\alpha_n} b_{2,n} + O(\eta_n),$$

where $b_{1,n}, b_{2,n}$ are centered Gaussian variables with variances bounded above and below by positive constants uniformly in n . Hence, uniformly on $\mathcal{E}_n \cap \mathcal{A}_n$,

$$\mathbb{P}(T_{21} \geq T_{11} \mid a_1, \mathcal{E}_n, \mathcal{A}_n) \leq C \bar{\Phi}(u_{1,n} - w_n),$$

and similarly

$$\mathbb{P}(T_{12} \geq T_{22} \mid a_2, \mathcal{E}_n, \mathcal{A}_n) \leq C \bar{\Phi}(u_{2,n} - w_n),$$

where $\bar{\Phi} = 1 - \Phi$. On the other hand,

$$\psi_n(t) = 1 - (1 - \bar{\Phi}(t))^{n-2}.$$

Since $\bar{\Phi}(t) \rightarrow 0$ uniformly for $t \asymp \sqrt{\log n}$, one has

$$\frac{\bar{\Phi}(t)}{\psi_n(t)} \rightarrow 0$$

uniformly on that range: if $(n-2)\bar{\Phi}(t) \leq 1$, then $\psi_n(t) \geq \frac{1}{2}(n-2)\bar{\Phi}(t)$; if $(n-2)\bar{\Phi}(t) > 1$, then $\psi_n(t) \geq 1 - e^{-1}$ and $\bar{\Phi}(t) \rightarrow 0$. Therefore the single-entry competitors T_{21} and T_{12} are negligible relative to the lower-block maxima. Combining this with (113)–(114), we obtain, uniformly on $\mathcal{E}_n \cap \mathcal{A}_n$,

$$\mathbb{P}(F_1 \mid a_1, \mathcal{E}_n, \mathcal{A}_n) = \psi_n(u_{1,n})(1 + o(1)), \quad (115)$$

$$\mathbb{P}(F_2 \mid a_2, \mathcal{E}_n, \mathcal{A}_n) = \psi_n(u_{2,n})(1 + o(1)). \quad (116)$$

Moreover, by (113)–(114), the lower-block events for columns 1 and 2 are driven by g and h , which are independent. Thus, uniformly on $\mathcal{E}_n \cap \mathcal{A}_n$,

$$\mathbb{P}(F_1 \cap F_2 \mid a_1, a_2, \mathcal{E}_n, \mathcal{A}_n) = \psi_n(u_{1,n})\psi_n(u_{2,n})(1 + o(1)). \quad (117)$$

It remains to remove the perturbation w_n . By Mills' inequality, for $t \asymp \sqrt{\log n}$ and $w_n = o((\log n)^{-1/2})$,

$$\bar{\Phi}(t \pm w_n) = \bar{\Phi}(t)(1 + o(1))$$

uniformly. It follows that $\psi_n(t \pm w_n) = \psi_n(t)(1 + o(1))$ uniformly on the same range. Since $|a_1|, |a_2| \leq L_n = \sqrt{8 \log n}$ on $\mathcal{A}_n$, both $u_{1,n}$ and $u_{2,n}$ lie in a compact multiple of $\sqrt{\log n}$, so the above uniform estimate applies.

Finally, by (101), (102), (110), and Lemma 7, we have

$$\mathbb{P}(\mathcal{E}_n^c \cup \mathcal{A}_n^c \cup \mathcal{G}_n^c) = o(\mathbb{P}(F_1)\mathbb{P}(F_2)),$$

because $n\mathbb{P}(F_1) \rightarrow \infty$ by assumption and $\mathbb{P}(F_1) = \mathbb{P}(F_2)$ by symmetry. Averaging (115), (116), and (117) over (a_1, a_2) thus gives

$$\begin{aligned} \mathbb{P}(F_1) &= \mathbb{E}[\psi_n(\alpha_n^{-1/2} + a_1)](1 + o(1)), \\ \mathbb{P}(F_2) &= \mathbb{E}[\psi_n(\alpha_n^{-1/2} + a_2)](1 + o(1)), \\ \mathbb{P}(F_1 \cap F_2) &= \mathbb{E}[\psi_n(\alpha_n^{-1/2} + a_1)\psi_n(\alpha_n^{-1/2} + a_2)](1 + o(1)). \end{aligned}$$

Since a_1 and a_2 are independent standard Gaussians, we have

$$\mathbb{E}[\psi_n(\alpha_n^{-1/2} + a_1)\psi_n(\alpha_n^{-1/2} + a_2)] = \mathbb{E}[\psi_n(\alpha_n^{-1/2} + a_1)]\mathbb{E}[\psi_n(\alpha_n^{-1/2} + a_2)].$$

Hence,

$$\mathbb{P}(F_1 \cap F_2) = \mathbb{P}(F_1)\mathbb{P}(F_2)(1 + o(1))$$

as desired. □

We are now equipped to prove Proposition 1 which we restate below.

Proposition 10 (Retrieval threshold for correlation matrix memory). *Consider the limiting regime*

$$n, d \rightarrow \infty, \quad \frac{d^2}{n \log n} \rightarrow \varrho$$

for some constant $\varrho \in [0, \infty]$. Then it holds that

$$\lim_{d \rightarrow \infty} \mathbb{P}(\text{PR}_n(W^\times)) = \begin{cases} 1 & \text{if } \varrho > 8, \\ 0 & \text{if } \varrho < 8. \end{cases}$$

Proof. By symmetry of the solution $W^\times$, the events $\{F_i\}_{i=1}^n$ are exchangeable. Hence when $\varrho > 8$, by Lemma 7 and a union bound,

$$\mathbb{P}(\text{PR}_n(W^\times)^c) \leq n\mathbb{P}(F_1) \xrightarrow{n \rightarrow \infty} 0.$$

Now suppose $\varrho < 8$ and define the nonnegative random variable

$$X_n = \sum_{i=1}^n \mathbf{1}_{F_i}$$

so that $\mathbb{P}(\text{PR}_n(W^\times)^c) = \mathbb{P}(X_n \geq 1)$. By the Paley-Zygmund inequality and the exchangeability of $\{F_i\}_{i=1}^n$, we have

$$\begin{aligned} \mathbb{P}(X_n \geq 1) &\geq \frac{(\mathbb{E}[X])^2}{\mathbb{E}[X^2]} = \frac{n^2 \mathbb{P}(F_1)^2}{n\mathbb{P}(F_1) + n(n-1)\mathbb{P}(F_1 \cap F_2)} \\ &\geq \frac{n^2 \mathbb{P}(F_1)^2}{n\mathbb{P}(F_1) + (1 + \omega_n)n^2 \mathbb{P}(F_1)^2}. \end{aligned}$$

where we have used Lemma 8 for the second inequality and $\omega_n = o_n(1)$. By Lemma 7, we have $n\mathbb{P}(F_1) \rightarrow \infty$ as $n \rightarrow \infty$ and so the right-hand side above tends to 1. Consequently $\lim_{n \rightarrow \infty} \mathbb{P}(\text{PR}_n(W^\times)^c) = 1$ under this regime. $\square$

A.4 Percentile law for correlation matrix memory

Lemma 9. *Let $s_{j,i}$ denote the entries of the score matrix $S = U^\top U V^\top V$. For each column $i \in [n]$, define*

$$\Omega_i^{(n)} := \frac{1}{n-1} \sum_{j \neq i} \mathbf{1}_{\{s_{i,i} \geq s_{j,i}\}}, \quad F_n(\omega) := \frac{1}{n} \sum_{i=1}^n \mathbf{1}_{\{\Omega_i^{(n)} \leq \omega\}}, \quad \omega \in [0, 1].$$

Let Ω be the $[0, 1]$ -valued random variable with CDF

$$F(\omega) = \Phi\left(\Phi^{-1}(\omega) - \frac{1}{\sqrt{\alpha}}\right), \quad 0 < \omega < 1,$$

and density

$$f_\Omega(\omega) = \exp\left(\frac{\Phi^{-1}(\omega)}{\sqrt{\alpha}} - \frac{1}{2\alpha}\right), \quad 0 < \omega < 1.$$

If $n/d^2 \rightarrow \alpha \in (0, \infty)$, then

1. for each fixed i , $\Omega_i^{(n)} \Rightarrow \Omega$;
2. for every $\omega \in [0, 1]$, $F_n(\omega) \rightarrow F(\omega)$ in probability.

We first prove Lemma 9(1).

Lemma 10 (One-column distributional limit). *For each $i \in [n]$, if $n/d^2 \rightarrow \alpha \in (0, \infty)$, then*

$$\Omega_i^{(n)} \Rightarrow \Omega.$$

Proof. By exchangeability, it suffices to consider $i = 1$. Using Lemma 3, we have

$$\Omega_1^{(n)} = \frac{1}{n-1} \sum_{j=1}^{n-1} \mathbf{1}\{\theta \geq \xi_j\} \quad \text{with} \quad \begin{bmatrix} \theta \\ \xi \end{bmatrix} \stackrel{(d)}{=} \rho_1 \begin{bmatrix} \frac{\rho_2 s}{r} \\ \frac{\|g\|}{\sqrt{n}} g \end{bmatrix}.$$

Write

$$\Omega_1^{(n)} \stackrel{(d)}{=} \frac{1}{n-1} \sum_{j=1}^{n-1} \mathbf{1}\{g_j \leq T_n H_n\}, \quad T_n := \frac{\rho_2 s}{r}, \quad H_n := \frac{\|g\|}{\sqrt{n}}.$$

Since $\rho_k \rightarrow 1$ in probability for each k and $n/d^2 \rightarrow \alpha$, Slutsky's theorem gives $s \Rightarrow \mathcal{N}(1, \alpha)$. Moreover, it is easy to check $r \xrightarrow{\mathbb{P}} \sqrt{\alpha}$, and therefore

$$T_n \Rightarrow \alpha^{-1/2} + Z, \quad Z \sim \mathcal{N}(0, 1). \quad (118)$$

Also, $H_n \rightarrow 1$ in probability. Now, defining

$$\tilde{\Omega}_n := \frac{1}{n-1} \sum_{j=1}^{n-1} \mathbf{1}\{g_j \leq T_n\}, \quad (119)$$

conditional on T_n , the summands in (119) are i.i.d. Bernoulli with success probability $\Phi(T_n)$, so

$$\mathbb{E}[\tilde{\Omega}_n \mid T_n] = \Phi(T_n), \quad \text{Var}(\tilde{\Omega}_n \mid T_n) = \frac{\Phi(T_n)(1 - \Phi(T_n))}{n-1} \leq \frac{1}{4(n-1)}.$$

Hence, by Chebyshev's inequality,

$$\tilde{\Omega}_n - \Phi(T_n) \xrightarrow{\mathbb{P}} 0. \quad (120)$$

It remains to replace T_n by $T_n H_n$. Since the sequence $(T_n)_{n \geq 1}$ is tight and $H_n \rightarrow 1$ in probability, for $\varepsilon > 0$ one may choose $M < \infty$ and $\delta > 0$ such that

$$\limsup_{n \rightarrow \infty} \mathbb{P}(|T_n| > M) < \varepsilon, \quad \limsup_{n \rightarrow \infty} \mathbb{P}(|H_n - 1| > \delta) < \varepsilon.$$

On the event $\{|T_n| \leq M, |H_n - 1| \leq \delta\}$, we have $|T_n H_n - T_n| \leq M\delta$, and therefore

$$|\Omega_1^{(n)} - \tilde{\Omega}_n| \leq \frac{1}{n-1} \sum_{j=1}^{n-1} \mathbf{1}\{|g_j - T_n| \leq M\delta\}. \quad (121)$$

Conditioning on T_n , the expectation of the right-hand side of (121) is bounded by

$$\sup_{x \in \mathbb{R}} \mathbb{P}(|g_j - x| \leq M\delta) \leq \frac{2M\delta}{\sqrt{2\pi}}$$

where $(2\pi)^{-1/2}$ appears from the bound on the standard normal density. Hence, taking δ arbitrarily small, by Markov's inequality,

$$\Omega_1^{(n)} - \tilde{\Omega}_n \xrightarrow{\mathbb{P}} 0. \quad (122)$$

Combining (120) and (122) implies

$$\Omega_1^{(n)} - \Phi(T_n) \xrightarrow{\mathbb{P}} 0.$$

Recalling (118), since Φ is continuous, we have

$$\Omega_1^{(n)} \Rightarrow \Omega = \Phi(\alpha^{-1/2} + Z).$$

Finally, for $\omega \in (0, 1)$,

$$\mathbb{P}(\Omega \leq \omega) = \mathbb{P}\left(Z \leq \Phi^{-1}(\omega) - \alpha^{-1/2}\right) = \Phi\left(\Phi^{-1}(\omega) - \alpha^{-1/2}\right).$$

□

For the proof of Lemma 9(2), the key ingredient is the following joint-limiting law.

Lemma 11 (Joint two-column percentile limit). *Assume $n/d^2 \rightarrow \alpha \in (0, \infty)$. Then*

$$(\Omega_1^{(n)}, \Omega_2^{(n)}) \Rightarrow (\Omega^{(1)}, \Omega^{(2)}),$$

where $\Omega^{(1)}$ and $\Omega^{(2)}$ are independent copies of Ω .

Proof. Set

$$\alpha_n := \frac{n}{d^2}, \quad S_{:,1:2} = \begin{bmatrix} T \\ \Xi \end{bmatrix},$$

where $T \in \mathbb{R}^{2 \times 2}$ and $\Xi \in \mathbb{R}^{(n-2) \times 2}$ are the top and lower blocks of the first two columns of S .

Diagonal entries. The explicit formulas from Appendix A.2 give

$$\begin{aligned} T_{11} &= c_1^2 c_7^2 + c_1 c_4 c_7 c_{10} + c_1 c_2 c_7 c_8, \\ T_{22} &= (c_2^2 + c_3^2)(c_8^2 + c_9^2) + c_1 c_2 c_7 c_8 + c_2 c_4 (c_{10} c_8 + c_{12} c_9) \\ &\quad + c_3 c_5 (c_{10} c_8 + c_{12} c_9) + c_3 c_6 (c_{11} c_8 + c_{13} c_9). \end{aligned}$$

Define

$$a_{1,n} := \sqrt{d} c_{10}, \quad a_{2,n} := \sqrt{d} c_{13}.$$

Then $a_{1,n}, a_{2,n}$ are independent standard Gaussians. Since

$$c_1, c_3, c_7, c_9 \xrightarrow{\mathbb{P}} 1, \quad c_2, c_5, c_8, c_{10}, c_{11}, c_{12}, c_{13} = O_{\mathbb{P}}(d^{-1/2}),$$

and

$$c_4 = \frac{\chi_{n-2}}{\sqrt{d}}, \quad c_6 = \frac{\chi_{n-3}}{\sqrt{d}},$$

we have

$$c_1 c_4 c_7 c_{10} = \sqrt{\alpha_n} a_{1,n} \left(c_1 c_7 \frac{\chi_{n-2}}{\sqrt{n}} \right) = \sqrt{\alpha_n} a_{1,n} + o_{\mathbb{P}}(1),$$

and likewise

$$c_3 c_6 c_{13} c_9 = \sqrt{\alpha_n} a_{2,n} \left(c_3 c_9 \frac{\chi_{n-3}}{\sqrt{n}} \right) = \sqrt{\alpha_n} a_{2,n} + o_{\mathbb{P}}(1).$$

All remaining terms are $o_{\mathbb{P}}(1)$, so

$$T_{11} = 1 + \sqrt{\alpha_n} a_{1,n} + o_{\mathbb{P}}(1), \quad T_{22} = 1 + \sqrt{\alpha_n} a_{2,n} + o_{\mathbb{P}}(1).$$

Since $\alpha_n \rightarrow \alpha$, it follows that

$$\left(\frac{T_{11}}{\sqrt{\alpha_n}}, \frac{T_{22}}{\sqrt{\alpha_n}} \right) \Rightarrow (\alpha^{-1/2} + Z_1, \alpha^{-1/2} + Z_2), \quad (123)$$

where Z_1, Z_2 are independent standard Gaussians.

Lower-block Gram matrix. Again from Appendix A.2, one has

$$\Xi = H_Z Y, \quad \Xi^\top \Xi = Y^\top Y,$$

and the last $n - 4$ rows of Y are equal to ΓB , where

$$B = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \\ b_{31} & b_{32} \\ 0 & b_{42} \end{pmatrix}$$

with

$$b_{31} = c_1 c_4 c_{14}, \quad b_{42} = c_3 c_6 c_{16},$$

and

$$b_{11}, b_{12}, b_{21}, b_{22}, b_{32} = O_{\mathbb{P}}(1).$$

Since $c_{14}, c_{16} \rightarrow 1$ in probability,

$$\frac{b_{31}^2}{d} = \alpha_n + o_{\mathbb{P}}(1), \quad \frac{b_{42}^2}{d} = \alpha_n + o_{\mathbb{P}}(1),$$

while every other entry of $d^{-1}B^{\top}B$ is $o_{\mathbb{P}}(1)$. Hence,

$$\frac{1}{d}B^{\top}B = \begin{pmatrix} \alpha_n & 0 \\ 0 & \alpha_n \end{pmatrix} + o_{\mathbb{P}}(1) \quad (124)$$

in operator norm.

The first two rows of Y are $O_{\mathbb{P}}(1)$ entrywise, so their contribution to $(n-2)^{-1}Y^{\top}Y$ is $o_{\mathbb{P}}(1)$. Therefore

$$\frac{1}{n-2}\Xi^{\top}\Xi = \frac{1}{n-2}Y^{\top}Y = \frac{1}{n-2}B^{\top}\Gamma^{\top}\Gamma B + o_{\mathbb{P}}(1).$$

Because $\Gamma \in \mathbb{R}^{(n-4) \times 4}$ has i.i.d. $\mathcal{N}(0, 1/d)$ entries and fixed column dimension,

$$\frac{d}{n-2}\Gamma^{\top}\Gamma \xrightarrow{\mathbb{P}} I_4$$

in operator norm. Combining this with (124), we obtain

$$\frac{1}{n-2}\Xi^{\top}\Xi = \begin{pmatrix} \sigma_{1,n}^2 & \rho_n \\ \rho_n & \sigma_{2,n}^2 \end{pmatrix}, \quad \sigma_{1,n}^2 = \alpha_n + o_{\mathbb{P}}(1), \quad \sigma_{2,n}^2 = \alpha_n + o_{\mathbb{P}}(1), \quad \rho_n = o_{\mathbb{P}}(1). \quad (125)$$

Gaussian representation of the lower block. For every deterministic orthogonal matrix $O \in O(n-2)$,

$$(T, \Xi) \stackrel{(d)}{=} (T, O\Xi).$$

Thus, conditionally on $(T, \Xi^{\top}\Xi)$, the law of Ξ is left-orthogonally invariant. Define

$$\tau_n^2 := \sigma_{2,n}^2 - \frac{\rho_n^2}{\sigma_{1,n}^2}.$$

Let $g_n, h_n \in \mathbb{R}^{n-2}$ be independent standard Gaussian vectors, independent of $(T, \Xi^{\top}\Xi)$, and set

$$A_n := \frac{\sqrt{n-2}}{\|g_n\|}, \quad \beta_n := \frac{\langle g_n, h_n \rangle}{\|g_n\|^2}, \quad h_n^{\perp} := h_n - \beta_n g_n, \quad C_n := \frac{\sqrt{n-2}}{\|h_n^{\perp}\|}.$$

Then, conditionally on $(T, \Xi^{\top}\Xi)$,

$$\Xi_{:,1} \stackrel{(d)}{=} \sigma_{1,n} A_n g_n, \quad \Xi_{:,2} \stackrel{(d)}{=} \frac{\rho_n}{\sigma_{1,n}} A_n g_n + \tau_n C_n h_n^{\perp}. \quad (126)$$

As $n \rightarrow \infty$, the law of large numbers gives

$$A_n \xrightarrow{\mathbb{P}} 1, \quad \beta_n \xrightarrow{\mathbb{P}} 0, \quad C_n \xrightarrow{\mathbb{P}} 1.$$

Also, by (125),

$$\tau_n^2 = \alpha_n + o_{\mathbb{P}}(1), \quad \tau_n = \sqrt{\alpha_n} + o_{\mathbb{P}}(1).$$

Using the representation (126), define

$$R_{1,n} := \frac{1}{n-2} \sum_{j=1}^{n-2} |\Xi_{j1} - \sqrt{\alpha_n} g_{n,j}|, \quad R_{2,n} := \frac{1}{n-2} \sum_{j=1}^{n-2} |\Xi_{j2} - \sqrt{\alpha_n} h_{n,j}|.$$

Then

$$R_{1,n} \leq |\sigma_{1,n} A_n - \sqrt{\alpha_n}| \cdot \frac{1}{n-2} \sum_{j=1}^{n-2} |g_{n,j}|,$$

hence $R_{1,n} \rightarrow 0$ in probability. For the second column,

$$\Xi_{:,2} - \sqrt{\alpha_n} h_n = \left(\frac{\rho_n}{\sigma_{1,n}} A_n - \tau_n C_n \beta_n \right) g_n + (\tau_n C_n - \sqrt{\alpha_n}) h_n,$$

and therefore

$$R_{2,n} \leq \left| \frac{\rho_n}{\sigma_{1,n}} A_n - \tau_n C_n \beta_n \right| \cdot \frac{1}{n-2} \sum_{j=1}^{n-2} |g_{n,j}| + |\tau_n C_n - \sqrt{\alpha_n}| \cdot \frac{1}{n-2} \sum_{j=1}^{n-2} |h_{n,j}|.$$

Since $\rho_n = o_{\mathbb{P}}(1)$ and the empirical means of $|g_{n,j}|$ and $|h_{n,j}|$ are tight, we conclude that

$$R_{1,n} \xrightarrow{\mathbb{P}} 0, \quad R_{2,n} \xrightarrow{\mathbb{P}} 0. \quad (127)$$

Replacement by Gaussian comparison variables. Define

$$\bar{\Omega}_{1,n} := \frac{1}{n-2} \sum_{j=1}^{n-2} \mathbf{1}_{\{\Xi_{j1} \leq T_{11}\}}, \quad \bar{\Omega}_{2,n} := \frac{1}{n-2} \sum_{j=1}^{n-2} \mathbf{1}_{\{\Xi_{j2} \leq T_{22}\}},$$

and

$$\hat{\Omega}_{1,n} := \frac{1}{n-2} \sum_{j=1}^{n-2} \mathbf{1}_{\{\sqrt{\alpha_n} g_{n,j} \leq T_{11}\}}, \quad \hat{\Omega}_{2,n} := \frac{1}{n-2} \sum_{j=1}^{n-2} \mathbf{1}_{\{\sqrt{\alpha_n} h_{n,j} \leq T_{22}\}}.$$

Because the original percentiles use denominator $n-1$ and include one additional competitor from the top block, we have the deterministic bounds

$$|\Omega_1^{(n)} - \bar{\Omega}_{1,n}| \leq \frac{2}{n-1}, \quad |\Omega_2^{(n)} - \bar{\Omega}_{2,n}| \leq \frac{2}{n-1}. \quad (128)$$

We next show that $\bar{\Omega}_{1,n} - \hat{\Omega}_{1,n} \rightarrow 0$ in probability. For every $x, y, t \in \mathbb{R}$ and every $\delta > 0$,

$$|\mathbf{1}_{\{x \leq t\}} - \mathbf{1}_{\{y \leq t\}}| \leq \mathbf{1}_{\{|x-y| > \delta\}} + \mathbf{1}_{\{|y-t| \leq \delta\}} \leq \frac{|x-y|}{\delta} + \mathbf{1}_{\{|y-t| \leq \delta\}}.$$

Applying this with $x = \Xi_{j1}$, $y = \sqrt{\alpha_n}g_{n,j}$, $t = T_{11}$ and averaging over j ,

$$|\bar{\Omega}_{1,n} - \hat{\Omega}_{1,n}| \leq \frac{R_{1,n}}{\delta} + \frac{1}{n-2} \sum_{j=1}^{n-2} \mathbf{1}_{\{|\sqrt{\alpha_n}g_{n,j} - T_{11}| \leq \delta\}}.$$

Now g_n is independent of T_{11} , and since $\alpha_n \rightarrow \alpha > 0$, the density of $\sqrt{\alpha_n}G$ is uniformly bounded for all large n . Therefore

$$\sup_{x \in \mathbb{R}} \mathbb{P}(|\sqrt{\alpha_n}G - x| \leq \delta) \leq C\delta$$

for some constant $C < \infty$ and all sufficiently large n . Using Markov's inequality together with (127), we obtain

$$\bar{\Omega}_{1,n} - \hat{\Omega}_{1,n} \xrightarrow{\mathbb{P}} 0.$$

The same argument gives

$$\bar{\Omega}_{2,n} - \hat{\Omega}_{2,n} \xrightarrow{\mathbb{P}} 0.$$

Together with (128), this yields

$$(\Omega_1^{(n)}, \Omega_2^{(n)}) - (\hat{\Omega}_{1,n}, \hat{\Omega}_{2,n}) \xrightarrow{\mathbb{P}} (0, 0). \quad (129)$$

Conditional law of large numbers. Conditionally on (T_{11}, T_{22}) , the indicators defining $\hat{\Omega}_{1,n}$ are i.i.d. Bernoulli with mean

$$p_{1,n} := \Phi\left(\frac{T_{11}}{\sqrt{\alpha_n}}\right),$$

and the indicators defining $\hat{\Omega}_{2,n}$ are i.i.d. Bernoulli with mean

$$p_{2,n} := \Phi\left(\frac{T_{22}}{\sqrt{\alpha_n}}\right).$$

Hence

$$\mathbb{E}\left[(\hat{\Omega}_{1,n} - p_{1,n})^2 \mid T_{11}, T_{22}\right] \leq \frac{1}{4(n-2)}, \quad \mathbb{E}\left[(\hat{\Omega}_{2,n} - p_{2,n})^2 \mid T_{11}, T_{22}\right] \leq \frac{1}{4(n-2)},$$

so

$$(\hat{\Omega}_{1,n}, \hat{\Omega}_{2,n}) - (p_{1,n}, p_{2,n}) \xrightarrow{\mathbb{P}} (0, 0). \quad (130)$$

Finally, (123) and the continuity of Φ imply

$$(p_{1,n}, p_{2,n}) = \left(\Phi\left(\frac{T_{11}}{\sqrt{\alpha_n}}\right), \Phi\left(\frac{T_{22}}{\sqrt{\alpha_n}}\right)\right) \Rightarrow (\Phi(\alpha^{-1/2} + Z_1), \Phi(\alpha^{-1/2} + Z_2)).$$

The two coordinates on the right-hand side are independent copies of Ω . Combining this with (129) and (130), Slutsky's theorem yields

$$(\Omega_1^{(n)}, \Omega_2^{(n)}) \Rightarrow (\Omega^{(1)}, \Omega^{(2)}).$$

□

We are now ready to prove Lemma 9(2) which we restate below.

Lemma 12 (Empirical distribution across columns). *For every $\omega \in [0, 1]$,*

$$F_n(\omega) \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \mathbb{P}(\Omega \leq \omega).$$

Proof. Fix $\omega \in [0, 1]$. By exchangeability,

$$\text{Var}(F_n(\omega)) = \frac{1}{n} \text{Var}(\mathbf{1}\{\Omega_1^{(n)} \leq \omega\}) + \frac{n-1}{n} \text{Cov}(\mathbf{1}\{\Omega_1^{(n)} \leq \omega\}, \mathbf{1}\{\Omega_2^{(n)} \leq \omega\}).$$

The first term is at most $1/(4n)$. For the second term, since $\mathbb{P}(\Omega = \omega) = 0$, the extended continuous mapping theorem and Lemma 11 give

$$(\mathbf{1}\{\Omega_1^{(n)} \leq \omega\}, \mathbf{1}\{\Omega_2^{(n)} \leq \omega\}) \Rightarrow (\mathbf{1}\{\Omega^{(1)} \leq \omega\}, \mathbf{1}\{\Omega^{(2)} \leq \omega\}), \quad (131)$$

where $\Omega^{(1)}, \Omega^{(2)}$ are independent copies of Ω . Obviously, the indicators on the right-hand side of (131) are uniformly bounded and so $\text{Cov}(\mathbf{1}\{\Omega_1^{(n)} \leq \omega\}, \mathbf{1}\{\Omega_2^{(n)} \leq \omega\}) \rightarrow 0$. Hence, $\text{Var}(F_n(\omega)) \rightarrow 0$, and so Chebyshev's inequality gives

$$F_n(\omega) - \mathbb{E}[F_n(\omega)] \xrightarrow{\mathbb{P}} 0. \quad (132)$$

Now, by exchangeability,

$$\mathbb{E}[F_n(\omega)] = \mathbb{P}(\Omega_1^{(n)} \leq \omega) \longrightarrow \mathbb{P}(\Omega \leq \omega), \quad (133)$$

where the limit follows from Lemma 9(1) and the continuity of the law of Ω . Combining (132) and (133) gives the desired result. $\square$

B Auxiliary estimates for the universal top-1 bound

Proof of Lemma 1. Let $\mathcal{M} \subset S^{d-1}$ be a $1/d$ -net of the sphere and fix $y \in \mathcal{M}$. By the standard volumetric bound, $|\mathcal{M}| \leq (1+2d)^d \leq e^{Cd \log d}$ for some $C > 0$. Since $u_j^\top y \sim \mathcal{N}(0, \frac{1}{d})$ are i.i.d., Mills' inequality gives

$$\mathbb{P} \left(u_j^\top y \geq \sqrt{\frac{(2-\epsilon) \log d}{d}} \right) \geq \frac{\sqrt{(2-\epsilon) \log d}}{(2-\epsilon) \log d + 1} \cdot \frac{e^{-\frac{1}{2}(2-\epsilon) \log d}}{\sqrt{2\pi}} \geq \frac{cd^{\epsilon/2-1}}{\sqrt{\log d}}$$

for some $c > 0$. By a union bound, the probability that no such j exists for some $y \in \mathcal{M}$ is at most

$$|\mathcal{M}| \left(1 - \frac{cd^{\epsilon/2-1}}{\sqrt{\log d}} \right)^n \leq \exp \left(Cd \log d - \Omega \left(\frac{d^{\epsilon/2+1}}{(\log d)^{3/2}} \right) \right) = e^{-\Omega(d)}.$$

On the complement of this event, for arbitrary $x \in S^{d-1}$, choose $y \in \mathcal{M}$ with $\|x - y\| \leq d^{-1}$ and then choose j as above. Using $\|u_j\| \leq 2$ with probability $1 - e^{-\Omega(d)}$, we get

$$u_j^\top x \geq u_j^\top y - \|u_j\| \|x - y\| \geq \sqrt{\frac{(2-\epsilon) \log d}{d}} - \frac{2}{d} \geq \sqrt{\frac{(2-2\epsilon) \log d}{d}}$$

for all sufficiently large d . The statement follows by replacing ϵ with $\epsilon/2$. $\square$

Proof of Lemma 2. First consider the Gaussian case. Recall $V = [v_1 \ \cdots \ v_n] \in \mathbb{R}^{d \times n}$ and define, for $x \in S^{d-1}$,

$$f_x(V) := \frac{1}{n} \sum_{i=1}^n |x^\top v_i|.$$

It holds that $\mathbb{E}[f_x(V)] = \sqrt{2/\pi d}$ and f_x is $n^{-1/2}$ -Lipschitz:

$$|f_x(V) - f_x(V')| \leq \frac{1}{n} \sum_{i=1}^n \|v_i - v'_i\| \leq \frac{\|V - V'\|_{\text{F}}}{\sqrt{n}}.$$

Since $\sqrt{d}V$ is standard Gaussian, Lipschitz concentration gives, for every $t > 0$,

$$\mathbb{P}\left(|f_x(V) - \mathbb{E}f_x(V)| \geq \frac{t}{\sqrt{d}}\right) \leq 2 \exp\left(-\frac{nt^2}{2}\right).$$

Thus $f_x(V) \geq \sqrt{(2-\epsilon)/\pi d}$ with probability $1 - e^{-\Omega(n)}$ for any fixed $\epsilon > 0$. Choosing a $1/d$ -net $\mathcal{M} \subset S^{d-1}$ and applying a union bound, we have $f_y(V) \geq \sqrt{(2-\epsilon)/\pi d}$ simultaneously for all $y \in \mathcal{M}$ with failure probability $|\mathcal{M}|e^{-\Omega(n)} = e^{Cd \log d - \Omega(n)} = e^{-\Omega(n)}$. For any $x \in S^{d-1}$, choose $y \in \mathcal{M}$ with $\|x - y\| \leq d^{-1}$. Using $\|a\| - \|b\| \leq \|a - b\|$ and $\|v_i\| \leq 2$ with high probability,

$$f_x(V) \geq f_y(V) - \frac{1}{n} \sum_{i=1}^n \|x - y\| \|v_i\| \geq \sqrt{\frac{2-\epsilon}{\pi d}} - \frac{1}{d} \geq \sqrt{\frac{2-2\epsilon}{\pi d}}.$$

The Gaussian statement follows by replacing ϵ with $\epsilon/2$.

Alternatively, suppose (11) holds. Define the k -tensor $M_k := n^{-1} \sum_{i=1}^n v_i^{\otimes k}$. By Hölder's inequality, for every $x \in S^{d-1}$,

$$\left(\frac{1}{n} \sum_{i=1}^n (x^\top v_i)^2\right)^{3/2} \leq \left(\frac{1}{n} \sum_{i=1}^n |x^\top v_i|\right) \left(\frac{1}{n} \sum_{i=1}^n (x^\top v_i)^4\right)^{1/2}.$$

Consequently,

$$\frac{1}{n} \sum_{i=1}^n |x^\top v_i| \geq \frac{(x^\top M_2 x)^{3/2}}{\langle M_4, x^{\otimes 4} \rangle^{1/2}} \gtrsim \frac{d^{-3/2}}{\|M_4\|^{1/2}} \gtrsim \frac{1}{\sqrt{d}}.$$

Taking the infimum over $x \in S^{d-1}$ completes the proof. $\square$

Lemma 13. For any fixed $\epsilon > 0$, it holds w.p. $1 - e^{-\Omega(d)}$ that

$$\|UV^\top\|_{\text{F}} \leq (1 + \epsilon)\sqrt{n}.$$

Proof. By standard Gaussian concentration, with probability $1 - e^{-\Omega(d)}$ we have

$$\|V\|_{\text{F}} \leq (1 + \epsilon)\sqrt{n}, \quad \|V\|_{\text{op}} \lesssim \sqrt{\frac{n}{d}}.$$

We condition on this event. Defining $f(U) := \|UV^\top\|_F$, it holds that

$$\mathbb{E}[f(U) \mid V] \leq \mathbb{E}[f(U)^2 \mid V]^{1/2} = \text{tr}(V \mathbb{E}[U^\top U] V^\top)^{1/2} = \|V\|_F \leq (1 + \epsilon)\sqrt{n}.$$

Moreover f is $\|V\|_{\text{op}}$ -Lipschitz, hence Lipschitz concentration yields

$$f(U) \leq \mathbb{E}[f(U) \mid V] + \|V\|_{\text{op}} \leq (1 + \epsilon)\sqrt{n} + o(\sqrt{n}) \leq (1 + 2\epsilon)\sqrt{n}$$

with probability $1 - e^{-\Omega(d)}$ for sufficiently large d . The statement follows by replacing ϵ with $\epsilon/2$. $\square$

C Taylor expansion of the interior remainder

This appendix carries out the first-order Taylor expansion of the interior remainder $\mathcal{R}^{\setminus h}$ defined in (31) of Section 4.2, derives the explicit form of the residual $\hat{E}$ appearing in the main identity (34), and establishes the entry-size bounds (35) used there. All notation is as in Section 4.2: $(\widehat{W}, \widehat{\mu})$ and $(\widehat{W}^{\setminus h}, \widehat{\mu}^{\setminus h})$ are the full and leave-one-out minimizers, $\Delta S := U^\top(\widehat{W} - \widehat{W}^{\setminus h})V$, $(\Delta\mu)_i := \widehat{\mu}_i - \mu_i^{\setminus h}$; a hat denotes a quantity evaluated at $(\widehat{W}, \widehat{\mu})$ and a superscript $\setminus h$ one evaluated at $(\widehat{W}^{\setminus h}, \widehat{\mu}^{\setminus h})$. The analysis is conditional on the LOO postulates (A1)–(A3).

C.1 Taylor identities

We record three Taylor identities around the LOO point. Throughout, for a scalar quantity $X(W, \mu)$ we write $\Delta X := X(\widehat{W}, \widehat{\mu}) - X(\widehat{W}^{\setminus h}, \widehat{\mu}^{\setminus h})$.

Distractor sigmoid. Since $q_{j,i} = \phi'_\beta(s_{j,i} - \mu_i) = \sigma(\beta(s_{j,i} - \mu_i))$ and $\sigma'(t) = \sigma(t)(1 - \sigma(t))$, a Taylor expansion in the single variable $s_{j,i} - \mu_i$ gives, for $j, i \neq h$ and $j \neq i$,

$$\begin{aligned} \hat{q}_{j,i} - q_{j,i}^{\setminus h} &= k \widetilde{R}_{j,i}^{\setminus h} (\Delta S_{j,i} - (\Delta\mu)_i) + \rho_{j,i}^{(q)}, \\ \rho_{j,i}^{(q)} &= \mathcal{O}_{\prec}(\beta^2 |\Delta S_{j,i} - (\Delta\mu)_i|^2) = \mathcal{O}_{\prec}(\beta^2/d^2), \end{aligned} \quad (134)$$

where $k \widetilde{R}_{j,i}^{\setminus h} = \beta q_{j,i}^{\setminus h} (1 - q_{j,i}^{\setminus h}) = \phi''_\beta(s_{j,i}^{\setminus h} - \mu_i^{\setminus h})$ and the last bound uses $\max_{j,i \neq h} |\Delta S_{j,i}| = \mathcal{O}_{\prec}(d^{-1})$ by (A2) and $|\Delta\mu_i| = \mathcal{O}_{\prec}(n^{-1}) = \mathcal{O}_{\prec}(d^{-2})$ by (A1).

Margin. For $i \neq h$, the full margin $\hat{m}_i = s_{i,i}(\widehat{W}) - \hat{c}_i$ and the LOO margin $m_i^{\setminus h} = s_{i,i}(\widehat{W}^{\setminus h}) - c_i^{\setminus h}$ differ both through the shifts $(\Delta S_{i,i}, (\Delta\mu)_i)$ and through the missing $j = h$ interferer in the LOO inner sum. Taylor-expanding the inner softplus summation in the column- i score perturbations and peeling off the missing interferer term,

$$\hat{m}_i - m_i^{\setminus h} = - (Q_{:,i}^{\setminus h})^\top \Delta S_{:,i} + [(Q_{:,i}^{\setminus h})^\top \mathbf{1}_n] (\Delta\mu)_i - \frac{1}{k} \phi_\beta(s_{h,i}^{\setminus h} - \mu_i^{\setminus h}) + \rho_i^{(m)}, \quad (135)$$

where the first two terms assemble the linear dependence on $(\Delta S_{:,i}, (\Delta\mu)_i)$ (using $\partial m_i / \partial s_{j,i} = -Q_{j,i}$ with the convention $Q_{i,i} = -1$, $Q_{j,i} = q_{j,i}/k$ for $j \neq i$), the third term is the removed-interferer contribution, and $\rho_i^{(m)}$ collects second-order Taylor remainders of the ϕ_β terms and

cross contributions. Using the second-order softplus bound $|\phi_\beta(x+\delta) - \phi_\beta(x) - \phi'_\beta(x)\delta| \leq \frac{\beta}{8} \delta^2$ together with (A1)–(A2),

$$\begin{aligned} |\rho_i^{(m)}| &\leq \frac{\beta}{8k} \sum_{j \neq i, j \neq h} (\Delta S_{j,i} - (\Delta\mu)_i)^2 + \mathcal{O}_{\prec}(|\Delta S_{h,i}| + |(\Delta\mu)_i|) \cdot \frac{1}{k} \\ &= \mathcal{O}_{\prec}(\beta/d^2) + \mathcal{O}_{\prec}(1/(nd)) = \mathcal{O}_{\prec}(1/d^2). \end{aligned} \quad (136)$$

Activation. Since $\ell'(t) = -\sigma(-t)$, $\ell''(t) = \sigma'(-t) = a(1-a) = b$, and ℓ''' is uniformly bounded,

$$\begin{aligned} \hat{a}_i - a_i^{\setminus h} &= -b_i^{\setminus h} (\hat{m}_i - m_i^{\setminus h}) + \rho_i^{(a)}, \\ |\rho_i^{(a)}| &= \mathcal{O}_{\prec}((\hat{m}_i - m_i^{\setminus h})^2) = \mathcal{O}_{\prec}(1/d^2) = \mathcal{O}_{\prec}(1/n). \end{aligned} \quad (137)$$

C.2 Explicit form of the residual

We now unfold the two cases of $\mathcal{R}^{\setminus h}$ in (31) and identify the linear part with $(M_i^{\setminus h} \Delta S_{:,i})_j$ from (27), so that $\hat{E}_{j,i}$ is what remains.

Off-diagonal case ($j \neq i, j, i \neq h$). Using the algebraic identity

$$\hat{a}_i \hat{q}_{j,i} - a_i^{\setminus h} q_{j,i}^{\setminus h} = a_i^{\setminus h} (\hat{q}_{j,i} - q_{j,i}^{\setminus h}) + (\hat{a}_i - a_i^{\setminus h}) q_{j,i}^{\setminus h} + (\hat{a}_i - a_i^{\setminus h}) (\hat{q}_{j,i} - q_{j,i}^{\setminus h}),$$

and substituting (134), (135), (137),

$$\begin{aligned} \mathcal{R}_{j,i}^{\setminus h} &= \frac{a_i^{\setminus h} \cdot k \tilde{R}_{j,i}^{\setminus h}}{k} (\Delta S_{j,i} - (\Delta\mu)_i) + \frac{q_{j,i}^{\setminus h}}{k} \cdot [-b_i^{\setminus h} (\hat{m}_i - m_i^{\setminus h})] + \hat{E}_{j,i}^{(\text{off})} \\ &= a_i^{\setminus h} \tilde{R}_{j,i}^{\setminus h} (\Delta S_{j,i} - (\Delta\mu)_i) + b_i^{\setminus h} Q_{j,i}^{\setminus h} (Q_{:,i}^{\setminus h})^\top \Delta S_{:,i} + \hat{E}_{j,i}^{(\text{off})}, \end{aligned}$$

where in the second equality we used $Q_{j,i}^{\setminus h} = q_{j,i}^{\setminus h}/k$ for $j \neq i, j \neq h$, and absorbed the $(\Delta\mu)_i$ -dependent piece of $\hat{m}_i - m_i^{\setminus h}$ from (135) into $\hat{E}_{j,i}^{(\text{off})}$. The first two terms match $(M_i^{\setminus h} \Delta S_{:,i})_j$ for $j \neq i$ from (27), provided we recognize that the $a_i^{\setminus h} \tilde{R}$ piece also absorbs the $-a_i^{\setminus h} \tilde{R}_{j,i}^{\setminus h} (\Delta\mu)_i$ contribution into $\hat{E}$. Collecting the residual pieces,

$$\begin{aligned} \hat{E}_{j,i}^{(\text{off})} &= -\frac{b_i^{\setminus h} q_{j,i}^{\setminus h}}{k^2} \phi_\beta(s_{h,i}^{\setminus h} - \mu_i^{\setminus h}) + \frac{a_i^{\setminus h}}{k} \rho_{j,i}^{(q)} - a_i^{\setminus h} \tilde{R}_{j,i}^{\setminus h} (\Delta\mu)_i \\ &\quad + \frac{q_{j,i}^{\setminus h}}{k} [b_i^{\setminus h} (Q_{:,i}^{\setminus h})^\top \mathbf{1}_n (\Delta\mu)_i - \rho_i^{(a)}] + \frac{(\hat{a}_i - a_i^{\setminus h})(\hat{q}_{j,i} - q_{j,i}^{\setminus h})}{k}. \end{aligned} \quad (138)$$

Diagonal case ($j = i \neq h$). We have $\mathcal{R}_{i,i}^{\setminus h} = a_i^{\setminus h} - \hat{a}_i = b_i^{\setminus h} (\hat{m}_i - m_i^{\setminus h}) - \rho_i^{(a)}$. Substituting (135),

$$\begin{aligned} \mathcal{R}_{i,i}^{\setminus h} &= b_i^{\setminus h} \cdot \left[-(Q_{:,i}^{\setminus h})^\top \Delta S_{:,i} + [(Q_{:,i}^{\setminus h})^\top \mathbf{1}_n] (\Delta\mu)_i - \frac{1}{k} \phi_\beta(s_{h,i}^{\setminus h} - \mu_i^{\setminus h}) + \rho_i^{(m)} \right] - \rho_i^{(a)} \\ &= -b_i^{\setminus h} (Q_{:,i}^{\setminus h})^\top \Delta S_{:,i} + \hat{E}_{i,i}, \end{aligned}$$

where the first term equals $(M_i^{\setminus h} \Delta S_{:,i})_i = b_i^{\setminus h} Q_{i,i}^{\setminus h} (Q_{:,i}^{\setminus h})^\top \Delta S_{:,i} = -b_i^{\setminus h} (Q_{:,i}^{\setminus h})^\top \Delta S_{:,i}$ (using $Q_{i,i}^{\setminus h} = -1$ and $\tilde{R}_{i,i}^{\setminus h} = 0$), and

$$\hat{E}_{i,i} = -\frac{b_i^{\setminus h}}{k} \phi_\beta(s_{h,i}^{\setminus h} - \mu_i^{\setminus h}) + b_i^{\setminus h} [(Q_{:,i}^{\setminus h})^\top \mathbf{1}_n] (\Delta \mu)_i + b_i^{\setminus h} \rho_i^{(m)} - \rho_i^{(a)}. \quad (139)$$

C.3 Entry-size bounds

We now bound each term in (138) and (139) under (A1)–(A3), arriving at the claim (35). Throughout, we use the deterministic bounds $a_i, b_i, q_{j,i} \in [0, 1]$ (with $b_i \leq 1/4$) and $|\tilde{R}_{j,i}^{\setminus h}| \leq \beta/(4k)$.

We first record the only pointwise consequence of (A3) needed below. The μ -stationarity condition for $\Psi^{\setminus h}$ gives $\frac{1}{n-1} \sum_{j \neq i, h} \sigma(\beta(s_{j,i}^{\setminus h} - \mu_i^{\setminus h})) \approx r_d$ for each $i \neq h$. Since the full scores are bounded by (A3), the LOO scores with $i \neq h$ are bounded by (A2). As $r_d \in (0, 1)$, this implies $\mu_i^{\setminus h} = \mathcal{O}_{\prec}(1)$ uniformly. Therefore, by Lipschitz continuity of the softplus,

$$|\phi_\beta(s_{j,i}^{\setminus h} - \mu_i^{\setminus h})| \leq \frac{\log 2}{\beta} + |s_{j,i}^{\setminus h} - \mu_i^{\setminus h}| = \mathcal{O}_{\prec}(1) \quad (140)$$

uniformly in (j, i) .

Off-diagonal entries. For $j \neq i$, $j, i \neq h$, each term in (138):

- (i) $\left| \frac{b_i^{\setminus h} q_{j,i}^{\setminus h}}{k^2} \phi_\beta(\cdot) \right| \leq \frac{1}{4k^2} \cdot \mathcal{O}_{\prec}(1) = \mathcal{O}_{\prec}(n^{-2})$ by (140);
- (ii) $\left| \frac{a_i^{\setminus h}}{k} \rho_{j,i}^{(q)} \right| \leq \frac{1}{k} \cdot \mathcal{O}_{\prec}(\beta^2/d^2) = \mathcal{O}_{\prec}(\beta^2/(nd^2)) = \mathcal{O}_{\prec}(n^{-2})$, using $\beta = \mathcal{O}_{\prec}(1)$ in the retrieval regime, or absorbing β as a constant;
- (iii) $|a_i^{\setminus h} \tilde{R}_{j,i}^{\setminus h} (\Delta \mu)_i| \leq \frac{\beta}{4k} \cdot \mathcal{O}_{\prec}(n^{-1}) = \mathcal{O}_{\prec}(n^{-2})$ by (A1);
- (iv) $\left| \frac{q_{j,i}^{\setminus h}}{k} b_i^{\setminus h} (Q_{:,i}^{\setminus h})^\top \mathbf{1}_n (\Delta \mu)_i \right| \leq \frac{1}{k} \cdot \frac{1}{4} \cdot \mathcal{O}_{\prec}(1) \cdot \mathcal{O}_{\prec}(n^{-1}) = \mathcal{O}_{\prec}(n^{-2})$, using $|(Q_{:,i}^{\setminus h})^\top \mathbf{1}_n| = |-1 + (1/k) \sum q| = \mathcal{O}_{\prec}(1)$;
- (v) $\left| \frac{q_{j,i}^{\setminus h}}{k} \rho_i^{(a)} \right| \leq \frac{1}{k} \cdot \mathcal{O}_{\prec}(1/n) = \mathcal{O}_{\prec}(n^{-2})$ by (137);
- (vi) $\left| \frac{(\hat{a}_i - a_i^{\setminus h})(\hat{q}_{j,i} - q_{j,i}^{\setminus h})}{k} \right| \leq \frac{1}{k} \cdot \mathcal{O}_{\prec}(d^{-1}) \cdot \mathcal{O}_{\prec}(d^{-1}) = \mathcal{O}_{\prec}(1/(nd^2)) = \mathcal{O}_{\prec}(n^{-2})$, using (134), (137) and $|\hat{m} - m^{\setminus h}| = \mathcal{O}_{\prec}(d^{-1})$.

Summing, $|\hat{E}_{j,i}^{(\text{off})}| = \mathcal{O}_{\prec}(n^{-2})$.

Diagonal entries. For $i \neq h$, each term in (139):

- (i) $\left| \frac{b_i^{\setminus h}}{k} \phi_\beta(\cdot) \right| \leq \frac{1}{4k} \cdot \mathcal{O}_{\prec}(1) = \mathcal{O}_{\prec}(n^{-1})$ by (140);
- (ii) $|b_i^{\setminus h} (Q_{:,i}^{\setminus h})^\top \mathbf{1}_n (\Delta \mu)_i| \leq \frac{1}{4} \cdot \mathcal{O}_{\prec}(1) \cdot \mathcal{O}_{\prec}(n^{-1}) = \mathcal{O}_{\prec}(n^{-1})$ by (A1);

(iii) $|b_i^{\setminus h} \rho_i^{(m)}| \leq \frac{1}{4} \cdot \mathcal{O}_{\prec}(d^{-2}) = \mathcal{O}_{\prec}(n^{-1})$ by (136);

(iv) $|\rho_i^{(a)}| = \mathcal{O}_{\prec}(n^{-1})$ by (137).

Summing, $|\hat{E}_{i,i}| = \mathcal{O}_{\prec}(n^{-1})$.

This completes the proof of (35) under (A1)–(A3).

D A priori bounds on the Frobenius norm of $\widehat{W}$

This appendix establishes two-sided control on $\|\widehat{W}\|_{\text{F}}/d$ for the joint TAM minimizer of (18). The upper bound is deterministic and uniform in the data; the lower bound holds with overwhelming probability ($1 - d^{-D}$ for every $D > 0$) in the regime $n/d^2 \rightarrow \alpha \in (0, \infty)$. Together, they pin

$$c_0 - O(d^{-D}) \leq \nu_d \leq C_0,$$

for deterministic constants $0 < c_0 \leq C_0 < \infty$ depending only on the parameters $(\alpha, r, \beta, \lambda)$, where $\nu_d^2 = \mathbb{E}[\|\widehat{W}\|_{\text{F}}^2/d^2]$ as in (47).

The two bounds provide the a priori scale for $\|\widehat{W}\|_{\text{F}}$ needed in Section 4.3 and Corollary 1. The corresponding Frobenius concentration estimate is treated there as a consequence of the empirical concentration postulate (A5), applied to the observables in the energy identity.

Throughout the appendix, C, c denote positive constants that may depend on $(\alpha, r, \beta, \lambda)$ but not on d , and may vary line to line.

D.1 Upper bound

Lemma 14 (Deterministic upper bound). *For any $\lambda > 0$, $\beta > 0$, $r \in (0, 1)$, and any data matrices $U, V \in \mathbb{R}^{d \times n}$,*

$$\frac{\|\widehat{W}\|_{\text{F}}^2}{d^2} \leq \frac{2}{\lambda} \left[\log 2 + \frac{\log 2}{r\beta} \right] =: C_0. \quad (141)$$

In particular, $\nu_d^2 \leq C_0$.

Proof. Evaluate Ψ at the test point $W = 0$, $\mu = 0$. All scores vanish, so

$$\Psi(0, 0) = n \cdot \ell \left(-\frac{1}{k} (n-1) \phi_{\beta}(0) \right) = n \cdot \ell \left(-\frac{\log 2}{r\beta} \right),$$

using $\phi_{\beta}(0) = (\log 2)/\beta$ and $r = k/(n-1)$. The function ℓ is convex with $|\ell'| \leq 1$, so

$$\ell(t) \leq \ell(0) + |t| = \log 2 + |t|. \quad (142)$$

Applying (142) at $t = -(\log 2)/(r\beta)$ gives

$$\Psi(0, 0) \leq n \left[\log 2 + \frac{\log 2}{r\beta} \right]. \quad (143)$$

By optimality of $(\widehat{W}, \widehat{\mu})$ and nonnegativity of ℓ ,

$$\frac{\lambda n}{2d^2} \|\widehat{W}\|_{\text{F}}^2 \leq \Psi(\widehat{W}, \widehat{\mu}) \leq \Psi(0, 0). \quad (144)$$

Combining (143) and (144) and dividing by $\lambda n/(2d^2)$ yields (141). The bound on ν_d^2 follows from $\nu_d^2 = \mathbb{E}\|\widehat{W}\|_F^2/d^2$. $\square$ $\square$

Remark 7 (Dependence on β). *The dependence on β enters as $1/\beta$, which vanishes in the large- β limit, leaving the cleaner asymptotic bound $C_0 \rightarrow 2 \log 2/\lambda$.*

D.2 Lower bound

Lemma 15 (High probability lower bound). *Fix $r, \alpha \in (0, 1) \times (0, \infty)$ and $\beta, \lambda > 0$, and consider the regime $n/d^2 \rightarrow \alpha$. There exists a constant $c_0 > 0$ such that, for every $D > 0$, there is $d_0 = d_0(D)$ for which*

$$\mathbb{P}\left(\frac{\|\widehat{W}\|_F}{d} \geq c_0\right) \geq 1 - d^{-D}, \quad d \geq d_0. \quad (145)$$

In particular, $\nu_d^2 \geq c_0^2 - O(d^{-D})$ for every $D > 0$.

Remark 8 (Strategy). *The argument is by objective gap. We exhibit a one-dimensional family $W_t = t W^\times$ along the direction*

$$W^\times := UV^\top$$

along which $\Psi(W_t, \mu_0 \mathbf{1})$ decreases by $\Theta(n) = \Theta(d^2)$ at small $t > 0$, and complement this with the convex linearization $\Psi(W, \mu_0 \mathbf{1}) \geq \Psi(0, \mu_0 \mathbf{1}) + \langle \nabla_W \Psi(0, \mu_0 \mathbf{1}), W \rangle$, whose linear term is bounded by $\|\nabla_W \Psi(0, \mu_0 \mathbf{1})\|_F \|W\|_F = O(d \|W\|_F)$. Comparing the two excludes a Frobenius ball of radius ηd for small η .

Proof. The variable μ is unregularized; we choose a fixed $\mu_0 \in \mathbb{R}$ and track the objective along the slice $\mu = \mu_0 \mathbf{1}_n$ throughout. Let

$$\mu_0 := -\frac{1}{\beta} \log \frac{r}{1-r}, \quad q_0 := \sigma(-\beta \mu_0) = r,$$

so that, with $f(\mu) := -\mu - \frac{1}{r} \phi_\beta(-\mu)$, we have $f'(\mu_0) = -1 + \frac{1}{r} q_0 = 0$. Define $a_0 := \sigma(-f(\mu_0)) \in (0, 1)$, the activation associated with the test profile $(0, \mu_0 \mathbf{1}_n)$.

Step 1: Gradient at the test point. By the gradient formula (24) evaluated at $W = 0$, $\mu = \mu_0 \mathbf{1}_n$, all $a_i = a_0$ and $q_{j,i} = q_0 = r$, hence

$$\nabla_W \Psi(0, \mu_0 \mathbf{1}_n) = -a_0 U M V^\top, \quad M_{ii} = 1, \quad M_{ij} = -\frac{1}{n-1} \text{ for } j \neq i. \quad (146)$$

The matrix M is symmetric with eigenvalues 0 (on $\mathbf{1}_n$) and $n/(n-1)$ (on $\mathbf{1}_n^\perp$, with multiplicity $n-1$); in particular, $\|M\|_F^2 = n^2/(n-1)$ and $\|M\|_{\text{op}} = n/(n-1)$.

By independence of U and V together with $\mathbb{E}[U^\top U] = \mathbb{E}[V^\top V] = I_n$ (since each column has variance I_d/d),

$$\mathbb{E}\|U M V^\top\|_F^2 = \text{tr}(M \mathbb{E}[U^\top U] M \mathbb{E}[V^\top V]) = \text{tr}(M^2) = \frac{n^2}{n-1}. \quad (147)$$

Hanson–Wright applied to the Gaussian quadratic form $\|UMV^\top\|_F^2$, viewed as a degree-4 polynomial in the i.i.d. $\mathcal{N}(0, 1)$ entries of $\sqrt{d}U$ and $\sqrt{d}V$, has variance $O(n)$ and operator-norm scale $O(1)$, yielding sub-exponential concentration: for any $\delta > 0$,

$$\mathbb{P}(\|UMV^\top\|_F^2 > (1 + \delta) \operatorname{tr}(M^2)) \leq \exp(-c d^2),$$

for a constant $c > 0$ depending only on δ . Hence there exists a constant $K = K(\alpha) > 0$ such that, for every $D > 0$, the event

$$\Omega_d^{(1)} := \{ \|\nabla_W \Psi(0, \mu_0 \mathbf{1}_n)\|_F \leq a_0 K d \} \quad (148)$$

satisfies $\mathbb{P}(\Omega_d^{(1)}) \geq 1 - d^{-D}$ for all sufficiently large d .

Step 2: Directional derivative along $W^\times$. Define $g(t) := \Psi(tW^\times, \mu_0 \mathbf{1}_n)$, viewed as a one-dimensional convex function of $t \geq 0$. By (146) and the symmetry of M ,

$$g'(0) = \langle \nabla_W \Psi(0, \mu_0 \mathbf{1}_n), W^\times \rangle = -a_0 \operatorname{tr}(M V^\top V U^\top U).$$

The quantity $\operatorname{tr}(MV^\top V U^\top U)$ is a degree-4 polynomial in the i.i.d. $\mathcal{N}(0, 1)$ entries of $(\sqrt{d}U, \sqrt{d}V)$, with mean $\operatorname{tr}(M) = n$ (by independence of U, V and $\mathbb{E}[U^\top U] = \mathbb{E}[V^\top V] = I_n$) and variance $O(n)$. Sub-exponential concentration via Hanson–Wright (or, equivalently, Gaussian hypercontractivity for finite-degree chaos) yields

$$\Omega_d^{(2)} := \{ g'(0) \leq -\frac{1}{2} a_0 n \}, \quad \mathbb{P}(\Omega_d^{(2)}) \geq 1 - d^{-D}, \quad (149)$$

for every $D > 0$ and all sufficiently large d .

Step 3: Curvature along the path. Computing the curvature $g''(t)$ for the smooth function $\Psi(\cdot, \mu_0 \mathbf{1}_n)$ and bounding $|\ell''| \leq 1/4$, $|\phi''_\beta| \leq \beta/4$:

$$g''(t) \leq \frac{1}{4} \sum_{i=1}^n (t'_i(t))^2 + \frac{\beta}{4k} \sum_{i=1}^n \sum_{j \neq i} (S_{ji}^\times)^2 + \frac{\lambda n}{d^2} \|W^\times\|_F^2,$$

where $t_i(t)$ is the loss argument at sample i along the path. Using $|t'_i(t)|^2 \leq 2(S_{ii}^\times)^2 + (2/(rk)) \sum_{j \neq i} (S_{ji}^\times)^2$, the first two terms collapse into

$$\frac{1}{2} \sum_i (S_{ii}^\times)^2 + \left(\frac{1}{2rk} + \frac{\beta}{4k} \right) \sum_{i,j \neq i} (S_{ji}^\times)^2. \quad (150)$$

The summed second moments of $S^\times$ admit the matching expectations

$$\mathbb{E} \sum_i (S_{ii}^\times)^2 \asymp n, \quad \mathbb{E} \sum_{i,j \neq i} (S_{ji}^\times)^2 \asymp n^2,$$

both inherited from the Wishart structure of $U^\top U$ and $V^\top V$. The ridge contribution $\|W^\times\|_F^2 = \|UV^\top\|_F^2$ has mean n (by the same calculation as (147) with $M = I$), so that $(\lambda n/d^2) \|W^\times\|_F^2 \asymp \lambda \alpha n$. Each of the three sums is a polynomial of bounded degree (at most

8) in the i.i.d. $\mathcal{N}(0, 1)$ entries of $\sqrt{d}U$ and $\sqrt{d}V$ with mean of order n^a for $a \in \{1, 2, 1\}$, respectively, and variance of strictly smaller order. Gaussian hypercontractivity for finite-degree chaos (or, equivalently, Latala's moment inequalities) yields stretched-exponential concentration $\mathbb{P}(|X - \mathbb{E}X| > \delta \mathbb{E}X) \leq \exp(-c d^{2/q})$ for any constant $\delta > 0$ and degree- q chaos. Combining the three concentration statements via a union bound, there is a constant $L = L(\alpha, r, \beta, \lambda) > 0$ such that, for every $D > 0$, the event

$$\Omega_d^{(3)} := \left\{ \sup_{t \in [0, 1]} g''(t) \leq L n \right\} \quad (151)$$

satisfies $\mathbb{P}(\Omega_d^{(3)}) \geq 1 - d^{-D}$ for all sufficiently large d . (The supremum over t is harmless: the data-dependent factors in (150) are independent of t , while the loss curvatures $|\ell''|, |\phi''_\beta|$ admit deterministic bounds $1/4, \beta/4$ that hold uniformly in t .)

Step 4: Objective gap. Set $t_\star := a_0/(2L) \wedge 1$. By Taylor's theorem with remainder,

$$g(t_\star) \leq g(0) + g'(0)t_\star + \frac{1}{2} \sup_{[0, t_\star]} g'' \cdot t_\star^2. \quad (152)$$

On $\Omega_d^{(2)} \cap \Omega_d^{(3)}$, the bounds (149) and (151) reduce (152) to

$$g(t_\star) \leq g(0) - \frac{1}{2} a_0 n t_\star + \frac{1}{2} L n t_\star^2 \leq g(0) - \frac{1}{4} a_0 n t_\star =: g(0) - c_1 n,$$

with $c_1 := a_0 t_\star/4 > 0$.

Step 5: Convex linearization and contradiction. By optimality of $(\widehat{W}, \widehat{\mu})$ in Ψ and convexity of $\Psi(\cdot, \mu_0 \mathbf{1}_n)$ in W ,

$$\begin{aligned} g(0) - c_1 n &\geq g(t_\star) \geq \Psi(\widehat{W}, \widehat{\mu}) \geq \Psi(0, \mu_0 \mathbf{1}_n) + \langle \nabla_W \Psi(0, \mu_0 \mathbf{1}_n), \widehat{W} \rangle \\ &\geq g(0) - \|\nabla_W \Psi(0, \mu_0 \mathbf{1}_n)\|_F \cdot \|\widehat{W}\|_F, \end{aligned}$$

where the third inequality follows from the convex subgradient inequality around $(0, \mu_0 \mathbf{1}_n)$, and the optimality $\nabla_\mu \Psi(0, \mu_0 \mathbf{1}_n) = 0$ at μ_0 (which follows from $f'(\mu_0) = 0$) eliminates the μ -component. Rearranging, on $\Omega_d := \bigcap_{j=1}^3 \Omega_d^{(j)}$,

$$\|\widehat{W}\|_F \geq \frac{c_1 n}{\|\nabla_W \Psi(0, \mu_0 \mathbf{1}_n)\|_F} \geq \frac{c_1 n}{a_0 K d} =: c_0 d,$$

using (148) and $n \asymp \alpha d^2$, with $c_0 := c_1 \alpha / (a_0 K) > 0$. This is (145).

A union bound over $\Omega_d^{(1)}, \Omega_d^{(2)}, \Omega_d^{(3)}$ gives $\mathbb{P}(\Omega_d) \leq 3d^{-D}$, which is at most $d^{-(D-1)}$ for $d \geq 3$; since D is arbitrary, the rate d^{-D} is preserved.

For the conclusion on ν_d^2 : combining Lemma 14 with the bound just established,

$$\|\widehat{W}\|_F^2/d^2 \geq c_0^2 \mathbf{1}_{\Omega_d}, \quad \|\widehat{W}\|_F^2/d^2 \leq C_0 \text{ everywhere,}$$

so $\nu_d^2 \geq c_0^2 \mathbb{P}(\Omega_d) \geq c_0^2(1 - d^{-D})$ for every $D > 0$. $\square$

Together, Lemma 14 and Lemma 15 certify that ν_d is bounded away from 0 and ∞ for all sufficiently large d . This is the *a priori* input that subsequent analyses—in particular the bilinear coupling Corollary 1—use implicitly. The Frobenius-norm concentration statement $\|\widehat{W}\|_F^2/d^2 - \nu_d^2 = \mathcal{O}_\prec(d^{-1/2})$ is not a separate postulate; it is folded into (A5) through the energy identity in Section 4.3.

E A coupling lemma for bilinear Gaussian forms

This appendix establishes a coupling estimate for the joint law of bilinear forms $u_j^\top W v_0$ that share a common right-side Gaussian vector v_0 . The estimate is the technical input behind the joint Gaussian limit for competitor scores derived in Section 4.3, where it is applied with $W = \widehat{W}^{\setminus i, j_1, \dots, j_m}$ and $v_0 = v_i$.

We first prove the estimate for a deterministic matrix W , then transfer it to the random case via concentration of $\|W\|_{\mathbb{F}}^2/d^2$.

E.1 Deterministic matrix

Lemma 16 (Bilinear Gaussian coupling, deterministic matrix). *Let $W \in \mathbb{R}^{d \times d}$ be a deterministic matrix with $V_0 := \|W\|_{\mathbb{F}}^2/d^2 > 0$, and let $u_1, \dots, u_m, v_0 \in \mathbb{R}^d$ be independent Gaussian vectors with $u_k, v_0 \sim \mathcal{N}(0, I_d/d)$. Set*

$$X_j := u_j^\top W v_0, \quad j = 1, \dots, m.$$

On a common probability space, there exist i.i.d. random variables $Y_1, \dots, Y_m \sim \mathcal{N}(0, V_0)$ such that

$$\|(X_1, \dots, X_m) - (Y_1, \dots, Y_m)\|_2 = \mathcal{O}_{\prec}(\|W^\top W\|_{\mathbb{F}}/(d^2 \sqrt{V_0})). \quad (153)$$

Remark 9 (Interpretation). *The deviation $\|W^\top W\|_{\mathbb{F}}/d^2$ measures the spread of the eigenvalues of $W^\top W/d^2$: when the eigenvalues are concentrated, $\|W^\top W\|_{\mathbb{F}}/d^2$ is small relative to $\|W\|_{\mathbb{F}}^2/d^2 = \text{tr}(W^\top W)/d^2$. In the proof below it is exactly the Hanson–Wright scale for the random conditional variance $\|W v_0\|^2/d$.*

Proof. The argument exploits the conditional Gaussian structure of $\{X_j\}$ given v_0 .

Step 1: Conditional Gaussian. Set $w := W v_0 \in \mathbb{R}^d$. Conditional on v_0 , w is deterministic, and $X_j = u_j^\top w$ is a Gaussian linear functional of u_j . Since $u_1, \dots, u_m \sim \mathcal{N}(0, I_d/d)$ are independent across j ,

$$(X_1, \dots, X_m) \mid v_0 \sim \mathcal{N}(0, V \cdot I_m), \quad V := \frac{\|W v_0\|^2}{d}.$$

Step 2: Hanson–Wright on the conditional variance. We have

$$\mathbb{E}[V] = \frac{1}{d} \mathbb{E}[v_0^\top W^\top W v_0] = \frac{1}{d^2} \text{tr}(W^\top W) = \frac{\|W\|_{\mathbb{F}}^2}{d^2} =: V_0,$$

using $\mathbb{E}[v_0 v_0^\top] = I_d/d$. The Hanson–Wright inequality applied to the centered quadratic form $V - V_0 = (1/d)(v_0^\top W^\top W v_0 - \|W\|_{\mathbb{F}}^2/d)$ in the Gaussian vector v_0 gives

$$V - V_0 = \mathcal{O}_{\prec}(\|W^\top W\|_{\mathbb{F}}/d^2), \quad \text{Var } V = \frac{2}{d^4} \|W^\top W\|_{\mathbb{F}}^2. \quad (154)$$

Step 3: Coupling. Define $Y_j := X_j \cdot \sqrt{V_0/V}$ for each j . Conditional on v_0 , $\{Y_j\}$ inherit the Gaussianity and independence of $\{X_j\}$, with conditional variance $V \cdot (V_0/V) = V_0$. Since the conditional law $\mathcal{N}(0, V_0 \cdot I_m)$ is independent of v_0 , the unconditional joint law of $(Y_1, \dots, Y_m)$ is i.i.d. $\mathcal{N}(0, V_0)$ as required.

The deviation factors as

$$X_j - Y_j = X_j(1 - \sqrt{V_0/V}) = X_j \cdot \frac{\sqrt{V} - \sqrt{V_0}}{\sqrt{V}}.$$

Two facts: conditional on v_0 , the standardized form $X_j/\sqrt{V} \sim \mathcal{N}(0, 1)$, so $|X_j/\sqrt{V}| = \mathcal{O}_{\prec}(1)$ uniformly in j ; and $|\sqrt{V} - \sqrt{V_0}| \leq |V - V_0|/\sqrt{V_0}$, by $\sqrt{V} + \sqrt{V_0} \geq \sqrt{V_0}$. Combined with (154),

$$|X_j - Y_j| = |X_j/\sqrt{V}| \cdot |\sqrt{V} - \sqrt{V_0}| = \mathcal{O}_{\prec}(\|W^\top W\|_{\text{F}}/(d^2 \sqrt{V_0})),$$

uniformly in j . For fixed m , we have

$$\|(X) - (Y)\|_2^2 = \sum_{j=1}^m |X_j - Y_j|^2 = \mathcal{O}_{\prec}((\|W^\top W\|_{\text{F}}/(d^2 \sqrt{V_0}))^2),$$

which gives (153). □

E.2 Random matrix with concentrated Frobenius norm

Corollary 1 (Bilinear Gaussian coupling, random matrix). *Let W be a random $d \times d$ matrix with $\|W\|_{\text{F}} > 0$, independent of $u_1, \dots, u_m, v_0$ which are i.i.d. $\mathcal{N}(0, I_d/d)$. Let $\nu_d^2 > 0$ be a deterministic constant. Suppose $\rho_f, \rho_{\text{op}} > 0$ are deterministic rates such that*

$$\frac{\|W\|_{\text{F}}^2}{d^2} - \nu_d^2 = \mathcal{O}_{\prec}(\rho_f), \quad \frac{\|W^\top W\|_{\text{F}}}{d^2} = \mathcal{O}_{\prec}(\rho_{\text{op}}). \quad (155)$$

Set $X_j := u_j^\top W v_0$ for $j = 1, \dots, m$. On a common probability space, there exist i.i.d. random variables $Y_1, \dots, Y_m \sim \mathcal{N}(0, \nu_d^2)$ such that

$$\|(X_1, \dots, X_m) - (Y_1, \dots, Y_m)\|_2 = \mathcal{O}_{\prec}((\rho_{\text{op}} + \rho_f)/\nu_d). \quad (156)$$

Proof. The key observation is that, conditional on (W, v_0) , $X_j/\sqrt{V} \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$ across j , where $V := \|W v_0\|^2/d$ is the conditional variance.

Define

$$Y_j := X_j \sqrt{\nu_d^2/V}, \quad j = 1, \dots, m.$$

Conditional on (W, v_0) , $\{Y_j\}$ inherit the Gaussianity and independence of $\{X_j\}$ with variance $V \cdot \nu_d^2/V = \nu_d^2$, so the unconditional joint law of $(Y_1, \dots, Y_m)$ is i.i.d. $\mathcal{N}(0, \nu_d^2)$ as required. The deviation factors as $X_j - Y_j = X_j(1 - \sqrt{\nu_d^2/V}) = X_j(\sqrt{V} - \sqrt{\nu_d^2})/\sqrt{V}$, and using $|X_j/\sqrt{V}| = \mathcal{O}_{\prec}(1)$ together with $|\sqrt{V} - \sqrt{\nu_d^2}| \leq |V - \nu_d^2|/\sqrt{\nu_d^2}$,

$$|X_j - Y_j| = |X_j/\sqrt{V}| \cdot |\sqrt{V} - \sqrt{\nu_d^2}| = \mathcal{O}_{\prec}(|V - \nu_d^2|/\sqrt{\nu_d^2}). \quad (157)$$

Decompose

$$V - \nu_d^2 = (V - V_0(W)) + (V_0(W) - \nu_d^2), \quad V_0(W) := \|W\|_{\mathbb{F}}^2/d^2.$$

Hanson–Wright applied to $V - V_0(W)$ conditional on W gives

$$V - V_0(W) = \mathcal{O}_{\prec}\left(\|W^\top W\|_{\mathbb{F}}/d^2\right) = \mathcal{O}_{\prec}(\rho_{\text{op}}),$$

using (154). The second piece is $\mathcal{O}_{\prec}(\rho_f)$ by (155). Combining with (157) and summing over j yields (156). $\square$

F A trace estimate for the inverse Hessian

This appendix records several elementary estimates that will be used in the characterization of the susceptibility parameter χ . The first says that a fresh rank-one Gaussian direction probes the normalized trace of a bounded linear operator on $\mathbb{R}^{d \times d}$. The next two show that the normalized inverse trace is stable under the perturbations needed to simplify the TAM Hessian.

We will use the following moment notation. For a sequence of random variables X_d and deterministic positive numbers a_d , we write

$$X_d = \mathcal{O}_{\mathcal{L}}(a_d)$$

if, for every fixed $p < \infty$, there is a constant $C_p < \infty$ such that $\|X_d\|_{L^p} \leq C_p a_d$ for all sufficiently large d .

Lemma 17 (Fresh rank-one trace estimate). *Let $u, v \in \mathbb{R}^d$ be independent Gaussian vectors with distribution $\mathcal{N}(0, I_d/d)$. Let $T : \mathbb{R}^{d \times d} \rightarrow \mathbb{R}^{d \times d}$ be a random symmetric linear operator, independent of (u, v) , and assume that*

$$\|T\|_{\text{op}} = \mathcal{O}_{\mathcal{L}}(1).$$

Then

$$\left\langle uv^\top, T[uv^\top] \right\rangle = \frac{\text{tr}(T)}{d^2} + \mathcal{O}_{\mathcal{L}}(d^{-1/2}).$$

Equivalently, for every fixed $p < \infty$,

$$\left\| \left\langle uv^\top, T[uv^\top] \right\rangle - \frac{\text{tr}(T)}{d^2} \right\|_{L^p} \leq \frac{C_p}{\sqrt{d}}. \quad (158)$$

Proof. Write $u = g/\sqrt{d}$ and $v = h/\sqrt{d}$, where $g, h \in \mathbb{R}^d$ are independent standard Gaussian vectors. Define

$$F(g, h, T) := \left\langle uv^\top, T[uv^\top] \right\rangle = \frac{1}{d^2} \langle g \otimes h, T[g \otimes h] \rangle.$$

Since T is independent of (g, h) , we have

$$\mathbb{E}[F(g, h, T) \mid T] = \frac{\text{tr}(T)}{d^2}.$$

We first bound the fluctuation in L^2 . By the Gaussian Poincare inequality applied to the variables (g, h) , with T kept fixed,

$$\text{Var}(F(g, h, T) \mid T) \leq \mathbb{E}[\|\nabla_g F\|^2 + \|\nabla_h F\|^2 \mid T]. \quad (159)$$

For fixed h , define the compressed operator

$$T_h := (I \otimes h)^* T (I \otimes h) \quad \text{on } \mathbb{R}^d,$$

where $(I \otimes h)g = g \otimes h$. Then

$$F(g, h, T) = \frac{1}{d^2} g^\top T_h g, \quad \nabla_g F = \frac{2}{d^2} T_h g,$$

using the symmetry of T . It follows that

$$\mathbb{E}[\|\nabla_g F\|^2 \mid T] = \frac{4}{d^4} \mathbb{E}[\text{tr}(T_h^2) \mid T]. \quad (160)$$

Moreover,

$$\|T_h\|_{\text{op}} \leq \|T\|_{\text{op}} \|h\|^2,$$

and hence

$$\text{tr}(T_h^2) \leq d \|T_h\|_{\text{op}}^2 \leq d \|T\|_{\text{op}}^2 \|h\|^4.$$

Since $\mathbb{E}\|h\|^4 = d^2 + 2d$, (160) gives

$$\mathbb{E}[\|\nabla_g F\|^2 \mid T] \leq \frac{C \|T\|_{\text{op}}^2}{d}. \quad (161)$$

The same argument, with the roles of g and h interchanged, gives

$$\mathbb{E}[\|\nabla_h F\|^2 \mid T] \leq \frac{C \|T\|_{\text{op}}^2}{d}. \quad (162)$$

Combining (159), (161), and (162), we obtain

$$\text{Var}(F(g, h, T) \mid T) \leq \frac{C \|T\|_{\text{op}}^2}{d}. \quad (163)$$

We now upgrade this L^2 bound to fixed moments. The centered random variable

$$F(g, h, T) - \frac{\text{tr}(T)}{d^2}$$

is, for fixed T , a Gaussian polynomial of degree at most 4 in (g, h) . By Gaussian hypercontractivity, for every fixed $p < \infty$,

$$\left\| F(g, h, T) - \frac{\text{tr}(T)}{d^2} \right\|_{L^p(g, h \mid T)} \leq C_p \left\| F(g, h, T) - \frac{\text{tr}(T)}{d^2} \right\|_{L^2(g, h \mid T)} \leq \frac{C_p \|T\|_{\text{op}}}{\sqrt{d}},$$

where the last step uses (163). Taking expectation over T and using $\|T\|_{\text{op}} = O_{\mathcal{L}}(1)$ yields (158). This proves the lemma. $\square$

We next record two deterministic perturbation estimates for the Hessian operators in (26)–(28). We identify $\mathbb{R}^{d \times d}$ with $\mathbb{R}^{d^2}$ throughout this discussion. If $A, B : \mathbb{R}^{d \times d} \rightarrow \mathbb{R}^{d \times d}$ are positive-definite self-adjoint operators, equivalently $d^2 \times d^2$ symmetric matrices after vectorization, and $E := A - B$, then the resolvent identity gives the following bound: whenever $\|A^{-1}\|_{\text{op}}, \|B^{-1}\|_{\text{op}} \leq \tau^{-1}$,

$$\left| \frac{1}{d^2} \text{tr}(A^{-1}) - \frac{1}{d^2} \text{tr}(B^{-1}) \right| \leq \frac{1}{\tau^2 d^2} \|E\|_* = \frac{1}{\tau^2 d^2} \|A - B\|_*, \quad (164)$$

where $\|\cdot\|_*$ denotes the nuclear norm on this d^2 -dimensional operator space.

For a fixed cavity index h , let $H^{\setminus h}$ be the Hessian in (26). Define the simplified Hessian $\tilde{H}^{\setminus h}$ by replacing the block $M_i^{\setminus h}$ in (28), for every $i \neq h$, by

$$\tilde{M}_i^{\setminus h} := b_i^{\setminus h} e_i e_i^\top + a_i^{\setminus h} \text{diag}(\tilde{R}_{:,i}^{\setminus h}),$$

and keeping the h -th block equal to zero. Equivalently, $\tilde{H}^{\setminus h}$ is obtained from $H^{\setminus h}$ by replacing $b_i^{\setminus h} Q_{:,i}^{\setminus h} (Q_{:,i}^{\setminus h})^\top$ by $b_i^{\setminus h} e_i e_i^\top$ in each active block.

Lemma 18 (Trace reduction to the simplified Hessian). *Assume $\lambda n/d^2 \geq \tau > 0$ and $\max_i b_i^{\setminus h} \leq B$. For $i \neq h$, write*

$$Q_{:,i}^{\setminus h} = -e_i + \rho_i^{\setminus h}, \quad p_i^{\setminus h} := U \rho_i^{\setminus h}.$$

Then

$$\left| \frac{1}{d^2} \text{tr}((H^{\setminus h})^{-1}) - \frac{1}{d^2} \text{tr}((\tilde{H}^{\setminus h})^{-1}) \right| \leq \frac{B}{\tau^2 d^2} \sum_{i \neq h} \|v_i\|^2 \left(2\|u_i\| \|p_i^{\setminus h}\| + \|p_i^{\setminus h}\|^2 \right). \quad (165)$$

Consequently, if $n/d^2 = O(1)$, $\max_i \{\|u_i\|, \|v_i\|\} = O(1)$, and $\max_{i \neq h} \|p_i^{\setminus h}\| \leq \eta$, then the normalized inverse trace of $H^{\setminus h}$ differs from that of the simplified Hessian $\tilde{H}^{\setminus h}$ by $O(\eta)$.

Proof. Since

$$Q_{:,i}^{\setminus h} (Q_{:,i}^{\setminus h})^\top - e_i e_i^\top = -e_i (\rho_i^{\setminus h})^\top - \rho_i^{\setminus h} e_i^\top + \rho_i^{\setminus h} (\rho_i^{\setminus h})^\top,$$

the i th summand in $H^{\setminus h} - \tilde{H}^{\setminus h}$ is

$$\Delta \mapsto b_i^{\setminus h} \left[-u_i (p_i^{\setminus h})^\top - p_i^{\setminus h} u_i^\top + p_i^{\setminus h} (p_i^{\setminus h})^\top \right] \Delta v_i v_i^\top.$$

Each of the three displayed terms is rank one as an operator on $\mathbb{R}^{d \times d}$. For example,

$$u_i (p_i^{\setminus h})^\top \Delta v_i v_i^\top = \langle p_i^{\setminus h} v_i^\top, \Delta \rangle u_i v_i^\top,$$

and hence its nuclear norm is $\|u_i v_i^\top\|_{\text{F}} \|p_i^{\setminus h} v_i^\top\|_{\text{F}} = \|u_i\| \|p_i^{\setminus h}\| \|v_i\|^2$. The same bound holds for the transposed cross term, while the $p_i^{\setminus h} (p_i^{\setminus h})^\top$ term contributes $\|p_i^{\setminus h}\|^2 \|v_i\|^2$. Thus

$$\|H^{\setminus h} - \tilde{H}^{\setminus h}\|_* \leq B \sum_{i \neq h} \|v_i\|^2 \left(2\|u_i\| \|p_i^{\setminus h}\| + \|p_i^{\setminus h}\|^2 \right).$$

Both $H^{\setminus h}$ and $\tilde{H}^{\setminus h}$ are bounded below by the ridge term $(\lambda n/d^2)I$, so their inverse operator norms are at most τ^{-1} . Applying (164) gives (165). $\square$

Lemma 19 (Stability at the natural coefficient scale). *Let $\tilde{H}^{\setminus h}$ be the simplified Hessian above. Let $\bar{H}^{\setminus h}$ be the simplified Hessian obtained by replacing, for $i \neq h$, the coefficients $a_i^{\setminus h}$ and $\tilde{R}_{j,i}^{\setminus h}$ by $\bar{a}_i$ and $\bar{R}_{j,i}$, while keeping the coefficients $b_i^{\setminus h}$ unchanged. Assume*

$$\frac{\lambda n}{d^2} \geq \tau > 0, \quad 0 \leq a_i^{\setminus h}, \bar{a}_i \leq B, \quad \bar{R}_{j,i} \geq 0,$$

and

$$\max_{i \neq h} |\bar{a}_i - a_i^{\setminus h}| \leq \delta_a, \quad \max_{i \neq h} \sum_{j \neq i, h} \tilde{R}_{j,i}^{\setminus h} \leq C_R, \quad \max_{j \neq i, h} |\bar{R}_{j,i} - \tilde{R}_{j,i}^{\setminus h}| \leq \frac{\delta_R}{k}.$$

Then

$$\begin{aligned} & \left| \frac{1}{d^2} \text{tr}((\bar{H}^{\setminus h})^{-1}) - \frac{1}{d^2} \text{tr}((\tilde{H}^{\setminus h})^{-1}) \right| \\ & \leq \frac{1}{\tau^2 d^2} \sum_{i \neq h} \|v_i\|^2 \sum_{j \neq i, h} \left(\delta_a \tilde{R}_{j,i}^{\setminus h} + \frac{B \delta_R}{k} \right) \|u_j\|^2. \end{aligned} \quad (166)$$

Consequently, if $n/d^2 = O(1)$, $n/k = O(1)$, $\max_i \{\|u_i\|, \|v_i\|\} = O(1)$, and $C_R = O(1)$, then the normalized inverse trace changes by

$$O(\delta_a + \delta_R).$$

Proof. Each elementary map

$$\Delta \mapsto u_j u_j^\top \Delta v_i v_i^\top = \langle u_j v_i^\top, \Delta \rangle u_j v_i^\top$$

has rank one and nuclear norm $\|u_j\|^2 \|v_i\|^2$. Therefore

$$\|\bar{H}^{\setminus h} - \tilde{H}^{\setminus h}\|_* \leq \sum_{i \neq h} \sum_{j \neq i, h} |\bar{a}_i \bar{R}_{j,i} - a_i^{\setminus h} \tilde{R}_{j,i}^{\setminus h}| \|u_j\|^2 \|v_i\|^2. \quad (167)$$

The coefficient difference satisfies

$$|\bar{a}_i \bar{R}_{j,i} - a_i^{\setminus h} \tilde{R}_{j,i}^{\setminus h}| \leq |\bar{a}_i - a_i^{\setminus h}| \tilde{R}_{j,i}^{\setminus h} + \bar{a}_i |\bar{R}_{j,i} - \tilde{R}_{j,i}^{\setminus h}| \leq \delta_a \tilde{R}_{j,i}^{\setminus h} + \frac{B \delta_R}{k}.$$

Substituting this into (167), and using that both Hessians are bounded below by the ridge term $(\lambda n/d^2)I$, (164) proves (166). $\square$

Quadratic forms of the simplified Hessian

We now use the simplified Hessian to extract the scalar equation for χ . In this subsection we suppress the leave-one-out superscript and write

$$f_{ji} := \text{vec}(u_j v_i^\top), \quad r_{j,i} := \beta q_{j,i}(1 - q_{j,i}), \quad \tilde{R}_{j,i} = \frac{r_{j,i}}{k}.$$

The full simplified Hessian is

$$\tilde{H} = \frac{\lambda n}{d^2} I + \sum_{i=1}^n b_i f_{ii} f_{ii}^\top + \sum_{i=1}^n \sum_{j \neq i} \frac{a_i r_{j,i}}{k} f_{ji} f_{ji}^\top.$$

Here $a_i, b_i, q_{j,i}$ are evaluated at the full minimizer. For the i -cavity problem, superscript (i) denotes the corresponding quantities after removing sample i from both tensor legs.

Feature spectral bounds. We use the following spectral estimates for the feature matrices: uniformly in i and ℓ ,

$$\left\| [f_{\ell\ell}]_{\ell \neq i} \right\|_{\text{op}} = \mathcal{O}_{\prec}(1), \quad \left\| [f_{j\ell}]_{j \neq \ell, i} \right\|_{\text{op}} = \mathcal{O}_{\prec}(\sqrt{d}). \quad (168)$$

Lemma 20 (LOO coefficient stability). *Assume (A1)–(A3). Then, uniformly in i ,*

$$\max_{\ell \neq i} |b_{\ell} - b_{\ell}^{(i)}| = \mathcal{O}_{\prec}(d^{-1}), \quad (169)$$

and

$$\max_{\ell \neq i} \max_{j \neq \ell, i} \left| a_{\ell} r_{j,\ell} - a_{\ell}^{(i)} r_{j,\ell}^{(i)} \right| = \mathcal{O}_{\prec}(d^{-1}). \quad (170)$$

Proof. The functions $a = \sigma(-m)$, $b = a(1 - a)$, and $r = \beta q(1 - q)$ are uniformly Lipschitz on the bounded-score event supplied by (A3); the corresponding LOO scores are bounded by (A2). The LOO perturbations of the relevant scores and thresholds are $\mathcal{O}_{\prec}(d^{-1})$ by (A1)–(A2), since $n \asymp d^2$ and the μ perturbation is smaller. Applying the Lipschitz bounds entrywise gives (169)–(170). $\square$

Lemma 21 (Diagonal quadratic form). *Assume (A1)–(A6) and the feature spectral bounds (168). Let χ_d be as in (48). Then, uniformly in i ,*

$$f_{ii}^{\top} \tilde{H}^{-1} f_{ii} = \frac{\chi_d}{1 + b_i \chi_d} + \mathcal{O}_{\prec}(d^{-1/2}). \quad (171)$$

Proof. Let H_i be the simplified i -cavity Hessian that keeps only feature directions $f_{j\ell}$ with $j, \ell \neq i$, and uses the i -cavity coefficients:

$$H_i = \frac{\lambda n}{d^2} I + \sum_{\ell \neq i} b_{\ell}^{(i)} f_{\ell\ell} f_{\ell\ell}^{\top} + \sum_{\ell \neq i} \sum_{j \neq \ell, i} \frac{a_{\ell}^{(i)} r_{j,\ell}^{(i)}}{k} f_{j\ell} f_{j\ell}^{\top}.$$

This operator is independent of (u_i, v_i) . Set

$$A_i := H_i + b_i f_{ii} f_{ii}^{\top}.$$

We first compare $\tilde{H}$ with A_i . Write

$$\tilde{H} = A_i + C_i^{\text{feat}} + C_i^{\text{coef}},$$

where

$$C_i^{\text{feat}} = \sum_{\ell \neq i} \frac{a_{\ell} r_{i,\ell}}{k} f_{i\ell} f_{i\ell}^{\top} + \sum_{j \neq i} \frac{a_i r_{j,i}}{k} f_{ji} f_{ji}^{\top}$$

contains the feature directions involving u_i or v_i , except for the diagonal rank-one term, and

$$\begin{aligned} C_i^{\text{coef}} &= \sum_{\ell \neq i} (b_{\ell} - b_{\ell}^{(i)}) f_{\ell\ell} f_{\ell\ell}^{\top} \\ &\quad + \sum_{\ell \neq i} \sum_{j \neq \ell, i} \frac{a_{\ell} r_{j,\ell} - a_{\ell}^{(i)} r_{j,\ell}^{(i)}}{k} f_{j\ell} f_{j\ell}^{\top}. \end{aligned}$$

The resolvent identity gives

$$f_{ii}^\top \tilde{H}^{-1} f_{ii} - f_{ii}^\top A_i^{-1} f_{ii} = -f_{ii}^\top \tilde{H}^{-1} (C_i^{\text{feat}} + C_i^{\text{coef}}) A_i^{-1} f_{ii}.$$

By Woodbury,

$$A_i^{-1} f_{ii} = \frac{H_i^{-1} f_{ii}}{1 + b_i f_{ii}^\top H_i^{-1} f_{ii}},$$

and the denominator is at least one. Thus it suffices to bound

$$f_{ii}^\top \tilde{H}^{-1} (C_i^{\text{feat}} + C_i^{\text{coef}}) H_i^{-1} f_{ii}. \quad (172)$$

Let $\xi_i := \tilde{H}^{-1} f_{ii}$. The ridge lower bound gives $\|\xi_i\| = \mathcal{O}_\prec(1)$. For the two terms in C_i^{feat} , each summand has a factor of the form

$$f_{i\ell}^\top H_i^{-1} f_{ii} \quad \text{or} \quad f_{ji}^\top H_i^{-1} f_{ii}.$$

In the first expression v_i is an unmatched Gaussian vector, and in the second expression u_i is unmatched. Equivalently, if $M = \text{mat}(H_i^{-1} f_{i\ell})$ or $M = \text{mat}(H_i^{-1} f_{ji})$, then $\|M\|_F = \mathcal{O}_\prec(1)$ and the expression is $u_i^\top M v_i$ with one Gaussian leg independent of M and the other leg. Hence each such factor is $\mathcal{O}_\prec(d^{-1/2})$. Since the coefficients have total scale $n/k = O(1)$, the contribution of C_i^{feat} to (172) is $\mathcal{O}_\prec(d^{-1/2})$.

We next consider C_i^{coef} . The diagonal coefficient part can be written as

$$[f_{\ell\ell}]_{\ell \neq i} \text{diag}(b_\ell - b_\ell^{(i)})_{\ell \neq i} [f_{\ell\ell}]_{\ell \neq i}^\top.$$

By Lemma 20 and (168), its operator norm is $\mathcal{O}_\prec(d^{-1})$, so its contribution to (172) is $\mathcal{O}_\prec(d^{-1})$.

For the off-diagonal coefficient part, fix $\ell \neq i$ and write

$$F_\ell^{(i)} := [f_{j\ell}]_{j \neq \ell, i}, \quad D_\ell^{(i)} := \text{diag}\left(a_\ell r_{j,\ell} - a_\ell^{(i)} r_{j,\ell}^{(i)}\right)_{j \neq \ell, i}.$$

The corresponding block is $k^{-1} F_\ell^{(i)} D_\ell^{(i)} (F_\ell^{(i)})^\top$. Lemma 20 gives $\|D_\ell^{(i)}\|_{\text{op}} = \mathcal{O}_\prec(d^{-1})$, while (168) gives $\|F_\ell^{(i)}\|_{\text{op}} = \mathcal{O}_\prec(\sqrt{d})$. Moreover,

$$\left\| (F_\ell^{(i)})^\top H_i^{-1} f_{ii} \right\| = \mathcal{O}_\prec(1),$$

because H_i and $F_\ell^{(i)}$ are independent of f_{ii} and

$$\frac{1}{d^2} \text{tr}\left(H_i^{-1} F_\ell^{(i)} (F_\ell^{(i)})^\top H_i^{-1}\right) = \mathcal{O}_\prec(1).$$

Therefore the fixed- ℓ contribution is $\mathcal{O}_\prec(1/(k\sqrt{d}))$, and summing over ℓ gives $\mathcal{O}_\prec(d^{-1/2})$. We have proved

$$f_{ii}^\top \tilde{H}^{-1} f_{ii} = f_{ii}^\top A_i^{-1} f_{ii} + \mathcal{O}_\prec(d^{-1/2}). \quad (173)$$

It remains to evaluate the right-hand side. Woodbury gives

$$f_{ii}^\top A_i^{-1} f_{ii} = \frac{x_i}{1 + b_i x_i}, \quad x_i := f_{ii}^\top H_i^{-1} f_{ii}.$$

Since H_i is independent of f_{ii} , Lemma 17, (A6), and the finite-removal trace stability transferred to the simplified Hessian by the perturbation estimates above give

$$x_i = \chi_d + \mathcal{O}_{\prec}(d^{-1/2}).$$

The map $x \mapsto x/(1 + b_i x)$ is uniformly Lipschitz for $b_i \geq 0$, and (171) follows from (173). $\square$

Lemma 22 (Off-diagonal quadratic form). *Under the assumptions of Lemma 21, uniformly for $j \neq i$,*

$$f_{ji}^\top \tilde{H}^{-1} f_{ji} = \chi_d + \mathcal{O}_{\prec}(d^{-1/2}).$$

Proof. The proof is the same as that of Lemma 21. One removes all terms involving u_j or v_i to create a cavity independent of f_{ji} . The direct Woodbury correction along f_{ji} has coefficient $a_i r_{j,i}/k = O(k^{-1})$ and is therefore absorbed into the $\mathcal{O}_{\prec}(d^{-1/2})$ error. The feature-removal and coefficient-removal errors are controlled by the same unmatched-Gaussian and feature-matrix estimates used above. $\square$

Remark 10 (Use for χ). *In the leave-one-out analysis, the operator T will be a Hessian inverse, for example $T = (H^{\setminus i})^{-1}$, evaluated on a sigma-field that is independent of a fresh pair (u_i, v_i) . Lemma 17 then gives*

$$\langle u_i v_i^\top, (H^{\setminus i})^{-1} [u_i v_i^\top] \rangle = \frac{1}{d^2} \text{tr}((H^{\setminus i})^{-1}) + O_{\mathcal{L}}(d^{-1/2}),$$

provided $\|(H^{\setminus i})^{-1}\|_{\text{op}} = O_{\mathcal{L}}(1)$. Thus the susceptibility can be characterized through the normalized trace of the inverse cavity Hessian. Lemmas 18 and 19 then justify replacing the original Hessian by simpler approximations without changing this normalized trace at leading order.

G Variational formulation of the scalar equations

This appendix verifies the variational representation used in Section 4.3. Throughout, Z and G denote independent standard normal variables, and $\nu, \chi > 0$.

G.1 The scalar TAM functional

Recall

$$c_r(\nu) = \min_{\mu \in \mathbb{R}} \left\{ \mu + \frac{1}{r} \mathbb{E}_Z [\phi_\beta(\nu Z - \mu)] \right\}.$$

The minimizer, denoted $\mu_r(\nu)$, is characterized by

$$r = \mathbb{E}_Z [\sigma(\beta(\nu Z - \mu_r(\nu)))].$$

$$c'_r(\nu) = \frac{1}{r} \mathbb{E}_Z [Z \phi'_\beta(\nu Z - \mu_r(\nu))].$$

Stein's identity gives

$$c'_r(\nu) = \frac{\nu}{r} \mathbb{E}_Z [\phi''_\beta(\nu Z - \mu_r(\nu))]. \quad (174)$$

We will also use that c_r is convex in ν . Indeed, the function

$$(\nu, \mu) \mapsto \mu + \frac{1}{r} \mathbb{E}_Z [\phi_\beta(\nu Z - \mu)]$$

is jointly convex, and partial minimization over μ preserves convexity.

G.2 Moreau envelope identities

Let $S = S_{\nu, \chi}(G)$ be the minimizer in (41), and set

$$A := \sigma(-S + c_r(\nu)), \quad B := A(1 - A).$$

The first-order condition in s is

$$S = \nu G + \chi A. \tag{175}$$

Envelope differentiation in χ gives

$$\partial_\chi M_r(\nu, \chi) = -\frac{1}{2} \mathbb{E}[A^2]. \tag{176}$$

For the derivative in ν , the envelope theorem gives

$$\partial_\nu M_r(\nu, \chi) = -\mathbb{E}[AG] + \mathbb{E}[A]c'_r(\nu). \tag{177}$$

Differentiating (175) with respect to the standard Gaussian coordinate G yields

$$\frac{dA}{dG} = -\frac{\nu B}{1 + \chi B}.$$

Therefore Stein's identity gives

$$\mathbb{E}[AG] = -\nu \mathbb{E}\left[\frac{B}{1 + \chi B}\right]. \tag{178}$$

Combining (174), (177), and (178),

$$\partial_\nu M_r(\nu, \chi) = \nu \mathbb{E}\left[\frac{B}{1 + \chi B}\right] + \frac{\nu}{r} \mathbb{E}[A] \mathbb{E}_Z [\phi''_\beta(\nu Z - \mu_r(\nu))]. \tag{179}$$

G.3 Saddle-point equations

The saddle potential is

$$\mathcal{P}_{\alpha, r}(\nu, \chi) = \frac{\lambda}{2} \nu^2 - \frac{\nu^2}{2\alpha\chi} + M_r(\nu, \chi).$$

Using (176),

$$\partial_\chi \mathcal{P}_{\alpha, r}(\nu, \chi) = \frac{\nu^2}{2\alpha\chi^2} - \frac{1}{2} \mathbb{E}[A^2]. \tag{180}$$

Hence the χ saddle-point equation is

$$\nu^2 = \alpha\chi^2 \mathbb{E}[A^2]. \tag{181}$$

Using (179),

$$\partial_\nu \mathcal{P}_{\alpha,r}(\nu, \chi) = \lambda\nu - \frac{\nu}{\alpha\chi} + \nu \mathbb{E} \left[\frac{B}{1 + \chi B} \right] + \frac{\nu}{r} \mathbb{E}[A] \mathbb{E}_Z \left[\phi''_\beta(\nu Z - \mu_r(\nu)) \right].$$

Since $\nu > 0$, the ν saddle-point equation is

$$\frac{1}{\alpha} = \lambda\chi + \chi \mathbb{E} \left[\frac{B}{1 + \chi B} \right] + \frac{\chi}{r} \mathbb{E}[A] \mathbb{E}_Z \left[\phi''_\beta(\nu Z - \mu_r(\nu)) \right]. \quad (182)$$

This is exactly the χ -equation obtained from the zero-residual limit of (67).

It remains to connect (181) with the variance equation. From (175),

$$\mathbb{E}[AS] = \nu \mathbb{E}[AG] + \chi \mathbb{E}[A^2].$$

Using (178) and (181),

$$\mathbb{E}[AS] = -\nu^2 \mathbb{E} \left[\frac{B}{1 + \chi B} \right] + \frac{\nu^2}{\alpha\chi}.$$

Substituting (182) into the last display gives

$$\mathbb{E}[AS] = \nu^2 \left[\lambda + \frac{1}{r} \mathbb{E}[A] \mathbb{E}_Z \left[\phi''_\beta(\nu Z - \mu_r(\nu)) \right] \right],$$

which is the corresponding ν -equation.

G.4 Maximization in χ and convexity in ν

For fixed ν , the inner problem is concave after the change of variables $t = 1/\chi$. Indeed,

$$\mathcal{P}_{\alpha,r}(\nu, 1/t) = \frac{\lambda}{2} \nu^2 - \frac{\nu^2}{2\alpha} t + \mathbb{E} \min_{s \in \mathbb{R}} \left\{ \frac{t}{2} (s - \nu G)^2 + \ell(s - c_r(\nu)) \right\}.$$

The last term is the infimum of affine functions of t , and hence is concave in t . Thus the inner extremization may equivalently be viewed as a one-dimensional concave maximization in $1/\chi$.

For fixed ν , differentiating $A = \sigma(-S + c_r(\nu))$ with respect to χ gives

$$\partial_\chi A = -\frac{AB}{1 + \chi B}.$$

Consequently,

$$\frac{d}{d\chi} (\chi^2 \mathbb{E}[A^2]) = 2\chi \mathbb{E} \left[\frac{A^2}{1 + \chi B} \right] > 0.$$

Moreover $\chi^2 \mathbb{E}[A^2] \rightarrow 0$ as $\chi \downarrow 0$, while $\chi^2 \mathbb{E}[A^2] \rightarrow \infty$ as $\chi \rightarrow \infty$. By (180), $\partial_\chi \mathcal{P}$ changes sign exactly once, from positive to negative. Hence $\chi \mapsto \mathcal{P}_{\alpha,r}(\nu, \chi)$ has a unique global maximizer.

Finally we prove strong convexity of the reduced potential $\mathcal{V}_{\alpha,r}$ defined in (42). The logistic loss has the Fenchel representation

$$\ell(t) = \sup_{a \in [0,1]} \{-at - I(a)\}, \quad I(a) := a \log a + (1-a) \log(1-a), \quad (183)$$

with the convention $0 \log 0 = 0$. Applying this pointwise in the Moreau envelope and minimizing over s gives

$$M_r(\nu, \chi) = \sup_{a(\cdot) \in [0,1]} \mathbb{E} \left[a(G)(c_r(\nu) - \nu G) - I(a(G)) - \frac{\chi}{2} a(G)^2 \right]. \quad (184)$$

Taking the supremum over $\chi > 0$ yields

$$\mathcal{V}_{\alpha,r}(\nu) = \frac{\lambda}{2} \nu^2 + \sup_{a(\cdot) \in [0,1]} \left\{ \mathbb{E} \left[a(G)(c_r(\nu) - \nu G) - I(a(G)) \right] - \frac{\nu}{\sqrt{\alpha}} (\mathbb{E}[a(G)^2])^{1/2} \right\}. \quad (185)$$

For each fixed measurable $a(\cdot)$, the expression inside the supremum is a convex function of ν , because c_r is convex and $\mathbb{E}[a(G)] \geq 0$; adding $\lambda \nu^2/2$ makes it λ -strongly convex. The supremum of λ -strongly convex functions is λ -strongly convex. This proves the convexity and uniqueness assertions in Proposition 6.

Remark 11 (Meaning of the dual optimizer). *The optimizer $a(G)$ in (184) is the scalar limit of the sample-wise logistic weights. Indeed, equality in the Fenchel representation (183) is attained at*

$$a = \sigma(-t).$$

Taking $t = S - c_r(\nu)$, where $S = S_{\nu,\chi}(G)$ is the proximal minimizer, gives

$$a_*(G) = \sigma(-S_{\nu,\chi}(G) + c_r(\nu)).$$

Equivalently, the proximal first-order condition gives

$$S_{\nu,\chi}(G) = \nu G + \chi a_*(G).$$

Thus the dual optimizer is the limiting counterpart of the finite-dimensional coefficients $a_i = \sigma(-m_i)$.