

Breast Cancer Statistics, 2015: Convergence of Incidence Rates Between Black and White Women

Carol E. DeSantis, MPH^{1*}; Stacey A. Fedewa, MPH²; Ann Goding Sauer, MPSH³; Joan L. Kramer, MD⁴; Robert A. Smith, PhD⁵; Ahmedin Jemal, DVM, PhD⁶

In this article, the American Cancer Society provides an overview of female breast cancer statistics in the United States, including data on incidence, mortality, survival, and screening. Approximately 231,840 new cases of invasive breast cancer and 40,290 breast cancer deaths are expected to occur among US women in 2015. Breast cancer incidence rates increased among non-Hispanic black (black) and Asian/Pacific Islander women and were stable among non-Hispanic white (white), Hispanic, and American Indian/Alaska Native women from 2008 to 2012. Although white women have historically had higher incidence rates than black women, in 2012, the rates converged. Notably, during 2008 through 2012, incidence rates were significantly higher in black women compared with white women in 7 states, primarily located in the South. From 1989 to 2012, breast cancer death rates decreased by 36%, which translates to 249,000 breast cancer deaths averted in the United States over this period. This decrease in death rates was evident in all racial/ethnic groups except American Indians/Alaska Natives. However, the mortality disparity between black and white women nationwide has continued to widen; and, by 2012, death rates were 42% higher in black women than in white women. During 2003 through 2012, breast cancer death rates declined for white women in all 50 states; but, for black women, declines occurred in 27 of 30 states that had sufficient data to analyze trends. In 3 states (Mississippi, Oklahoma, and Wisconsin), breast cancer death rates in black women were stable during 2003 through 2012. Widening racial disparities in breast cancer mortality are likely to continue, at least in the short term, in view of the increasing trends in breast cancer incidence rates in black women. *CA Cancer J Clin* 2016;66:31-42. © 2015 American Cancer Society.

Keywords: breast neoplasms, epidemiology, screening and early detection

Introduction

Excluding skin cancers, breast cancer is the most common cancer diagnosed among US women, accounting for nearly one in three cancers. It is also the second leading cause of cancer death among women after lung cancer. In this article, we describe trends in breast cancer incidence, mortality, survival, and screening by race/ethnicity in the United States as well as state variations in these measures. Additional data are available from the biennial publication of *Breast Cancer Facts & Figures* (available at cancer.org/statistics).

Materials and Methods

Data Sources

Data on incidence trends, probabilities of developing cancer, and cause-specific survival were obtained from the Surveillance, Epidemiology, and End Results (SEER) program of the National Cancer Institute.¹⁻⁴ The SEER program has been collecting clinical, pathological, and demographic information on cancer patients since 1973. Data are available for whites, blacks, and all races combined since 1973 and for American Indian/Alaska Natives (AI/ANs), Asian/Pacific Islanders (APIs), and Hispanics since 1992. SEER incidence rates were adjusted for reporting delay. Incidence rates by race/ethnicity, stage at diagnosis, and the distribution of breast cancer cases by hormone receptor (HR) and human epidermal growth factor receptor 2 (HER2) statuses were obtained using 2008 through 2012 data from the North American Association of Central Cancer Registries (NAACCR).⁵ NAACCR data were also used for state-level analyses of breast cancer incidence rates and proportions of breast cancers

¹Senior Epidemiologist, Surveillance and Health Services Research, American Cancer Society, Atlanta, GA; ²Director, Risk Factor Screening and Surveillance, Surveillance and Health Services Research, American Cancer Society, Atlanta, GA; ³Epidemiologist, Surveillance and Health Services Research, American Cancer Society, Atlanta, GA; ⁴Assistant Professor of Hematology and Medical Oncology, Emory University School of Medicine, Atlanta, GA; ⁵Vice President, Cancer Screening, Cancer Control Department, American Cancer Society, Atlanta, GA; ⁶Vice President, Surveillance and Health Services Research, American Cancer Society, Atlanta, GA

Corresponding author: Carol E. DeSantis, MPH, Surveillance and Health Services Research, American Cancer Society, 250 Williams Street NW, Atlanta, GA 30303; carol.desantis@cancer.org

DISCLOSURES: The authors report no conflicts of interest.

doi: 10.3322/caac.21320. Available online at cacancerjournal.com

diagnosed at in situ and regional/distant stages.⁵⁻⁷ Overall US rates based on data from NAACCR include all states except Arkansas, Minnesota, and Nevada, because these states failed to meet NAACCR high-quality standards for 1 or more years during 2008 through 2012. For analyses by HR and HER2 status, we limited our analyses to cases diagnosed during 2012, because HER2 status was not required by registries until 2010 and has become increasingly complete since that time. Thus, we excluded 15 additional states (Alabama, Arizona, Florida, Kansas, Louisiana, Maryland, New Jersey, New Mexico, New York, Oklahoma, South Carolina, Tennessee, Texas, West Virginia, and Wyoming) and Washington, DC because greater than 10% of cases were missing HER2 status.

Mortality data were obtained from the SEER program's SEER*Stat database as provided by the National Center for Health Statistics.⁸ Data beginning in 1969 are available for whites and blacks. Since 1990, data are available for the 5 major racial and ethnic groups: non-Hispanic whites, non-Hispanic blacks, APIs, AI/ANs, and Hispanics. Population data were obtained from the US Census Bureau.

Prevalence data on mammography by race/ethnicity and state were obtained from the 2012 Behavioral Risk Factor Surveillance System, an ongoing system of surveys conducted by the state health departments in cooperation with the Centers for Disease Control and Prevention.⁹ Mammography prevalence estimates do not distinguish between examinations for screening and diagnosis.

Statistical Analyses

Estimates of the total number of invasive and in situ breast cancer cases and breast cancer deaths for 2015 were published previously.¹⁰ We calculated the estimated number of breast cancer cases by age at diagnosis by applying the proportion of cases diagnosed in each age group during 2008 through 2012 from the NAACCR analytic file to the total number of estimated cases of invasive and in situ breast cancer in 2015. Similarly, we calculated the estimated number of breast cancer deaths by age at death by applying the proportion of deaths that occurred in each age group during 2008 through 2012 to the total estimated breast cancer deaths in 2015.

The estimated number of female breast cancer deaths averted because of the reduction in breast cancer death rates was calculated by first estimating the number of cancer deaths that would have occurred if the death rate had remained at its 1989 level. The expected number of deaths was estimated by applying the 5-year age-specific cancer death rates in 1989 to the corresponding age-specific female populations from 1990 through 2012. The total number of breast cancer deaths averted was the sum of the difference between the expected number and recorded number of cancer deaths in each age group and calendar year.

TABLE 1. Estimated New Female Breast Cancer Cases and Deaths by Age, United States, 2015*

AGES	IN SITU CASES		INVASIVE CASES		DEATHS	
	NUMBER	%	NUMBER	%	NUMBER	%
<40	1,650	3%	10,500	5%	1,010	3%
40-49	12,310	20%	35,850	15%	3,690	9%
50-59	16,970	28%	54,060	23%	7,600	19%
60-69	15,850	26%	59,990	26%	9,090	23%
70-79	9,650	16%	42,480	18%	8,040	20%
80+	3,860	6%	28,960	12%	10,860	27%
All ages	60,290		231,840		40,290	

*Rounded to the nearest 10 cases. Percentages may not sum to 100% due to rounding.

We examined incidence trends by race/ethnicity, age, and stage and mortality trends by state using the National Cancer Institute's Joinpoint regression analysis program (version 4.2.0.2).¹¹ The direction and magnitude of the resulting trends are described by the annual percent change (APC). In describing trends, the terms increase or decrease were used when the APC was statistically significant; otherwise the term stable was used. The relation between state-level mammography screening rates in 2012 and the percentage of breast cancer cases diagnosed at in situ and late stages during 2008 through 2012 was examined by using the Pearson correlation coefficient. We also calculated rate ratios to compare the incidence rates between black women and white women by state. Probabilities of developing breast cancer were calculated using the DevCan (version 6.7.3) probability of developing cancer software program developed by the National Cancer Institute.¹²

SELECTED FINDINGS

Expected Numbers of New Cases and Deaths by Age

Table 1 shows the estimated number of female breast cancer cases and deaths that are expected to occur in the

TABLE 2. Age-Specific Probabilities of Developing Invasive Female Breast Cancer in the United States*

IF CURRENT AGE IS...	THE PROBABILITY OF DEVELOPING BREAST CANCER IN THE NEXT 10 YEARS IS:	OR 1 IN:
20	0.06%	1,674
30	0.44%	225
40	1.44%	69
50	2.28%	44
60	3.46%	29
70	3.89%	26
Lifetime risk	12.32%	8

*Among those free of cancer at beginning of age interval. Based on cases diagnosed 2010-2012. Percentages and "1 in" numbers may not be numerically equivalent due to rounding.

Probabilities derived using NCI DevCan Software, Version 6.7.3.¹²




FIGURE 1. Breast Cancer Incidence and Mortality Rates by Race/Ethnicity, United States, 2008 to 2012.

Rates are per 100,000 females and age adjusted to the 2000 US standard population.

†Rates are based on data from Contract Health Service Delivery Area counties. NHW indicates non-Hispanic white; NHB, non-Hispanic black; AI/AN, Alaska Native/American Indian; API, Asian/Pacific Islander. Sources: Incidence: North American Association of Central Cancer Registries CiNA Analytic File, 1995 to 2012.⁵ Mortality: National Center for Health Statistics, Centers for Disease Control and Prevention, as provided by the Surveillance, Epidemiology, and End Results program, National Cancer Institute, 2015.⁸

United States in 2015 by age. Approximately 231,840 new cases of invasive breast cancer and 40,290 deaths are expected among US women in 2015. In addition to invasive breast cancers, about 60,290 new diagnoses of in situ breast cancer are expected among US women in 2015. The median age at diagnosis for female breast cancer is 61 years.⁴ The median age at diagnosis is younger for black women (58 years) than for white women (62 years). The median age at breast cancer death is 68 years overall: 69 years for white women and 62 years for black women.⁴

Probability of Developing Invasive Female Breast Cancer

About 12% of women in the US (or 1 in 8) will be diagnosed with breast cancer in their lifetime (Table 2). This lifetime risk represents an average of the risks of different women, rather than the risk of any one woman. Lifetime risk includes the possibility that women will die from other causes before being diagnosed with breast cancer and is often misinterpreted to apply only to women who live to very old ages. Age-specific probabilities for developing cancer over a 10-year period are also provided in Table 2. For example, the risk for a cancer-free woman aged 40 years of being diagnosed with breast cancer over the next 10 years is 1.4%. Equivalently, 1 in 69 women who are aged 40 years will be diagnosed with breast cancer by the age of 50 years.

Cancer Occurrence in the Most Recent Time Period (2008-2012)

Incidence and mortality rates

Female breast cancer incidence and mortality rates vary substantially by race/ethnicity (Fig. 1). Non-Hispanic white (white) and non-Hispanic black (black) women have higher breast cancer incidence and death rates than women of other race/ethnicities; API women have the lowest incidence and death rates. Although the overall breast cancer incidence rate is slightly lower in black women than in white women, the breast cancer death rate is 42% higher in blacks than in whites. This mortality difference likely reflects a combination of biologic and nonbiologic factors, including differences in stage at diagnosis, obesity and comorbidities, tumor characteristics, as well as access, adherence, and response to treatments.¹³⁻¹⁶ Lower breast cancer rates in AI/AN, Hispanic, and API women are thought to largely reflect variation in the prevalence of breast cancer risk factors.¹⁷ For example, Hispanic women tend to have a greater number of children and AI/AN women have their first child at younger ages compared to women of other race/ethnicities, both of which are protective against breast cancer.¹⁸ Conversely, API women are more likely to breastfeed for at least 12 months, less likely to consume alcohol, and have lower rates of obesity, which are generally associated with lower breast cancer risk.¹⁹⁻²¹

Distribution of breast cancer subtypes

Although often referred to as a single disease, breast cancer is distinguished by up to 21 distinct histologic subtypes and at least 4 different molecular subtypes, which




FIGURE 2. Distribution of Breast Cancer Subtypes by Race/Ethnicity, United States, 2012.

NH indicates non-Hispanic; API, Asian/Pacific Islander; HR, hormone receptor; HER2, human epidermal growth factor receptor 2; TNBC, triple-negative breast cancer. Source: North American Association of Central Cancer Registries CiNA Analytic File, 1995 to 2012.⁵

are associated with distinct risk factors and are biologically variable in presentation, response to treatment, and outcomes.^{22–27} Gene expression profiling techniques have allowed researchers to better understand the genetic variability among tumors; however, use of these techniques is currently not standard clinical practice. More convenient approximations of molecular subtypes have been identified using routinely evaluated biological markers, including the presence or absence of hormone (estrogen or progesterone) receptors (HR-positive/HR-negative [HR+/HR–]) and overexpression of the HER2 (HER2+/HER2–) protein.^{28,29} The clinical approximations of molecular subtypes appear to be better for some subtypes than for others.³⁰

Figure 2 shows the distribution of US breast cancer cases by subtype (as defined by HR and HER2 statuses) and race/ethnicity in 2012. Overall, 74% of breast cancer cases were

HR+/HER2– (luminal A), 12% were HR–/HER2– (triple negative), 10% were HR+/HER2+ (luminal B), and 4% were HR–/HER2+ (HER2-enriched), with the distributions varying substantially by race/ethnicity. Black women have the smallest proportion of HR+/HER2– breast cancers and the largest proportion of HR–/HER2– breast cancers compared with women of other race/ethnicities. Triple-negative breast cancers are considered to be more aggressive and have poorer prognoses, in part because there are currently no targeted therapies for these tumors.^{31,32} In contrast, white women have the highest proportion of HR+/HER2– breast cancers, whereas API women have the largest proportion of HR–/HER2+ breast cancers. HR+/HER2– (luminal A) breast cancers are associated with higher survival rates, particularly in the first 5 years after diagnosis, in part because expression of HRs is predictive of response to hormonal therapy.^{28,31}




FIGURE 3. Breast Cancer Stage Distribution and 5-Year Cause-Specific Survival by Race/Ethnicity.

Cause-specific survival rates represent the probability of not dying from breast cancer within 5 years of diagnosis. Patients were diagnosed from 2005 through 2011 and were followed through 2012. †Rates are based on data from Contract Health Service Delivery Area counties. NH indicates non-Hispanic; API, Asian/Pacific Islander; AI/AN, American Indian/Alaska Native. Sources: Stage distribution: North American Association of Central Cancer Registries CINA Analytic File, 1995 to 2012⁵; Survival: Surveillance, Epidemiology, and End Results program, National Cancer Institute, 2015.¹

Stage distribution and survival

Distribution of breast cancer cases by stage at diagnosis and 5-year cause-specific survival by race/ethnicity are shown in Figure 3. White and API women have the highest proportions of localized breast cancers (range, 63%-64%) and the smallest proportions of regional (range, 28%-30%) and distant (5%) stage disease. In contrast, black women have the smallest proportion of localized breast cancers (53%) and the largest proportions of regional (35%) and distant (8%) stage disease.

Cause-specific survival is used instead of relative survival to describe survival in racial and ethnic minorities, because estimates of life expectancy are not available for all racial/ethnic groups. Cause-specific survival is the probability of not dying of breast cancer within a specified number of years after diagnosis. Five-year breast cancer survival is highest for API women both overall and for each known stage, whereas black women have the lowest survival for each known stage at diagnosis. Racial differences in survival in part reflect differences in the distribution of breast cancer subtypes (Fig. 2). A study examining the risk of death among California breast cancer patients by tumor subtype and stage at diagnosis found the greatest disparities were for stage II and III HR+/HER2- breast cancers (31%-39% higher risk of death in blacks than whites).³³ Another study found that, among patients diagnosed with small breast tumors (≤ 2.0 cm) during 2004

through 2011, black patients were more likely to present with lymph node metastases than white patients (24% vs 18%, respectively).³⁴ Poverty, lower levels of educational attainment, and a lack of health insurance are also associated with lower breast cancer survival largely because of limited access to care.³⁵⁻³⁹

Long-term incidence trends by age

Trends in incidence rates for female breast cancer by age at diagnosis are presented in Figure 4. Much of the historic increases in breast cancer incidence rates reflect changes in reproductive patterns, such as delayed childbearing and having fewer children, which are recognized risk factors for breast cancer.^{40,41} In the 1980s, breast cancer incidence rates increased rapidly in women aged 40 years and older, largely due to greater use of mammography screening. Subsequently, incidence rates stabilized for women in their 40s but continued to increase through much of the 1990s in women aged 50 years and older. This increasing trend in older women may reflect rising rates of obesity and the use of menopausal hormones, both of which increase the risk of postmenopausal breast cancer, as well as further increases in the prevalence of mammography screening.⁴²

Around the year 2000, incidence rates began to decline in women aged 50 years and older; and, between 2002 and 2003, breast cancer rates decreased sharply, likely because of the decreased use of menopausal hormones after the 2002




FIGURE 4. Incidence and Mortality Rates of Female Breast Cancer by Age, United States, 1975 to 2012.

Rates are per 100,000 females and age adjusted to the 2000 US standard population. Incidence rates were adjusted for reporting delay. Sources: Incidence: Surveillance, Epidemiology, and End Results program.² Mortality: National Center for Health Statistics, Centers for Disease Control and Prevention, as provided by the Surveillance, Epidemiology, and End Results program, National Cancer Institute, 2015.⁸

publication of the Women's Health Initiative randomized trial results linking the use of combination hormone therapy to breast cancer and heart disease.^{43–45} The decline occurred primarily in white women and for HR+ disease.^{44,46} This trend also may reflect small declines in mammography screening since 2000. The percentage of women aged 40 years older who reported having a mammogram within the past 2 years increased from 29% in 1987 to 70% in 2000, declined 3.4 percentage points from 2000 to 2005, and has since been relatively stable.⁴⁷ Similar reversals in breast cancer incidence trends have been observed in other high-income countries.^{48–53}

Most recently, breast cancer incidence rates in the United States were stable for women in their 50s from 2006 to 2012 but increased for women in their 60s (1.0% per year since 2004) and women aged 70 years and older (1.2% per year since 2005). Incidence rates have been stable for women in their 40s since 1986. Among women ages 20 to 39 years, incidence rates increased slightly (0.6% per year) from 1994 to 2012.

Incidence trends by race/ethnicity

During 2008 through 2012 (the most recent 5 years of data available), overall breast cancer incidence rates increased among black women (0.4% per year) and API women (1.5% per year) but did not change significantly among white, Hispanic, or AI/AN women. Notably, breast cancer incidence rates for whites and blacks in the 9 SEER areas

converged in 2012, reflecting the increase in incidence in black women and relatively stable rates in white women (Fig. 5). Incidence rates for AI/AN women are less stable than for other racial and ethnic groups, because high-quality data for this group are only available from limited geographic areas. We previously noted that the increase in incidence rates in black women has been driven by increases in ER+ breast cancers.⁵⁴ This trend in part may reflect the rising rates of obesity in black women, which increased from 39% in 1999 through 2002, to 49% in 2003 through 2006, to 58% in 2009 through 2012.²¹ Obesity is a risk factor for postmenopausal breast cancer, and a recent study reported that a high recent body mass index was associated with an increased risk of ER+ breast cancer in black women (odds ratio, 1.31; 95% confidence interval, 1.02–1.67 for a body mass index of ≥ 35 kg/m² vs < 25 kg/m²).⁵⁵

We further examined breast cancer incidence trends by stage at diagnosis for 4 racial/ethnic groups (Fig. 6, Table 3). In the most recent period, incidence rates for localized breast cancers increased among white (0.9% per year during 2004–2012), black (2.4% per year during 2006–2012), and API (0.8% per year during 1992–2012) women but were stable among Hispanic women. Conversely, rates for regional-stage tumors decreased in white women (–1.3% per year during 1999–2012) and were stable in the most recent period for black and API women. Incidence rates for regional-stage breast cancer also declined for Hispanic




FIGURE 5. Trends in Female Breast Cancer Incidence and Mortality Rates by Race/Ethnicity, United States, 1975 to 2012.

Rates are per 100,000 females, age adjusted to the 2000 US standard population, and incidence rates are adjusted for reporting delay.

†Rates are 3-year moving averages. NH indicates non-Hispanic; API, Asian/Pacific Islander; AI/AN, American Indian/Alaska Native. Sources: Incidence: Surveillance, Epidemiology, and End Results program, National Cancer Institute, 2015.^{2,3} Mortality: National Center for Health Statistics, Centers for Disease Control and Prevention, as provided by the Surveillance, Epidemiology, and End Results program, National Cancer Institute, 2015.⁸ For Hispanics, incidence data do not include cases from the Alaska Native Registry and mortality data exclude New Hampshire and Oklahoma. Data for American Indians/Alaska Natives are based on Contract Health Service Delivery Area (CHSDA) counties.




FIGURE 6. Trends in Female Breast Cancer Incidence Rates by Stage at Diagnosis and Race/Ethnicity, United States, 1992 to 2012.

Rates are per 100,000 females, age adjusted to the 2000 US standard population, and adjusted for reporting delay. NH indicates non-Hispanic. Dots indicate observed, data and lines represent modeled trends based on Joinpoint analyses. Sources: Surveillance, Epidemiology, and End Results program, National Cancer Institute, 2015.³

women during 2000 through 2010 and have since stabilized. The decreasing trends in regional-stage disease among white and Hispanic women may reflect a shift toward earlier stage at diagnosis. Rates of distant-stage tumors increased in white, black, and API women, but not in Hispanic women; however, in all 4 groups, the rates of unstaged tumors declined sharply, which likely reflects more complete staging of advanced tumors.

Long-term mortality trends

Figure 4 also illustrates national trends in mortality rates for breast cancer by age at death. Overall breast cancer mortality rates decreased 36% from 1989 to 2012 after slowly increasing for many years (0.4% annually since 1975).⁴ As a result of this decline, 249,000 breast cancer deaths have been averted in US women from 1990 to 2012 (Fig. 7). Death rates from breast cancer have decreased in all age groups, ranging from a


TABLE 3. Trends in Female Breast Cancer Incidence Rates by Stage at Diagnosis and Race/Ethnicity, United States, 1992 to 2012

	JOINPOINT ANALYSES							
	TREND 1		TREND 2		TREND 3		TREND 4	
	YEARS	APC	YEARS	APC	YEARS	APC	YEARS	APC
Non-Hispanic white								
All stages	1992-1999	1.7*	1999-2004	-2.3*	2004-2012	0.3		
Localized	1992-1999	1.8*	1999-2004	-2.7*	2004-2012	0.9*		
Regional	1992-1996	-0.1	1996-1999	4.5	1999-2012	-1.3*		
Distant	1992-2006	0.2	2006-2012	2.8*				
Unstaged	1992-1997	0.3	1997-2000	-12.3*	2000-2012	-4.9*		
Non-Hispanic black								
All stages	1992-2012	0.4*						
Localized	1992-2006	0.1	2006-2012	2.4*				
Regional	1992-2012	0.2						
Distant	1992-2012	1.4						
Unstaged	1992-2012	-5.9*						
Asian/Pacific Islander								
All stages	1992-1994	-4.6	1994-1997	6.9	1997-2005	-0.7	2005-2012	1.5*
Localized	1992-2012	0.8*						
Regional	1992-2000	3.1*	2000-2012	-0.6				
Distant	1992-2012	1.3*						
Unstaged	1992-2012	-2.4*						
Hispanic								
All stages	1992-2012	0.2						
Localized	1992-2012	0.4*						
Regional	1992-1997	-0.5	1997-2000	5.3	2000-2010	-1.2*	2010-2012	4.0
Distant	1992-2012	0.6						
Unstaged	1992-2012	-4.1*						

APC indicates annual percent change.

*The APC is statistically different from zero.

Source: Surveillance, Epidemiology, and End Results Program.³

**FIGURE 7. Total Number of Female Breast Cancer Deaths Averted From 1990 to 2012.**

The blue line represents the actual number of breast cancer deaths recorded in each year, and the red line represents the number of breast cancer deaths that would have been expected if breast cancer death rates had remained at their peak rate in 1989.

decline of 1.5% per year in women aged 70 years and older to 2.8% per year in women ages 20 to 39 years. Declines in breast cancer mortality rates have been attributed to both improvements in treatment (eg, adjuvant chemotherapy and hormonal therapy in the 1980s and targeted therapies in the 1990s) and early detection.^{56,57}

Mortality trends by race/ethnicity

Trends in breast cancer mortality rates by race/ethnicity are also shown in Figure 5. From 2003 through 2012, breast cancer mortality rates declined annually by 1.8% in whites, 1.5% in Hispanics, 1.4% in blacks, and 1.0% in APIs but remained unchanged among AI/ANs.⁴ A striking divergence in long-term breast cancer mortality trends between black and white women emerged in the early 1980s and has since continued to widen (Fig. 5); and, by 2012, mortality rates were 42% higher in black women than in white women. This mortality difference likely reflects a combination of factors, including differences in incidence rates (at least during the recent time period), stage at diagnosis, tumor characteristics, obesity, and comorbidities as well as access, adherence, and response to state-of-the-art treatments.^{15,16,58} The racial disparity may also reflect differences in the quality of mammography screening and delayed follow-up for abnormal mammography findings.^{59,60}

TABLE 4. State Variation in Female Breast Cancer Incidence and Mortality Rates and Mammography Usage by Race

STATE	NON-HISPANIC WHITE					NON-HISPANIC BLACK					BLACK:WHITE INCIDENCE RATE RATIO
	RECENT MAMMO- GRAM (%), AGES 45+, 2012†	INCIDENCE 2008-2012			MORTALITY RATE 2008-2012*	RECENT MAMMO- GRAM (%), AGES 45+, 2012†	INCIDENCE 2008-2012			MORTALITY RATE 2008-2012*	
		% IN SITU‡	% REGIONAL/ DISTANT‡	INVASIVE OVERALL RATE*			% IN SITU‡	% REGIONAL/ DISTANT‡	INVASIVE OVERALL RATE*		
Alabama	77	18%	34%	117.5	20.4	81	18%	43%	125.9	30.7	1.07§
Alaska	70	22%	35%	126.7	21.2				141.7		1.12
Arizona	74	19%	32%	118.8	20.6		18%	39%	103.2	29.7	0.87§
Arkansas	68			107.7	21.4	74			106.1	31.4	
California	80	18%	33%	140.5	24.2	87	19%	42%	129.1	33.1	0.92§
Colorado	73	20%	33%	127.7	20.1	68	18%	46%	120.2	26.0	0.94
Connecticut	81	24%	30%	141.1	20.6	83	27%	40%	122.1	24.6	0.87§
Delaware	81	24%	31%	126.8	21.9	86	26%	39%	127.5	26.5	1.01
Dist. of Columbia	80	23%	31%	164.4	24.1	86	20%	43%	137.9	34.0	0.84§
Florida	75	18%	32%	120.4	21.3	72	20%	45%	109.7	28.5	0.91§
Georgia	77	20%	33%	125.8	21.2	83	20%	44%	124.1	29.5	0.99
Hawaii	75	19%	31%	138.3	18.9				134.0		0.97
Idaho	68	17%	35%	120.9	21.4						
Illinois	76	20%	34%	133.2	22.9	81	22%	45%	126.8	32.8	0.95§
Indiana	70	18%	34%	119.7	22.3	75	21%	42%	123.7	31.0	1.03
Iowa	78	19%	33%	124.3	20.9		24%	47%	111.6	25.6	0.90
Kansas	77	17%	33%	123.1	21.3	80	17%	45%	131.4	29.4	1.07
Kentucky	73	17%	34%	121.6	22.1	84	20%	39%	133.2	32.7	1.10§
Louisiana	75	18%	35%	121.2	21.9	80	17%	45%	130.0	34.8	1.07§
Maine	82	21%	32%	125.2	19.2						
Maryland	80	20%	32%	133.5	22.1	89	22%	41%	130.2	30.6	0.98
Massachusetts	86	24%	28%	141.7	20.9	88	26%	37%	115.1	23.7	0.81§
Michigan	78	21%	33%	121.2	22.1	81	22%	42%	122.7	33.1	1.01
Minnesota	81			131.2	20.3	75			94.0	21.7	
Mississippi	70	16%	35%	113.9	20.4	70	16%	47%	124.0	33.3	1.09§
Missouri	75	18%	35%	124.6	22.6	84	19%	47%	135.6	33.7	1.09§
Montana	68	19%	35%	123.5	20.3						
Nebraska	72	19%	33%	123.4	19.7	74	18%	42%	134.2	29.0	1.09
Nevada	72			121.2	25.4	78			109.9	29.3	
New Hampshire	82	21%	31%	136.2	20.5						
New Jersey	77	24%	32%	140.7	24.7	85	21%	44%	124.4	32.5	0.88§
New Mexico	70	16%	32%	125.6	22.4			39%	107.2	30.2	0.85
New York	78	25%	31%	138.9	21.8	78	23%	42%	119.2	28.4	0.86§
North Carolina	78	18%	33%	128.9	21.0	80	19%	43%	128.1	28.8	0.99
North Dakota	77	18%	36%	122.9	19.7						
Ohio	76	18%	35%	121.0	23.2	82	20%	44%	121.0	30.9	1.00
Oklahoma	69	17%	35%	116.9	23.0	70	15%	45%	131.3	35.4	1.12§
Oregon	74	19%	32%	130.4	21.5		21%	46%	121.6	28.5	0.93
Pennsylvania	77	20%	34%	129.0	22.6	87	23%	42%	131.1	33.1	1.02
Rhode Island	84	21%	30%	135.8	20.1		19%	35%	104.8	28.1	0.77§
South Carolina	72	19%	33%	125.9	21.1	80	19%	45%	125.1	29.2	0.99
South Dakota	76	20%	35%	128.0	20.9						
Tennessee	76	18%	34%	120.7	21.2	81	18%	44%	126.2	33.9	1.05§
Texas	72	18%	32%	124.2	21.1	78	19%	43%	120.3	33.7	0.97§
Utah	71	18%	38%	115.9	21.7				108.6		0.94
Vermont	79	22%	30%	129.1	18.7						
Virginia	79	22%	32%	127.8	21.6	80	23%	41%	129.8	31.7	1.02
Washington	75	21%	32%	139.6	21.3	87	24%	42%	127.3	25.0	0.91§
West Virginia	75	17%	34%	111.5	22.7	86	14%	42%	115.5	26.6	1.04
Wisconsin	80	20%	34%	126.5	20.7	88	24%	49%	126.0	32.1	1.00
Wyoming	66	15%	35%	113.3	19.9						
United States	76	20%	33%	128.1	21.9	81	20%	43%	124.3	31.0	0.97§
Range	66-86	15%-25%	28%-38%	107.7-164.4	18.7-25.4	68-89	14%-27%	35%-49%	94.0-141.7	21.7-35.0	—

*All rates are per 100,000 and age-adjusted to 2000 US standard population.

†Recent mammogram is defined as having had a mammogram within the past 2 years.

‡The denominators for percent in situ include all breast cancers. The denominators for percent regional/distant include all invasive breast cancers.

§The black:white incidence rate ratio is significantly different from 1.00 ($P < .05$).

||Statistic could not be calculated; for Behavioral Risk Factor Surveillance System estimate of mammography screening, percentage was not calculated if there were fewer than 50 respondents; for incidence and mortality, statistics were not calculated if there were 25 or fewer cases or deaths. Incidence data are not available for Minnesota. Data on stage distribution were not available for Arkansas and Nevada.

Sources: Mammography: Behavioral Risk Factor Surveillance System 2012, Centers for Disease Control and Prevention.⁹ Incidence: North American Association of Central Cancer Registries.⁵ Overall US incidence data do not include data from Arkansas, Minnesota, or Nevada. Mortality: National Center for Health Statistics, Centers for Disease Control and Prevention.⁸

Furthermore, although findings from national surveys indicate that current mammography screening rates are similar or even slightly higher in black women than in white women, studies suggest that these surveys may overestimate mammography rates, and more so for blacks than for whites.^{61–63}

Variation in mammography use, incidence, and mortality by state

State variations in mammography screening prevalence, breast cancer incidence and mortality rates, and the proportion of breast cancers diagnosed at in situ and regional/distant stages are presented in Table 4. In 2012, the prevalence of recent mammography screening within the past 2 years among white women aged 45 years and older ranged from 66% in Wyoming to 86% in Massachusetts. Thirty-four states had sample sizes large enough to estimate the prevalence of mammography screening within the past 2 years in black women aged 45 years and older, which ranged from 68% in Colorado to 89% in Maryland.

Breast cancer incidence rates ranged from 107.7 cases per 100,000 females in Arkansas to 164.4 cases per 100,000 females in Washington, DC among white women and from 94.0 per 100,000 females in Minnesota to 141.7 cases per 100,000 females in Alaska among black women. We examined rate ratios for 42 states and Washington, DC with incidence rates for both blacks and whites. In 7 states (Alabama, Kentucky, Louisiana, Mississippi, Missouri, Oklahoma, and Tennessee), breast cancer incidence rates were significantly higher in black women than in white women, whereas rates were lower for black women in 11 states and Washington, DC. In 24 other states, rates were not significantly different between blacks and whites. Reasons for higher breast cancer incidence rates among blacks than whites in certain states are not clear but may have resulted in part from the decline in incidence rates associated with a reduction in postmenopausal hormone therapy use that was observed in white women but not black women coupled with increasing incidence rates in black women.^{46,54}

When comparing incidence rates among states, it is also important to consider that incidence rates reflect the intensity of screening as well as disease occurrence. The percentage of in situ breast cancers, an indicator of mammography utilization, varied from 15% in Wyoming to 25% in New York among white women and from 14% in West Virginia to 27% in Connecticut among black women. The proportion of regional/distant stage cancers ranged from 28% in Massachusetts to 38% in Utah among white women and from 35% in Rhode Island to 49% in Wisconsin among black women.

We assessed the relationship between mammography screening rates in 2012 and breast cancer stage at diagnosis during 2008 through 2012. State-level mammography screening prevalence was positively correlated with the percentage of breast cancers diagnosed at in situ stage (Pearson correlation

coefficient [r] = 0.65; $P < .001$) and negatively correlated with the percentage of breast cancers diagnosed at late stages ($r = -0.63$; $P < .001$) among white women. Among black women, state-level mammography screening prevalence was also correlated with in situ diagnoses ($r = 0.49$; $P = .003$) but not with late-stage diagnoses ($r = -0.31$; $P = .08$).

Despite generally similar current and historically lower incidence rates, breast cancer death rates are higher in black women than in white women. Breast cancer death rates among white women range from 18.7 in Vermont to 25.4 in Nevada. In contrast, breast cancer death rates among black women range from 21.7 in Minnesota to 35.0 in Oklahoma. We also examined trends in breast cancer mortality rates from 2003 through 2012 by state (data not shown). Death rates decreased for white women in all 50 states and for black women in 27 of 30 states with sufficient data to examine trends. In contrast, breast cancer death rates for black women were stable during 2003 through 2012 in 3 states: Mississippi, Oklahoma, and Wisconsin. Although 88% of black women reported receiving a recent mammogram in Wisconsin, 49% of breast cancers were diagnosed at regional or distant stages during 2008 through 2012 (Table 4).

Conclusion

From 1989 to 2012, breast cancer death rates have decreased by 36%; and, as a result, 249,000 US breast cancer deaths have been averted during this time period. This decrease was evident in all racial/ethnic groups except AI/ANs. Nevertheless, disparities in breast cancer mortality rates between black and white women continue to widen in the United States, with rates 42% higher among blacks in 2012. This pattern is likely to continue—at least in the near future—in view of the increasing trends in breast cancer incidence rates in black women. Breast cancer incidence rates are higher among black women than white women in 7 states, most of which are located in the South. Black women are also disproportionately diagnosed with triple-negative breast cancers.

The World Cancer Research Fund International estimates that one third of breast cancers could be prevented through healthy behaviors, including maintaining a healthy body weight, engaging in regular physical activity, and not drinking alcohol.⁶⁴ There is growing evidence that high levels of fruit and vegetable consumption may reduce the risk of HR-negative breast cancer.⁶⁵ These findings are supported by studies linking lower breast cancer risk to higher blood levels of carotenoids.^{66,67} In addition, the Collaborative Group on Hormonal Factors in Breast Cancer reviewed 47 studies from 30 countries and concluded that the risk of breast cancer was reduced by 4% for every 12 months of breastfeeding.⁶⁸ More recent studies suggest that the protective effect may be stronger for or even limited to triple-negative breast cancers.^{69–75}

Clinicians should encourage their patients to have regular screening mammography. The American Cancer Society has recently updated their breast cancer screening guidelines for average-risk women.⁷⁶ Annual mammography is recommended for women who are ages 45 to 54 years; women age 55 years and older should transition to biennial screening or have the opportunity to continue screening annually, continuing as long as their overall health is good and life expectancy is 10 years or more.

Women between the ages of 40 and 44 should have the opportunity to begin annual mammography screening. It is also important that patients at high risk of breast cancer are identified and offered appropriate screening and follow-up.⁷⁷ In addition, increased efforts are required to ensure that all segments of the population receive consistent follow-up of abnormal results, prompt diagnosis, and the delivery of high-quality treatment for breast cancer. ■

References

1. Surveillance, Epidemiology, and End Results (SEER) Program. SEER*Stat Database: Incidence-SEER 18 Regs Research Data, Nov 2014 Sub (1992-2012) <Katrina/Rita Population Adjustment>-Linked To County Attributes-Total US, 1969-2013 Counties. Bethesda, MD: National Cancer Institute, Division of Cancer Control and Population Sciences, Surveillance Research Program, Surveillance Systems Branch; 2015.
2. Surveillance, Epidemiology, and End Results (SEER) Program. SEER*Stat Database: Incidence-SEER 9 Regs Research Data with Delay-Adjustment, Malignant Only, Nov 2014 Sub (1975-2012) <Katrina/Rita Population Adjustment>-Linked To County Attributes-Total US, 1969-2013 Counties. Bethesda, MD: National Cancer Institute, Division of Cancer Control and Population Sciences, Surveillance Research Program, Surveillance Systems Branch; 2015.
3. Surveillance, Epidemiology, and End Results (SEER) Program. SEER*Stat Database: Incidence-SEER 13 Regs Research Data with Delay-Adjustment, Malignant Only, Nov 2014 Sub (1992-2012) <Katrina/Rita Population Adjustment>-Linked To County Attributes-Total US, 1969-2013 Counties. Bethesda, MD: National Cancer Institute, Division of Cancer Control and Population Sciences, Surveillance Research Program, Surveillance Systems Branch; 2015.
4. Howlader N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2012, based on Nov 2014 SEER data submission. Bethesda, MD: National Cancer Institute; 2015.
5. Surveillance, Epidemiology and End Results (SEER) Program. SEER*Stat Database: North American Association of Central Cancer Registries (NAACCR) Incidence-CiNA Analytic File, 1995-2012, Custom File With County, North American Association of Central Cancer Registries. Bethesda, MD: National Cancer Institute, Division of Cancer Control and Population Sciences, Surveillance Research Program, Cancer Statistics Branch; 2015.
6. Copeland G, Lake A, Firth R, et al. Cancer in North America: 2008-2012. Volume One: Combined Cancer Incidence for the United States, Canada and North America. Springfield, IL: North American Association of Central Cancer Registries; 2015.
7. Copeland G, Lake A, Firth R, et al. Cancer in North America: 2008-2012. Volume Two: Registry-Specific Cancer Incidence for the United States, Canada and North America. Springfield, IL: North American Association of Central Cancer Registries; 2015.
8. Surveillance, Epidemiology, and End Results (SEER) Program. SEER*Stat Database: Mortality-All COD, Aggregated With State, Total US (1969-2012) <Katrina/Rita Population Adjustment>. Bethesda, MD: National Cancer Institute, Division of Cancer Control and Population Sciences, Surveillance Research Program, Surveillance Systems Branch; 2015. Underlying mortality data provided by the National Center for Health Statistics (cdc.gov/nchs).
9. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System Survey Data. Atlanta, GA: Centers for Disease Control and Prevention, US Department of Health and Human Services; 2012.
10. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin*. 2015;65:5-29.
11. Statistical Research and Applications Branch, National Cancer Institute. Joint Regression Program, version 4.2.0.2, June 2015. Bethesda, MD: Statistical Research and Applications Branch, National Cancer Institute; 2015.
12. Statistical Research and Applications Branch, National Cancer Institute. DevCan: Probability of Developing or Dying of Cancer. DevCan software, version 6.7.3. Statistical Research and Applications Branch, National Cancer Institute, April 2015.
13. Danforth DN Jr. Disparities in breast cancer outcomes between Caucasian and African American women: a model for describing the relationship of biological and nonbiological factors [serial online]. *Breast Cancer Res*. 2013;15:208.
14. Silber JH, Rosenbaum PR, Clark AS, et al. Characteristics associated with differences in survival among black and white women with breast cancer. *JAMA*. 2013;310:389-397.
15. Ooi SL, Martinez ME, Li CI. Disparities in breast cancer characteristics and outcomes by race/ethnicity. *Breast Cancer Res Treat*. 2011;127:729-738.
16. Curtis E, Quale C, Haggstrom D, Smith-Bindman R. Racial and ethnic differences in breast cancer survival: how much is explained by screening, tumor severity, biology, treatment, comorbidities, and demographics? *Cancer*. 2008;112:171-180.
17. Chlebowski RT, Chen Z, Anderson GL, et al. Ethnicity and breast cancer: factors influencing differences in incidence and outcome. *J Natl Cancer Inst*. 2005;97:439-448.
18. Martin JA, Hamilton BE, Osterman MJ, Curtin SC, Matthews TJ. Births: final data for 2013. *Natl Vital Stat Rep*. 2015;64:1-65.
19. Centers for Disease Control and Prevention (CDC). Breastfeeding among US Children Born 2002-2012: CDC National Immunization Surveys. cdc.gov/breastfeeding/data/nis_data/index.htm. Accessed August 4, 2015.
20. Blackwell DL, Lucas JW, Clarke TC. Summary health statistics for US adults: National Health Interview Survey, 2012. National Center for Health Statistics. *Vital Health Stat* 10. 2014;(260):1-161.
21. Fedewa SA, Sauer AG, Siegel RL, Jemal A. Prevalence of major risk factors and use of screening tests for cancer in the United States. *Cancer Epidemiol Biomarkers Prev*. 2015;24:637-652.
22. Dieci MV, Orvieto E, Dominici M, Conte P, Guarneri V. Rare breast cancer subtypes: histological, molecular, and clinical peculiarities. *Oncologist*. 2014;19:805-813.
23. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature*. 2012;490:61-70.
24. Perou CM, Sorlie T, Eisen MB, et al. Molecular portraits of human breast tumours. *Nature*. 2000;406:747-752.
25. Tamimi RM, Colditz GA, Hazra A, et al. Traditional breast cancer risk factors in relation to molecular subtypes of breast cancer. *Breast Cancer Res Treat*. 2012;131:159-167.
26. Yang XR, Chang-Claude J, Goode EL, et al. Associations of breast cancer risk factors with tumor subtypes: a pooled analysis from the Breast Cancer Association Consortium studies. *J Natl Cancer Inst*. 2011;103:250-263.
27. Barnard ME, Boeke CE, Tamimi RM. Established breast cancer risk factors and risk of intrinsic tumor subtypes. *Biochim Biophys Acta*. 2015;1856:73-85.
28. Anderson WF, Rosenberg PS, Prat A, Perou CM, Sherman ME. How many etiological subtypes of breast cancer: two, three, four, or more [serial online]? *J Natl Cancer Inst*. 2014;106. pii: dju165.
29. Anderson WF, Rosenberg PS, Katki HA. Tracking and evaluating molecular tumor markers with cancer registry data: HER2 and breast cancer [serial online]. *J Natl Cancer Inst*. 2014;106. pii: dju093.
30. Cheang MC, Martin M, Nielsen TO, et al. Defining breast cancer intrinsic subtypes by quantitative receptor expression. *Oncologist*. 2015;20:474-482.
31. Blows FM, Driver KE, Schmidt MK, et al. Subtyping of breast cancer by immunohistochemistry to investigate a relationship between subtype and short and long term survival: a collaborative analysis of data for 10,159 cases from 12 studies [serial online]. *PLoS Med*. 2010;7:e1000279.

32. Perou CM, Borresen-Dale AL. Systems biology and genomics of breast cancer [serial online]. *Cold Spring Harb Perspect Biol*. 2011;3. pii: a003293.
33. Tao L, Gomez SL, Keegan TH, Kurian AW, Clarke CA. Breast cancer mortality in African-American and non-Hispanic white women by molecular subtype and stage at diagnosis: a population-based study. *Cancer Epidemiol Biomarkers Prev*. 2015;24:1039-1045.
34. Iqbal J, Ginsburg O, Rochon PA, Sun P, Narod SA. Differences in breast cancer stage at diagnosis and cancer-specific survival by race and ethnicity in the United States. *JAMA*. 2015;313:165-173.
35. Parise CA, Caggiano V. Disparities in race/ethnicity and socioeconomic status: risk of mortality of breast cancer patients in the California Cancer Registry, 2000-2010 [serial online]. *BMC Cancer*. 2013;13:449.
36. Shi R, Taylor H, McLarty J, Liu L, Mills G, Burton G. Effects of payer status on breast cancer survival: a retrospective study [serial online]. *BMC Cancer*. 2015;15:211.
37. Sprague BL, Trentham-Dietz A, Gangnon RE, et al. Socioeconomic status and survival after an invasive breast cancer diagnosis. *Cancer*. 2011;117:1542-1551.
38. Halpern MT, Bian J, Ward EM, Schrag NM, Chen AY. Insurance status and stage of cancer at diagnosis among women with breast cancer. *Cancer*. 2007;110:403-411.
39. Harper S, Lynch J, Meersman SC, Breen N, Davis WW, Reichman MC. Trends in area-socioeconomic and race-ethnic disparities in breast cancer incidence, stage at diagnosis, screening, mortality, and survival among women ages 50 years and over (1987-2005). *Cancer Epidemiol Biomarkers Prev*. 2009;18:121-131.
40. Sullivan PD, Christine B, Connelly R, Barrett H. Analysis of trends in age-adjusted incidence rates for 10 major sites of cancer. *Am J Public Health*. 1972;62:1065-1071.
41. Devesa SS, Silverman DT, Young JL Jr, et al. Cancer incidence and mortality trends among whites in the United States, 1947-84. *J Natl Cancer Inst*. 1987;79:701-770.
42. Toriola AT, Colditz GA. Trends in breast cancer incidence and mortality in the United States: implications for prevention. *Breast Cancer Res Treat*. 2013;138:665-673.
43. Rossouw JE, Anderson GL, Prentice RL, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *JAMA*. 2002;288:321-333.
44. Ravdin PM, Cronin KA, Howlander N, et al. The decrease in breast cancer incidence in 2003 in the United States. *N Engl J Med*. 2007;356:1670-1674.
45. Coombs NJ, Cronin KA, Taylor RJ, Freedman AN, Boyages J. The impact of changes in hormone therapy on breast cancer incidence in the US population. *Cancer Causes Control*. 2010;21:83-90.
46. DeSantis C, Howlander N, Cronin KA, Jemal A. Breast cancer incidence rates in US women are no longer declining. *Cancer Epidemiol Biomarkers Prev*. 2011;20:733-739.
47. Breen N, Gentleman JF, Schiller JS. Update on mammography trends: comparisons of rates in 2000, 2005, and 2008. *Cancer*. 2011;117:2209-2218.
48. Parkin DM. Is the recent fall in incidence of post-menopausal breast cancer in UK related to changes in use of hormone replacement therapy? *Eur J Cancer*. 2009;45:1649-1653.
49. Antoine C, Ameye L, Paesmans M, Rozenberg S. Update of the evolution of breast cancer incidence in relation to hormone replacement therapy use in Belgium. *Maturitas*. 2012;72:317-323.
50. Hemminki E, Kyyronen P, Pukkala E. Post-menopausal hormone drugs and breast and colon cancer: Nordic countries 1995-2005. *Maturitas*. 2008;61:299-304.
51. Canfell K, Banks E, Moa AM, Beral V. Decrease in breast cancer incidence following a rapid fall in use of hormone replacement therapy in Australia. *Med J Aust*. 2008;188:641-644.
52. De P, Neutel CI, Olivetto I, Morrison H. Breast cancer incidence and hormone replacement therapy in Canada. *J Natl Cancer Inst*. 2010;102:1489-1495.
53. Zbuk K, Anand SS. Declining incidence of breast cancer after decreased use of hormone-replacement therapy: magnitude and time lags in different countries. *J Epidemiol Community Health*. 2012;66:1-7.
54. DeSantis C, Ma J, Bryan L, Jemal A. Breast cancer statistics, 2013. *CA Cancer J Clin*. 2014;64:52-62.
55. Bandera EV, Chandran U, Hong CC, et al. Obesity, body fat distribution, and risk of breast cancer subtypes in African American women participating in the AMBER Consortium. *Breast Cancer Res Treat*. 2015;150:655-666.
56. Berry DA, Cronin KA, Plevritis SK, et al. Effect of screening and adjuvant therapy on mortality from breast cancer. *N Engl J Med*. 2005;353:1784-1792.
57. Munoz D, Near AM, van Ravestein NT, et al. Effects of screening and systemic adjuvant therapy on ER-specific US breast cancer mortality [serial online]. *J Natl Cancer Inst*. 2014;106. pii: dju289.
58. Daly B, Olopade OI. A perfect storm: How tumor biology, genomics, and health care delivery patterns collide to create a racial survival disparity in breast cancer and proposed interventions for change. *CA Cancer J Clin*. 2015;65:221-238.
59. Goldman LE, Walker R, Miglioretti DL, Smith-Bindman R, Kerlikowske K; National Cancer Institute Breast Cancer Surveillance Consortium. Accuracy of diagnostic mammography at facilities serving vulnerable women. *Med Care*. 2011;49:67-75.
60. Gorin SS, Heck JE, Cheng B, Smith SJ. Delays in breast cancer diagnosis and treatment by racial/ethnic group. *Arch Intern Med*. 2006;166:2244-2252.
61. Allgood KL, Rauscher GH, Whitman S, Vasquez-Jones G, Shah AM. Validating self-reported mammography use in vulnerable communities: findings and recommendations. *Cancer Epidemiol Biomarkers Prev*. 2014;23:1649-1658.
62. Njai R, Siegel PZ, Miller JW, Liao Y. Misclassification of survey responses and black-white disparity in mammography use, Behavioral Risk Factor Surveillance System, 1995-2006 [serial online]. *Prev Chronic Dis*. 2011;8. pii: A59.
63. Cronin KA, Miglioretti DL, Krapcho M, et al. Bias associated with self-report of prior screening mammography. *Cancer Epidemiol Biomarkers Prev*. 2009;18:1699-1705.
64. World Cancer Research Fund/American Institute for Cancer Research. Continuous Update Project Report: Cancer preventability estimates for diet, nutrition, body fatness, and physical activity. Washington, DC: American Institute for Cancer Research; 2015.
65. Jung S, Spiegelman D, Baglietto L, et al. Fruit and vegetable intake and risk of breast cancer by hormone receptor status. *J Natl Cancer Inst*. 2013;105:219-236.
66. Eliassen AH, Liao X, Rosner B, Tamimi RM, Tworoger SS, Hankinson SE. Plasma carotenoids and risk of breast cancer over 20 y of follow-up. *Am J Clin Nutr*. 2015;101:1197-1205.
67. Wang Y, Gapstur SM, Gaudet MM, Furtado JD, Campos H, McCullough ML. Plasma carotenoids and breast cancer risk in the Cancer Prevention Study II Nutrition Cohort. *Cancer Causes Control*. 2015;26:1233-1244.
68. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and breast-feeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50,302 women with breast cancer and 96,973 women without the disease. *Lancet*. 2002;360:187-195.
69. Palmer JR, Viscidi E, Troester MA, et al. Parity, lactation, and breast cancer subtypes in African American women: results from the AMBER Consortium [serial online]. *J Natl Cancer Inst*. 2014;106. pii: dju237.
70. Millikan RC, Newman B, Tse CK, et al. Epidemiology of basal-like breast cancer. *Breast Cancer Res Treat*. 2008;109:123-139.
71. Ambrosone CB, Zirpoli G, Ruzsyczky M, et al. Parity and breastfeeding among African-American women: differential effects on breast cancer risk by estrogen receptor status in the Women's Circle of Health Study. *Cancer Causes Control*. 2014;25:259-265.
72. Faupel-Badger JM, Arcaro KF, Balkam JJ, et al. Postpartum remodeling, lactation, and breast cancer risk: summary of a National Cancer Institute-sponsored workshop. *J Natl Cancer Inst*. 2013;105:166-174.
73. Li CI, Beaber EF, Tang MT, Porter PL, Daling JR, Malone KE. Reproductive factors and risk of estrogen receptor positive, triple-negative, and HER2-neu overexpressing breast cancer among women 20-44 years of age. *Breast Cancer Res Treat*. 2013;137:579-587.
74. Gaudet MM, Press MF, Haile RW, et al. Risk factors by molecular subtypes of breast cancer across a population-based study of women 56 years or younger. *Breast Cancer Res Treat*. 2011;130:587-597.
75. Phipps AI, Malone KE, Porter PL, Daling JR, Li CI. Reproductive and hormonal risk factors for postmenopausal luminal, HER2-overexpressing, and triple-negative breast cancer. *Cancer*. 2008;113:1521-1526.
76. Oeffinger KC, Fontham ET, Etzioni R, et al. Breast Cancer Screening for Women at Average Risk: 2015 Guideline Update from the American Cancer Society. *JAMA*. In press.
77. Saslow D, Boetes C, Burke W, et al. American Cancer Society guidelines for breast screening with MRI as an adjunct to mammography. *CA Cancer J Clin*. 2007;57:75-89.