031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_negative_max_area_40C
Pearson r=-0.346 (p=0.0569) | Spearman ρ=-0.308 (p=0.0919) | Kendall τ=-0.223 (p=0.0875)
Pearson log1p(activity) r=-0.304 (p=0.0967)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.57, p=0.105)

Global statistics\nn = 31\nR² = 0.119\nRMSE = 69.36 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.65 (p=0.115)
  Spearman ρ=0.48 (p=0.274)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.64 (p=0.048)
  Spearman ρ=-0.64 (p=0.0444)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.57 (p=0.105)
  Spearman ρ=-0.68 (p=0.046)