
031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_positive_max_area_40C
Pearson r=0.228 (p=0.218) | Spearman ρ=-0.196 (p=0.291) | Kendall τ=-0.149 (p=0.252)
Pearson log1p(activity) r=0.037 (p=0.844)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
20
30
40
50
60
70
NylCa
NylCk
NylCp2_TS
NylCp2
UMGSP2
UMGSP1
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
HotPETase
TtA
NfpolyA
ThcCut1
FsC
Tfcut2
LCC_ICCG
ThcCut2
HiC
ph_NylC
AoC
MsmegA
Cut190
MGS0156
Trypsin
Subtilisin
NylB
Papain
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG3 Cutinase / PETase-like regression (r=0.06, p=0.88)
Global statistics\nn = 31\nR² = 0.052\nRMSE = 71.97 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.52 (p=0.232)
Spearman ρ=0.25 (p=0.596)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.07 (p=0.84)
Spearman ρ=0.04 (p=0.92)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.06 (p=0.88)
Spearman ρ=0.26 (p=0.5)