031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_positive_total_area_40C
Pearson r=0.600 (p=3.65e-04) | Spearman ρ=0.527 (p=0.00229) | Kendall τ=0.385 (p=0.00284)
Pearson log1p(activity) r=0.534 (p=0.00198)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.39, p=0.305)

Global statistics\nn = 31\nR² = 0.359\nRMSE = 59.16 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.16 (p=0.737)
  Spearman ρ=0.05 (p=0.908)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.14 (p=0.691)
  Spearman ρ=-0.15 (p=0.674)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.39 (p=0.305)
  Spearman ρ=0.20 (p=0.6)