050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_global_score_40C
Pearson r=0.067 (p=0.719) | Spearman ρ=0.140 (p=0.453) | Kendall τ=0.077 (p=0.551)
Pearson log1p(activity) r=0.097 (p=0.602)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.15, p=0.705)

Global statistics\nn = 31\nR² = 0.005\nRMSE = 73.74 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=0.01 (p=0.983)
Spearman ρ=-0.07 (p=0.878)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.16 (p=0.656)
Spearman ρ=-0.09 (p=0.802)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.15 (p=0.705)
Spearman ρ=0.13 (p=0.746)