050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_positive_max_area_40C
Pearson r=-0.202 (p=0.276) | Spearman ρ=-0.157 (p=0.398) | Kendall τ=-0.096 (p=0.462)
Pearson log1p(activity) r=-0.148 (p=0.426)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.09, p=0.818)

Global statistics\nn = 31\nR² = 0.041\nRMSE = 72.39 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.03 (p=0.95)
  Spearman ρ=-0.14 (p=0.758)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.30 (p=0.407)
  Spearman ρ=0.28 (p=0.437)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.09 (p=0.818)
  Spearman ρ=0.35 (p=0.353)