078 | orientation selected from PEARSON global_score
PEARSON_global_score_40C
Pearson r=-0.511 (p=0.00329) | Spearman ρ=-0.509 (p=0.00344) | Kendall τ=-0.344 (p=0.00745)
Pearson log1p(activity) r=-0.382 (p=0.0341)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.49, p=0.176)

Global statistics\nn = 31\nR² = 0.261\nRMSE = 63.52 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.73 (p=0.0619)
  Spearman ρ=-0.92 (p=0.00344)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.47 (p=0.175)
  Spearman ρ=0.57 (p=0.0832)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.49 (p=0.176)
  Spearman ρ=0.41 (p=0.279)