078 | orientation selected from PEARSON global_score
PEARSON_negative_max_area_40C
Pearson r=-0.286 (p=0.119) | Spearman ρ=-0.209 (p=0.259) | Kendall τ=-0.146 (p=0.271)
Pearson log1p(activity) r=-0.180 (p=0.333)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=0.48, p=0.191)

Global statistics\nn = 31\nR² = 0.082\nRMSE = 70.83 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.06 (p=0.898)
  Spearman ρ=0.01 (p=0.985)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.02 (p=0.962)
  Spearman ρ=0.17 (p=0.644)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.48 (p=0.191)
  Spearman ρ=0.29 (p=0.443)