
089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_global_score_40C
Pearson r=0.140 (p=0.452) | Spearman ρ=0.175 (p=0.346) | Kendall τ=0.112 (p=0.385)
Pearson log1p(activity) r=0.251 (p=0.172)
PA6 activity (µM)
SURFMATCH global score
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG3 Cutinase / PETase-like regression (r=0.40, p=0.281)
Global statistics\nn = 31\nR² = 0.020\nRMSE = 73.18 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.04 (p=0.932)
Spearman ρ=-0.23 (p=0.613)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.03 (p=0.94)
Spearman ρ=-0.09 (p=0.802)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.40 (p=0.281)
Spearman ρ=0.34 (p=0.374)
NylCa
NylCp2_TS
NylCk
NylCp2
UMGSP2
UMGSP1
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
NfpolyA
TtA
HotPETase
ThcCut1
FsC
ThcCut2
Tfcut2
LCC_ICCG
ph_NylC
AoC
HiC
Cut190
NylB
Trypsin
MGS0156
Subtilisin
Papain
MsmegA
chymotrypsin