108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_negative_max_area_40C
Pearson r=-0.257 (p=0.163) | Spearman ρ=-0.313 (p=0.0862) | Kendall τ=-0.216 (p=0.103)
Pearson log1p(activity) r=-0.177 (p=0.342)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.14, p=0.722)

Global statistics\nn = 31\nR² = 0.066\nRMSE = 71.43 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=-0.20 (p=0.669)
Spearman ρ=-0.23 (p=0.624)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.05 (p=0.901)
Spearman ρ=-0.48 (p=0.164)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.14 (p=0.722)
Spearman ρ=0.18 (p=0.643)