127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_negative_max_area_40C
Pearson r=-0.349 (p=0.054) | Spearman ρ=-0.326 (p=0.0731) | Kendall τ=-0.253 (p=0.0516)
Pearson log1p(activity) r=-0.288 (p=0.117)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.23, p=0.545)

Global statistics\nn = 31\nR² = 0.122\nRMSE = 69.25 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.05 (p=0.922)
  Spearman ρ=0.12 (p=0.799)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.33 (p=0.349)
  Spearman ρ=0.13 (p=0.71)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.23 (p=0.545)
  Spearman ρ=-0.18 (p=0.647)