127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_positive_max_area_40C
Pearson r=-0.221 (p=0.232) | Spearman ρ=-0.136 (p=0.466) | Kendall τ=-0.098 (p=0.452)
Pearson log1p(activity) r=-0.123 (p=0.508)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG3 Cutinase / PETase-like regression (r=-0.07, p=0.86)

Global statistics\nn = 31\nR² = 0.049\nRMSE = 72.08 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.48 (p=0.28)
  Spearman ρ=-0.33 (p=0.474)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.23 (p=0.53)
  Spearman ρ=0.20 (p=0.582)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.07 (p=0.86)
  Spearman ρ=0.11 (p=0.786)