001 | orientation selected from FORMULA global_score
FORMULA_global_score_40C
Pearson r=-0.522 (p=0.0026) | Spearman ρ=-0.423 (p=0.0177) | Kendall τ=-0.291 (p=0.0234)
Pearson log1p(activity) r=-0.411 (p=0.0217)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression

Global statistics\nn = 31\nR² = 0.272\nRMSE = 63.04 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=-0.09 (p=0.848)
Spearman ρ=-0.05 (p=0.908)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.55 (p=0.101)
Spearman ρ=0.61 (p=0.0612)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.04 (p=0.914)
Spearman ρ=0.08 (p=0.846)