
001 | orientation selected from FORMULA global_score
FORMULA_negative_max_area_40C
Pearson r=-0.356 (p=0.0493) | Spearman ρ=-0.217 (p=0.241) | Kendall τ=-0.194 (p=0.141)
Pearson log1p(activity) r=-0.194 (p=0.295)
PA6 activity (µM)
SURFMATCH score_negative_max_area
0
50
100
150
200
10.0
12.5
15.0
17.5
20.0
22.5
25.0
27.5
NylCp2_TS
NylCa
NylCk
NylCp2
UMGSP1
UMGSP2
GatA
UMGSP3
tt_NylC
ReU
Ooh
nc_NylC
HotPETase
TtA
NTpolyA
ThcCut1
FsC
Tfcut2
LCC_ICCG
ph_NylC
Cut190
ThcCut2
HiC
AoC
NylB
Papain
MsmegA
Subtilisin
Trypsin
MGS0156
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.127\nRMSE = 69.07 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.10 (p=0.839)
Spearman ρ=-0.07 (p=0.877)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.04 (p=0.916)
Spearman ρ=-0.05 (p=0.886)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.12 (p=0.765)
Spearman ρ=-0.04 (p=0.913)