
001 | orientation selected from FORMULA global_score
FORMULA_positive_max_area_40C
Pearson r=-0.190 (p=0.306) | Spearman ρ=-0.262 (p=0.155) | Kendall τ=-0.193 (p=0.137)
Pearson log1p(activity) r=-0.177 (p=0.341)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
20
30
40
50
60
70
NylCk
NylCa
NylCp2
NylCp2_TS
UMGSP2
UMGSP1
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
HotPETase
TfA
NfpolyA
ThcCut1
FsC
LCC_ICCG
ph_NylC
Trypsin
ThcCut2
Tfcut2
HiC
MGS0156
AoC
Cut190
MsmegA
NylB
Subtilisin
Papain
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.036\nRMSE = 72.56 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.19 (p=0.689)
Spearman ρ=-0.31 (p=0.504)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.05 (p=0.892)
Spearman ρ=-0.06 (p=0.866)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.28 (p=0.471)
Spearman ρ=-0.21 (p=0.586)