
012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_global_score_40C
Pearson r=0.035 (p=0.85) | Spearman ρ=0.094 (p=0.614) | Kendall τ=0.068 (p=0.597)
Pearson log1p(activity) r=0.023 (p=0.902)
PA6 activity (µM)
SURFMATCH global score
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.001\nRMSE = 73.86 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.01 (p=0.981)
Spearman ρ=-0.14 (p=0.758)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.48 (p=0.161)
Spearman ρ=0.29 (p=0.422)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.53 (p=0.146)
Spearman ρ=-0.31 (p=0.413)
NylCk
NylCa
NylCp2_TS
NylCp2
UMGSP2
UMGSP1
GatA
UMGSP3
tt_NylC
ReU
Ooh
nc_NylC
HotPETase
NfpolyA
TtA
ThcCut1
FsC
LCC_ICCG
Tfcut2
ph_NylC
Cut190
AoC
ThcCut2
HiC
Trypsin
NylB
MGS0156
chymotrypsin
Papain
Subtilisin
MsmegA