
012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_negative_max_area_40C
Pearson r=-0.325 (p=0.0747) | Spearman ρ=-0.218 (p=0.239) | Kendall τ=-0.167 (p=0.205)
Pearson log1p(activity) r=-0.180 (p=0.332)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.105\nRMSE = 69.90 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.50 (p=0.257)
Spearman ρ=-0.42 (p=0.35)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.40 (p=0.249)
Spearman ρ=-0.30 (p=0.396)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.30 (p=0.428)
Spearman ρ=0.19 (p=0.622)
NylCa
NylCp2_TS
NylCk
NylCp2
UMGSP1
UMGSP2
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
TtA
NfpolyA
FsC
ThcCut1
HotPETase
Tfcut2
LCC_ICCG
MGS0156
HiC
ThcCut2
AoC
MsmegA
Cut190
ph_NylC
NylB
Papain
Trypsin
Subtilisin
chymotrypsin