
012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_positive_max_area_40C
Pearson r=-0.073 (p=0.695) | Spearman ρ=-0.245 (p=0.184) | Kendall τ=-0.177 (p=0.172)
Pearson log1p(activity) r=-0.183 (p=0.325)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
30
40
50
60
70
80
90
NylCa
NylCk
NylCp2_TS
NylCp2
UMGSP1
UMGSP2
GatA
UMGSP3
tt_NylC
ReU
Oon
nc_NylC
HotPETase
TtA
NfpolyA
ThcCut1
FsC
ph_NylC
LCC_ICCG
Tfcut2
ThcCut2
HiC
Cut190
MGS0156
AoC
Trypsin
MsmegA
NylB
Subtilisin
Papain
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.005\nRMSE = 73.71 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.14 (p=0.767)
Spearman ρ=-0.16 (p=0.728)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.40 (p=0.252)
Spearman ρ=0.44 (p=0.201)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.45 (p=0.228)
Spearman ρ=-0.58 (p=0.103)