
050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_global_score_40C
Pearson r=0.067 (p=0.719) | Spearman ρ=0.140 (p=0.453) | Kendall τ=0.077 (p=0.551)
Pearson log1p(activity) r=0.097 (p=0.602)
PA6 activity (µM)
SURFMATCH global score
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.005\nRMSE = 73.74 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.01 (p=0.983)
Spearman ρ=-0.07 (p=0.878)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.16 (p=0.656)
Spearman ρ=-0.09 (p=0.802)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.15 (p=0.705)
Spearman ρ=0.13 (p=0.746)
NylCk
NylCp2_TS
NylCa
NylCp2
UMGSP1
UMGSP2
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
TtA
HotPETase
NfpolyA
ThcCut1
FsC
Tfcut2
LCC_ICCG
AoC
HiC
Trypsin
Papain
Cut190
ph_NylC
NylB
MGS0156
ThcCut2
Subtilisin
Msmeg
chymotrypsin