
069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_global_score_40C
Pearson r=0.005 (p=0.978) | Spearman ρ=-0.194 (p=0.295) | Kendall τ=-0.116 (p=0.366)
Pearson log1p(activity) r=-0.077 (p=0.681)
PA6 activity (µM)
SURFMATCH global score
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.000\nRMSE = 73.91 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.50 (p=0.253)
Spearman ρ=0.45 (p=0.31)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.24 (p=0.496)
Spearman ρ=-0.23 (p=0.519)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.17 (p=0.663)
Spearman ρ=-0.16 (p=0.68)
NylCk
NylCa
NylCp2_TS
NylCp2
UMGSP1
UMGSP2
GatA
tt_NylC
UMGSP3
ReU
Qqh
nc_NylC
HotPETase
TtA
NfpolyA
ThcCut1
FsC
HiC
AoC
LCC_ICCG
NylB
Trypsin
ThcCut2
Tfcut2
Cut190
chymotrypsin
ph_NylC
MsmegA
MGS0156
Papain
Subtilisin
200
150
100
50
0
−0.3
−0.2
−0.1
0.0