
069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_negative_max_area_40C
Pearson r=-0.109 (p=0.56) | Spearman ρ=-0.031 (p=0.867) | Kendall τ=-0.018 (p=0.891)
Pearson log1p(activity) r=0.014 (p=0.939)
PA6 activity (µM)
SURFMATCH score_negative_max_area
0
50
100
150
200
15
20
25
30
35
40
45
50
NylCp2_TS
NylCk
NylCa
NylCp2
UMGSP2
UMGSP1
GatA
UMGSP3
tt_NylC
ReU
Ooh
nc_NylC
TtA
NfpolyA
HotPETase
ThcCut1
FsC
ThcCut2
Tfcut2
ph_NylC
LCC_ICCG
HiC
Cut190
Trypsin
MGS0156
AoC
MsmegA
NylB
Papain
Subtilisin
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.012\nRMSE = 73.47 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.36 (p=0.43)
Spearman ρ=-0.44 (p=0.328)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.49 (p=0.151)
Spearman ρ=0.38 (p=0.273)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.09 (p=0.827)
Spearman ρ=0.16 (p=0.68)