
078 | orientation selected from PEARSON global_score
PEARSON_negative_max_area_40C
Pearson r=-0.286 (p=0.119) | Spearman ρ=-0.209 (p=0.259) | Kendall τ=-0.146 (p=0.271)
Pearson log1p(activity) r=-0.180 (p=0.333)
PA6 activity (µM)
SURFMATCH score_negative_max_area
0
50
100
150
200
6
8
10
12
14
16
NylCp2_TS
NylCa
NylCk
NylCp2
UMGSP2
UMGSP1
GatA
UMGSP3
tt_NylC
ReU
Ooh
nc_NylC
HotPETase
TtA
NfpolyA
ThcCut1
FsC
LCC_ICCG
Tfcut2
HiC
ph_NylC
Cut190
ThcCut2
MGS0156
AoC
NylB
MsmegA
Papain
Subtilisin
Trypsin
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.082\nRMSE = 70.83 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.06 (p=0.898)
Spearman ρ=0.01 (p=0.985)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.02 (p=0.962)
Spearman ρ=0.17 (p=0.644)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.48 (p=0.191)
Spearman ρ=0.29 (p=0.443)