
089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_positive_max_area_40C
Pearson r=-0.119 (p=0.524) | Spearman ρ=-0.235 (p=0.202) | Kendall τ=-0.172 (p=0.183)
Pearson log1p(activity) r=-0.161 (p=0.388)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
30
40
50
60
70
NylCa
NylCk
NylCp2_TS
NylCp2
UMGSP2
UMGSP1
GatA
tt_NylC
UMGSP3
ReU
Ooh_nc_NylC
TtA
NfpolyA
HotPETase
ThcCut1
FsC
LCC_ICCG
AoC
Tfcut2
NylB
ph_NylC
MGS0156
Cut190
ThcCut2
HiC
Trypsin
Subtilisin
Papain
chymotrypsin
MsmegA
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.014\nRMSE = 73.38 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.19 (p=0.677)
Spearman ρ=-0.13 (p=0.788)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.10 (p=0.78)
Spearman ρ=-0.14 (p=0.705)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.02 (p=0.956)
Spearman ρ=-0.08 (p=0.846)