127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_global_score_40C
Pearson r=0.319 (p=0.0808) | Spearman ρ=0.487 (p=0.00549) | Kendall τ=0.361 (p=0.00492)
Pearson log1p(activity) r=0.511 (p=0.0033)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression

Global statistics\nn = 31\nR² = 0.101\nRMSE = 70.06 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
Pearson r=-0.33 (p=0.47)
Spearman ρ=-0.09 (p=0.848)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.11 (p=0.757)
Spearman ρ=0.26 (p=0.464)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.35 (p=0.359)
Spearman ρ=0.53 (p=0.141)