
127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_positive_max_area_40C
Pearson r=-0.221 (p=0.232) | Spearman ρ=-0.136 (p=0.466) | Kendall τ=-0.098 (p=0.452)
Pearson log1p(activity) r=-0.123 (p=0.508)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
10
15
20
25
30
35
40
45
NylCa
NylCp2_TS
NylCp2
NylCk
UMGSP1
UMGSP2
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
TtA
HotPETase
NfpolyA
ThcCut1
FsC
LCC_ICCG
MGS0156
AoC
Tfcut2
ph_NylC
HiC
MsmegA
ThcCut2
Cut190
NylB
Papain
Trypsin
Subtilisin
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.049\nRMSE = 72.08 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.48 (p=0.28)
Spearman ρ=-0.33 (p=0.474)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.23 (p=0.53)
Spearman ρ=0.20 (p=0.582)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.07 (p=0.86)
Spearman ρ=0.11 (p=0.786)