
146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_negative_max_area_40C
Pearson r=-0.215 (p=0.246) | Spearman ρ=-0.075 (p=0.69) | Kendall τ=-0.049 (p=0.707)
Pearson log1p(activity) r=-0.215 (p=0.245)
PA6 activity (µM)
SURFMATCH score_negative_max_area
0
50
100
150
200
10
15
20
25
30
35
40
45
NylCk
NylCa
NylCp2_TS
NylCp2
UMGSP2
UMGSP1
GatA
UMGSP3
tt_NylC
ReU
Ooh
nc_NylC
TtA
NfpolyA
HotPETase
ThcCut1
FsC
ThcCut2
ph_NylC
MsmegA
MGS0156
NylB
Subtilisin
HiC
Tfcut2
Trypsin
Cut190
chymotrypsin
Papain
LCC_ICCG
AoC
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.046\nRMSE = 72.19 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.57 (p=0.178)
Spearman ρ=0.32 (p=0.487)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.64 (p=0.0472)
Spearman ρ=0.88 (p=8.01e-04)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.42 (p=0.263)
Spearman ρ=-0.30 (p=0.44)