146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_negative_total_area_40C
Pearson r=0.479 (p=0.00637) | Spearman ρ=0.470 (p=0.00768) | Kendall τ=0.328 (p=0.011)
Pearson log1p(activity) r=0.446 (p=0.0119)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression

Global statistics\nn = 31\nR² = 0.230\nRMSE = 64.87 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.67 (p=0.102)
  Spearman ρ=-0.52 (p=0.229)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.28 (p=0.442)
  Spearman ρ=-0.21 (p=0.564)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.62 (p=0.0724)
  Spearman ρ=0.45 (p=0.227)