
146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_positive_max_area_40C
Pearson r=-0.137 (p=0.463) | Spearman ρ=-0.250 (p=0.174) | Kendall τ=-0.200 (p=0.124)
Pearson log1p(activity) r=-0.131 (p=0.484)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
10
15
20
25
30
35
40
NylCa
NylCk
NylCp2_TS
NylCp2
UMGSP1
GatA
UMGSP2
UMGSP3
tt_NylC
ReU
Ooh
nc_NylC
HotPETase
TtA
NfpolyA
ThcCut1
FsC
LCC_ICCG
Tfcut2
Cut190
ph_NylC
AoC
Trypsin
ThcCut2
NylB
HiC
MGS0156
Papain
MsmegA
chymotrypsin
Subtilisin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
Global statistics\nn = 31\nR² = 0.019\nRMSE = 73.21 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.18 (p=0.692)
Spearman ρ=-0.13 (p=0.786)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.20 (p=0.585)
Spearman ρ=-0.38 (p=0.28)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.05 (p=0.895)
Spearman ρ=-0.24 (p=0.54)