001 | orientation selected from FORMULA global_score
FORMULA_positive_total_area_40C
Pearson r=0.505 (p=0.00375) | Spearman ρ=0.466 (p=0.00822) | Kendall τ=0.334 (p=0.00952)
Pearson log1p(activity) r=0.466 (p=0.00826)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG1 NylC-like regression (r=-0.23, p=0.624)

Global statistics\nn = 31\nR² = 0.255\nRMSE = 63.78 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.23 (p=0.624)
  Spearman ρ=-0.16 (p=0.726)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.07 (p=0.84)
  Spearman ρ=-0.20 (p=0.581)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.46 (p=0.216)
  Spearman ρ=0.24 (p=0.526)