
050 | orientation selected from FORMULA negative_max_area, threshold=0.50
FORMULA_positive_max_area_40C
Pearson r=-0.202 (p=0.276) | Spearman ρ=-0.157 (p=0.398) | Kendall τ=-0.096 (p=0.462)
Pearson log1p(activity) r=-0.148 (p=0.426)
PA6 activity (µM)
SURFMATCH score_positive_max_area
NylCa
NylCk
NylCp2_TS
NylCp2
UMGSP1
UMGSP2
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
TtA
NfpolyA
HotPETase
ThcCut1
FsC
LCC_ICCG
Tfcut2
NylB
MsmegA
chymotrypsin
Trypsin
Cut190
ThcCut2
Subtilisin
ph_NylC
MGS0156
AoC
HiC
Papain
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.03, p=0.95)
Global statistics\nn = 31\nR² = 0.041\nRMSE = 72.39 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.03 (p=0.95)
Spearman ρ=-0.14 (p=0.758)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.30 (p=0.407)
Spearman ρ=0.28 (p=0.437)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.09 (p=0.818)
Spearman ρ=0.35 (p=0.353)