
069 | orientation selected from FORMULA negative_total_area, threshold=0.50
FORMULA_positive_max_area_40C
Pearson r=-0.158 (p=0.396) | Spearman ρ=-0.199 (p=0.284) | Kendall τ=-0.139 (p=0.289)
Pearson log1p(activity) r=-0.142 (p=0.446)
PA6 activity (µM)
SURFMATCH score_positive_max_area
0
50
100
150
200
10
15
20
25
30
35
40
NylCa
NylCk
NylCp2_TS
NylCp2
UMGSP2
UMGSP1
GatA
UMGSP3
tt_NylC
ReU
Ooh
nc_NylC
HotPETase
TtA
NfpolyA
ThcCut1
FsC
LCC_ICCG
ThcCut2
AoC
Tfcut2
Cut190
ph_NylC
HiC
NylB
MsmegA
chymotrypsin
Trypsin
MGS0156
Papain
Subtilisin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=0.01, p=0.983)
Global statistics\nn = 31\nR² = 0.025\nRMSE = 72.98 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.01 (p=0.983)
Spearman ρ=-0.30 (p=0.513)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.32 (p=0.36)
Spearman ρ=-0.38 (p=0.275)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.01 (p=0.975)
Spearman ρ=-0.16 (p=0.687)