078 | orientation selected from PEARSON global_score
PEARSON_positive_max_area_40C
Pearson r=-0.284 (p=0.121) | Spearman ρ=-0.371 (p=0.0399) | Kendall τ=-0.277 (p=0.0339)
Pearson log1p(activity) r=-0.223 (p=0.227)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG1 NylC-like regression (r=-0.38, p=0.403)

Global statistics\nn = 31\nR² = 0.081\nRMSE = 70.87 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.38 (p=0.403)
  Spearman ρ=-0.28 (p=0.536)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.07 (p=0.85)
  Spearman ρ=-0.02 (p=0.946)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.36 (p=0.347)
  Spearman ρ=-0.41 (p=0.269)