078 | orientation selected from PEARSON global_score
PEARSON_positive_total_area_40C
Pearson r=0.486 (p=0.0056) | Spearman ρ=0.507 (p=0.00358) | Kendall τ=0.349 (p=0.00671)
Pearson log1p(activity) r=0.489 (p=0.00526)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG1 NylC-like regression (r=-0.14, p=0.757)

Global statistics\nn = 31\nR² = 0.236\nRMSE = 64.61 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.14 (p=0.757)
  Spearman ρ=-0.11 (p=0.818)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.22 (p=0.533)
  Spearman ρ=-0.30 (p=0.402)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.48 (p=0.186)
  Spearman ρ=0.36 (p=0.337)