
089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_negative_max_area_40C
Pearson r=-0.214 (p=0.247) | Spearman ρ=-0.203 (p=0.273) | Kendall τ=-0.148 (p=0.258)
Pearson log1p(activity) r=-0.202 (p=0.277)
PA6 activity (µM)
SURFMATCH score_negative_max_area
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=0.19, p=0.687)
Global statistics\nn = 31\nR² = 0.046\nRMSE = 72.20 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.19 (p=0.687)
Spearman ρ=0.06 (p=0.907)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.10 (p=0.787)
Spearman ρ=0.04 (p=0.913)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.31 (p=0.424)
Spearman ρ=-0.31 (p=0.416)
NylCk
NylCa
NylCp2_TS
NylCp2
UMGSP2
UMGSP1
GatA
UMGSP3
tt_NylC
ReU
Ooh
nc_NylC
HotPETase
NfpolyA
TtA
ThcCut1
FsC
Tfcut2
HiC
NylB
ph_NylC
ThcCut2
AoC
LCC_ICCG
Cut190
Trypsin
Subtilisin
Papain
MsmegA
chymotrypsin
MGS0156
0
50
100
150
200
10
15
20
25
30
35