089 | orientation selected from PEARSON positive_max_area, threshold=0.50
PEARSON_negative_total_area_40C
Pearson r=0.495 (p=0.00465) | Spearman ρ=0.444 (p=0.0124) | Kendall τ=0.327 (p=0.0111)
Pearson log1p(activity) r=0.386 (p=0.032)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG1 NylC-like regression (r=-0.17, p=0.721)

Global statistics\nn = 31\nR² = 0.245\nRMSE = 64.22 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.17 (p=0.721)
  Spearman ρ=-0.20 (p=0.67)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.32 (p=0.366)
  Spearman ρ=-0.31 (p=0.385)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.05 (p=0.889)
  Spearman ρ=0.01 (p=0.983)