
108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_negative_max_area_40C
Pearson r=-0.257 (p=0.163) | Spearman ρ=-0.313 (p=0.0862) | Kendall τ=-0.216 (p=0.103)
Pearson log1p(activity) r=-0.177 (p=0.342)
PA6 activity (µM)
SURFMATCH score_negative_max_area
0
50
100
150
200
10
15
20
25
30
35
NylCa
NylCp2_TS
NylCk
NylCp2
UMGSP2
UMGSP1
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
TtA
NfpolyA
HotPETase
ThcCut1
FsC
LCC_ICCG
ph_NylC
Tfcut2
HiC
NylB
Cut190
ThcCut2
MGS0156
AoC
Trypsin
Subtilisin
MsmegA
Papain
chymotrypsin
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.20, p=0.669)
Global statistics\nn = 31\nR² = 0.066\nRMSE = 71.43 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.20 (p=0.669)
Spearman ρ=-0.23 (p=0.624)
SG2 Amidase / urethane-like (n=10)
Pearson r=0.05 (p=0.901)
Spearman ρ=-0.48 (p=0.164)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.14 (p=0.722)
Spearman ρ=0.18 (p=0.643)