108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_negative_total_area_40C
Pearson r=0.595 (p=4.17e-04) | Spearman ρ=0.547 (p=0.00145) | Kendall τ=0.378 (p=0.00334)
Pearson log1p(activity) r=0.505 (p=0.00376)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG1 NylC-like regression (r=0.22, p=0.63)

Global statistics\nn = 31\nR² = 0.354\nRMSE = 59.41 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.22 (p=0.63)
  Spearman ρ=0.30 (p=0.513)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.30 (p=0.395)
  Spearman ρ=-0.29 (p=0.422)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.11 (p=0.776)
  Spearman ρ=0.00 (p=1)