
108 | orientation selected from PEARSON positive_total_area, threshold=0.50
PEARSON_positive_max_area_40C
Pearson r=-0.319 (p=0.08) | Spearman ρ=-0.348 (p=0.0549) | Kendall τ=-0.236 (p=0.0701)
Pearson log1p(activity) r=-0.380 (p=0.0352)
PA6 activity (μM)
SURFMATCH score_positive_max_area
0
50
100
150
200
10
20
30
40
50
NylCk
NylCa
NylCp2_TS
NylCp2
UMGSP2
UMGSP1
GatA
tt_NylC
UMGSP3
ReIJ
Ooh
nc_NylC
TtA
HotPETase
NfpolyA
ThcCut1
FsC
ThcCut2
Tfcut2
LCC_ICCG
MGS0156
AoC
ph_NylC
Cut190
HiC
Trypsin
chymotrypsin
MsmegA
NylB
Subtilisin
Papain
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=-0.40, p=0.378)
Global statistics\nn = 31\nR² = 0.102\nRMSE = 70.04 μM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=-0.40 (p=0.378)
Spearman ρ=-0.48 (p=0.274)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.11 (p=0.768)
Spearman ρ=-0.04 (p=0.906)
SG3 Cutinase / PETase-like (n=9)
Pearson r=-0.05 (p=0.898)
Spearman ρ=0.09 (p=0.812)