127 | orientation selected from PEARSON negative_max_area, threshold=0.50
PEARSON_negative_total_area_40C
Pearson r=0.458 (p=0.00963) | Spearman ρ=0.376 (p=0.0371) | Kendall τ=0.270 (p=0.0359)
Pearson log1p(activity) r=0.361 (p=0.0458)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG1 NylC-like regression (r=-0.03, p=0.954)

Global statistics\nn = 31\nR² = 0.209\nRMSE = 65.72 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.03 (p=0.954)
  Spearman ρ=-0.16 (p=0.728)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.57 (p=0.0869)
  Spearman ρ=-0.63 (p=0.0495)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.23 (p=0.543)
  Spearman ρ=0.18 (p=0.647)