
146 | orientation selected from PEARSON negative_total_area, threshold=0.50
PEARSON_global_score_40C
Pearson r=0.409 (p=0.0225) | Spearman ρ=0.366 (p=0.043) | Kendall τ=0.278 (p=0.0304)
Pearson log1p(activity) r=0.398 (p=0.0265)
PA6 activity (µM)
SURFMATCH global score
Structural groups (D=0.50)
SG1 NylC-like (n=7)
SG10 Subtilisin isolated (n=1)
SG2 Amidase / urethane-like (n=10)
SG3 Cutinase / PETase-like (n=9)
SG4 MGS0156 / LipBC (n=1)
SG6 Trypsin / chymotrypsin (n=2)
SG7 Papain / bromelain (n=1)
Global linear regression
SG1 NylC-like regression (r=0.68, p=0.0918)
Global statistics\nn = 31\nR² = 0.167\nRMSE = 67.46 µM
Within-group correlations (n ≥ 4)
exploratory
SG1 NylC-like (n=7)
Pearson r=0.68 (p=0.0918)
Spearman ρ=0.74 (p=0.0579)
SG2 Amidase / urethane-like (n=10)
Pearson r=-0.12 (p=0.746)
Spearman ρ=-0.07 (p=0.841)
SG3 Cutinase / PETase-like (n=9)
Pearson r=0.39 (p=0.304)
Spearman ρ=0.19 (p=0.632)
NylCa
NylCk
NylCp2_TS
NylCp2
UMGSP1
UMGSP2
GatA
tt_NylC
UMGSP3
ReU
Ooh
nc_NylC
TtA
HotPETase
NfpolyA
ThcCut1
FsC
Tfcut2
LCC_ICCG
HiC
ph_NylC
ThcCut2
Cut190
AoC
MGS0156
MsmegA
Trypsin
Papain
chymotrypsin
NylB
Subtilisin
200
150
100
50
0
−0.16
−0.14
−0.12
−0.10
−0.08
−0.06
−0.04
−0.02