001 | orientation selected from FORMULA global_score
FORMULA_positive_max_area_40C
Pearson r=-0.190 (p=0.306) | Spearman ρ=-0.262 (p=0.155) | Kendall τ=-0.193 (p=0.137)
Pearson log1p(activity) r=-0.177 (p=0.341)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.05, p=0.892)

Global statistics\nn = 31\nR² = 0.036\nRMSE = 72.56 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.19 (p=0.689)
  Spearman ρ=-0.31 (p=0.504)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.05 (p=0.892)
  Spearman ρ=-0.06 (p=0.866)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.28 (p=0.471)
  Spearman ρ=-0.21 (p=0.586)