012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_global_score_40C
Pearson r=0.035 (p=0.85) | Spearman ρ=0.094 (p=0.614) | Kendall τ=0.068 (p=0.597)
Pearson log1p(activity) r=0.023 (p=0.902)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.48, p=0.161)

Global statistics\nn = 31\nR² = 0.001\nRMSE = 73.86 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.01 (p=0.981)
  Spearman ρ=-0.14 (p=0.758)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.48 (p=0.161)
  Spearman ρ=0.29 (p=0.422)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.53 (p=0.146)
  Spearman ρ=-0.31 (p=0.413)