012 | orientation selected from FORMULA positive_max_area, threshold=0.50
FORMULA_negative_max_area_40C
Pearson r=-0.325 (p=0.0747) | Spearman ρ=-0.218 (p=0.239) | Kendall τ=-0.167 (p=0.205)
Pearson log1p(activity) r=-0.180 (p=0.332)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=-0.40, p=0.249)

Global statistics\nn = 31\nR² = 0.105\nRMSE = 69.90 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.50 (p=0.257)
  Spearman ρ=-0.42 (p=0.35)
SG2 Amidase / urethane-like (n=10)
  Pearson r=-0.40 (p=0.249)
  Spearman ρ=-0.30 (p=0.396)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.30 (p=0.428)
  Spearman ρ=0.19 (p=0.622)