031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_global_score_40C
Pearson r=-0.356 (p=0.0491) | Spearman ρ=-0.294 (p=0.108) | Kendall τ=-0.225 (p=0.0791)
Pearson log1p(activity) r=-0.256 (p=0.164)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.25, p=0.486)

Global statistics\nn = 31\nR² = 0.127\nRMSE = 69.06 µM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=-0.29 (p=0.535)
  Spearman ρ=-0.27 (p=0.558)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.25 (p=0.486)
  Spearman ρ=0.16 (p=0.662)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=0.53 (p=0.139)
  Spearman ρ=0.59 (p=0.0939)