031 | orientation selected from FORMULA positive_total_area, threshold=0.50
FORMULA_negative_total_area_40C
Pearson r=0.639 (p=1.08e-04) | Spearman ρ=0.522 (p=0.00257) | Kendall τ=0.372 (p=0.00375)
Pearson log1p(activity) r=0.494 (p=0.00478)



Structural groups (D=0.50)
- SG1 NylC-like (n=7)
- SG10 Subtilisin isolated (n=1)
- SG2 Amidase / urethane-like (n=10)
- SG3 Cutinase / PETase-like (n=9)
- SG4 MGS0156 / LipBC (n=1)
- SG6 Trypsin / chymotrypsin (n=2)
- SG7 Papain / bromelain (n=1)
- Global linear regression
- SG2 Amidase / urethane-like regression (r=0.15, p=0.683)

Global statistics\nn = 31\nR² = 0.409\nRMSE = 56.83 μM

Within-group correlations (n ≥ 4)
exploratory

SG1 NylC-like (n=7)
  Pearson r=0.87 (p=0.0101)
  Spearman ρ=0.95 (p=8.06e-04)
SG2 Amidase / urethane-like (n=10)
  Pearson r=0.15 (p=0.683)
  Spearman ρ=-0.06 (p=0.867)
SG3 Cutinase / PETase-like (n=9)
  Pearson r=-0.49 (p=0.18)
  Spearman ρ=-0.55 (p=0.126)